


ferti·lome[®]



Dollar Weed Control

Dollar Weed Control

Dollar Weed Control

Covers 5,000 Sq. Ft.

- For Postemergence Control Of Annual And Perennial Broadleaf Weeds in Established Home Lawns
 - Apply At Any Time (Spring, Summer, Fall)
- Apply (Anytime) When Weeds Are Actively Growing


ACTIVE INGREDIENT:	
Penoxsulam: 2-(2,2-difluoroethoxy)-N-(5,8-dimethoxy[1,2,4] triazolo [1,5c]pyrimidin-2-yl)-6-(trifluoromethyl) benzenesulfonamide.....	0.04%
OTHER INGREDIENTS:	99.96%
Total	100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

(SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS)

Net Weight 17 LBS. (7.71 KG)



Dollar Weed Control

Dollar Weed Control

NOTICE: Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on the label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.
Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Wear long-sleeved shirt, long pants, and shoes plus socks. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions for use carefully before applying.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in cool dry place in original container.

Container Handling: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-992-5994 for disposal instructions. Never place unused product down any indoor or outdoor drain.

HOW THIS WORKS

ferti•lome® DOLLAR WEED CONTROL provides postemergence control of annual, biennial and perennial broad leaf weeds in established home lawns.

GENERAL USE PRECAUTIONS AND RESTRICTIONS

- Do not apply **ferti•lome® DOLLAR WEED CONTROL** on dichondra, carpetgrass, bahiagrass, or to turfgrass containing desirable clovers.
- Do not apply **ferti•lome® DOLLAR WEED CONTROL** directly to, or otherwise permit **ferti•lome® DOLLAR WEED CONTROL** to come into contact with, flowers, ornamental shrubs or trees, or other desirable broadleaf plants, as serious plant injury may occur.
- Do not apply to centipedegrass during the period of winter dormancy. Grass should be green and actively growing at the time of application.
- The active ingredient in this product may leave residues in treated grass clippings which could be harmful to some plants and trees. Do not collect treated grass clippings for use as mulch around plants. The applicator must notify in writing the appropriate residences/individuals of this precaution when this product is applied by a lawn service/commercial applicator.
- To minimize the potential for turfgrass injury, allow four weeks between applications.
- Avoid unnecessary disturbances, including watering and mowing, for approximately 24 hours after application.
- Use **ferti•lome® DOLLAR WEED CONTROL** on solid (monoculture) stands of perennial ryegrass or tall fescue at a maximum rate of 1.2 lb per 1000 sq ft. Some yellowing of these turfgrasses may occur within two weeks of application. Any discoloration is temporary and will dissipate under adequate growing conditions.
- Apply **ferti•lome® DOLLAR WEED CONTROL** to newly seeded or sprigged turfgrass only after the third mowing or after the turfgrass seedlings have developed an adequate secondary root system. Reseeding is not recommended for at least three to four weeks after application.
- Do not apply more than 3.4 lb of **ferti•lome® DOLLAR WEED CONTROL** per 1000 sq ft per application.
- Do not apply more than 5.1 lb of **ferti•lome® DOLLAR WEED CONTROL** per 1000 sq ft per year.

TREATMENT ON TURFGRASS SPECIES NOT LISTED ON THE LABEL FOR ferti•lome® DOLLAR WEED CONTROL

Users who wish to use **ferti•lome® DOLLAR WEED CONTROL** on turf species not recommended on this label may determine the suitability for use by treating a small area at a recommended rate. Prior to treatment of large areas, the treated area should be observed for any sign of herbicidal injury during 30 days of normal growing conditions to determine if the target turfgrass species is tolerant. Applications should not be made to turfgrass under stress. The user assumes responsibility for any plant damage or other liability resulting from use of **ferti•lome® DOLLAR WEED CONTROL** on turfgrass species not recommended on this label.

ferti•lome® DOLLAR WEED CONTROL is recommended for use on the following established turfgrass species (numbers in parentheses (-) refer to footnotes below):

Established Cool Season Turfgrass(1)	Established Warm Season Turfgrass (3)
Annual bluegrass Bentgrass Bluegrass, Kentucky Fescue, chewing Fescue, creeping red Fescue, sheeps Fescue, tall (2) Ryegrass, perennial (2)	Bermudagrass (common or hybrid) Centipedegrass (4) Fescue, tall (growing in warm season areas) (2) Kikuyugrass St. Augustinegrass Zoysiagrass

- Use no more than 2.3 lbs. of **ferti•lome® DOLLAR WEED CONTROL** per 1,000 sq. ft. per application.
- Use Penoxsulam 0.04% on these species at a maximum rate of 1.2 lb per 1000 sq ft per application and only if some injury can be tolerated. A slight yellowing of turfgrass may occur within two weeks of application. The discoloration is temporary and is generally gone within four weeks of treatment under adequate growing conditions.
- Use no more than 3.4 lbs. of **ferti•lome® DOLLAR WEED CONTROL** per 1,000 sq. ft. per application.
- Do not apply to centipedegrass during the period of winter dormancy. Grass should be green and actively growing at the time of application.

WEEDS CONTROLLED OR SUPPRESSED

Generally, application rates at the lower end of the recommended rate range will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species and under conditions where control is more difficult (plant stress conditions such as drought or extreme temperatures, dense weed stands and/or larger weeds), the higher rate within the rate range or repeat applications will be needed. Weeds growing in the absence of competition from other vegetation generally require higher rates or repeat applications to obtain satisfactory control or suppression.

Weeds Controlled or Suppressed		Rate
		1 lb./1,000 sq. ft.
Chickweed, common Chickweed, mouseear Clover, hop Clover, red Clover, white Dandelion, common Dollarweed English lawn daisy False dandelion Florida betony	Ivy, ground Lespedeza, common Medic, black Parsley-piert Plantain, broadleaf Plantain, buckhorn Virginia buttonweed ¹	3.4
Kyllinga, annual and green (suppression only) ¹	Nutsedge, purple and yellow (suppression only) ¹	2.3 - 3.4

¹ Re-treatment may be necessary

HOW TO APPLY

Apply **ferti•lome® DOLLAR WEED CONTROL** using a drop or rotary-type spreader designed to apply granules. Calibrate application equipment prior to use according to manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first.

To optimize product performance mow the turfgrass two to three days before applying **ferti•lome® DOLLAR WEED CONTROL**. If the turfgrass is under moisture stress, a thorough sprinkler irrigation of at least ½ inch may be needed at least one to two days before application. For optimal results, apply **ferti•lome® DOLLAR WEED CONTROL** in the morning when dew is present. If necessary, sprinkler irrigate with less than ¼ inch of water just prior to application to maximize particle adherence to leaf surfaces. Avoid watering of treated area for 24 hours following application. Avoid making applications if rainfall is expected within 24 hours after treatment.

APPLICATION TECHNIQUES FOR APPLYING ferti•lome® DOLLAR WEED CONTROL

- When using a drop-type spreader, a splash board mounted under the hopper will provide more even granule distribution.
- A chain fastened to the side of the spreader and allowed to drag on the soil can be used to mark the edge of the treated swath and help prevent skips and overlaps.
- For treating smaller areas, a hand held or push-type rotary applicator such as a whirlybird or cyclone unit is recommended. For hand held units, walk and turn the crank at a constant rate of speed.
- A shaker-type applicator made from a small container with holes punched in the bottom is recommended for small, difficult to treat areas. Carefully measure the amount of product needed to avoid over application.

SPREADER SETTINGS AS A GUIDE FOR CALIBRATION

Note: These settings are provided as a suggested starting point in calibrating each individual spreader. Since no two spreaders are alike, these settings are not intended to be used as absolute recommendations by Voluntary Purchasing Groups, Inc. or the spreader manufacturer.

Spreader	Settings for Spreader	
	Recommended Rate of ferti•lome® DOLLAR WEED CONTROL	
Revise Spreader and Settings as follows:	2.3 lbs./1,000 sq. ft.	3.4 lbs./1,000 sq. ft.
Scotts®/Republic Accugreen (Drop)	2.75	3.75
Scotts® Republic Speedy Green (Broadcast)	2.5	3.5
ferti•lome®/Earthway Ev-N-Spred (Broadcast)	13	14

HOW MUCH TO APPLY

Apply **ferti•lome® DOLLAR WEED CONTROL** to established turfgrass. Possible re-treatment in approximately 30 days may be necessary if turfgrass is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed specified rates because damage to turfgrass may occur. Do not make more than two applications per season.

Note: When applying to warm season turfgrass under conditions of inadequate moisture, high temperature and high humidity, over application may cause minor temporary discoloration. It is recommended to reduce application rates by 50% during hot, dry conditions to avoid possible dehydration or browning of turfgrass.

WHEN TO APPLY

Best weed control results are obtained when **ferti•lome® DOLLAR WEED CONTROL** is applied to small, actively growing weeds in the spring or early fall. Extreme growing conditions such as drought or near freezing temperatures prior to, at, or following time of application may reduce weed control. Broadleaf weed species germinate at different times.

TERMS AND CONDITIONS OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. Disclaimer, Inherent Risks of Use and Limitations of Remedies

INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. To the extent permitted by law, all such risks shall be assumed by the buyer.

LIMITATION OF REMEDIES

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at the Seller's election, one of the following:

- Refund of purchase price paid by buyer or user for the product bought, or
- Replacement of amount of product used

To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

Terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

ferti•lome® is a registered trademark of Voluntary Purchasing Groups, Inc.

Manufactured by:



230 FM 87 • BONHAM, TEXAS 75418

EPA Reg. No. 62719-550-7401 EPA Est. No. 7401-TX-01

Visit Us At: www.fertilome.com

Product Questions? 855-270-4776

