

For All Seasons Lawn Food Plus Crabgrass and Weed Preventer

Containing Prodiamine

 For selective preemergence control of grass and broadleaf weeds in established lawns

ACTIVE INGREDIENTS: Prodiamine: (N ³ ,N ³ -Di-n-propyl-2,4-dinitro-6) (trifluor	
phenylenediamine	0.383%
OTHER INGREDIENTS:	99.617%
TOTAL:	100.000%

KEEP OUT OF REACH OF CHILDREN CAUTION

See the following section, Precautionary **Statements, Hazards To Humans and Domestic Animals,** for additional information.

Net Weight 20 LBS. (9.08 KG)

For All Seasons Lawn Food Plus Crabgrass and Weed Preventer

10914







mos.emolitrata.uww.t4 zU size vww.tati bit seriention and the internet at: http://www.saptco.org/metaninet. noticenteepression regeneration and the series of the series of the internet at: http://www.saptco.org/metaninet

Produced For:

2-04-1-40-2

by a duly authorized representative of Voluntary Purchasing Groups.





and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Narranty, which may be varied only by agreement in writing signed

Voluntary Purchasing Groups warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions for Use** subject to the inherent risks referred to above. **Voluntary Purhosaing Groups makes no other express or implied Warranty Purchasing Groups or the Seller Directors of the for consequential, or indirect damages resulting that for the Buyer and user acceler.** Voluntary Purchasing Groups makes **no other express or indirect damages resulting that for the Buyer and user acceler to the force of the forcentary Purchasing Groups or the Buyer and user acceleration the factor of the forcentary Purchasing Groups and the factor of the forcentary Purchasing Groups and the factor of the forcentary Purchasing Groups and the factor of the forcentary Purchasing Groups and the factor of the Seller Officer and the Buyer and user acceleration is the forcentary Purchasing Groups and the factor of the forcentary of the factor of the fact**

ply the buyer. ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Voluntary Purchasing Groups or the Seller. All such risks shall be assumed by the burden power

and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Grop injury, The Directions for Use of this product reflect the opinion of the experts based on field use and tests. The directions are believed to be reliable

VTNARRAW GNA BLAR AND WARRANTY

SPILL: In case of spill, sweep up and dispose of material according to Product disposal directions above.

outdoor drain.

available. If Partly Filled: Call your local waste agency for disposal instructions. Never place unused product down any indoor or be used, carefully following label directions and precautions. If Empty: Do not reuse this bag. Place in trash or offer for recycling if PRODUCT DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag

by storage or disposal.

STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, food-stuffs, feed, or seed

JASO92ID DNA 3DA90T2


F370	Derived from: Ammonium Sulfate, Urea, Muriate of Potash, Sodium Borate, Copper Sulfate, Ferrous Sulfate, Manganese Sulfate, and Zinc Sulfate.
%90.0	Zinc (AZ) 0.05% Water Soluble Zinc (Zn)
%90.0	(nM) əsənsgnaM 0.05% Water Soluble Manganese (nM)
%01.0	Iron (Fe). 0.10% Water Soluble Iron (Fe)
%90.0	Copper (Cu) 0.05% Water Soluble Copper (Cu)
%20.0	Boron (B)
%8	Soluble Potash (K ₂ 0)
	0.82% Ammoniscal Nitrogen 15.18% Urea Nitrogen

.. (N) nəgortiN lstoT *ΣΙΣΥΙΑΝΑ* **ΠΕΕΓΝΑΡΑΟΘ**

8-0-9L

Precision Drop FH-22D	6
Precision Broadcast/Red Devil	01
Republic Drop	9
Republic/Scotts Broadcast	ß
Scott Broadcast/Speedy Green	L
Earthway Drop 7300B	۷۱ کا
ferti•lome@ Deluxe Broadcast/Earthway 2050 S.U.	14
ferti•lome@ Plus Iron Prof. Broadcast/Earthway 28002	14
fedielome@ Plus Iron Broadcast/Earthway 2850	14
ferti•lome@ By Spyker	<u>S</u>

5 – Apply January 1 to March 20 02 donald A – After March 20 0S lingA notes and 20 km a 01 Yatter May 10



This product is recommended for use on the turt grass species listed in the following table. Do not exceed the maximum yearly **SATAR JAUNNA MUMIXAM**

Maximum use rate selection should be based on turf species. The length of time residual weed control provided by this product year. All applications must be made prior to germination of the target weeds. This product will not control established weeds.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the RATES OF APPLICATION

may result in reduced weed control. (6) Do not apply this product where dichondra, colonial bentgrass, velvet bentgrass or annual

low tertility, or pest damage may result in turf injury. (5) Disturbing the herbicide barrier with cultural practices such as disking after application. Do not apply to newly set sod until the following year. (4) Application of this product to turt stressed by drought,

seedling secondary roots are in the second inch of soil, not thatch plus soil. (3) Do not cut (harvest) treated sod before 120 days

Use PRECAUTIONS: The following precautions apply to the use of this product in turf grasses and lawns: (1) Application of this

seequid or nutil after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before product may thin emerged annual bluegrass and newly overseeded grasses. (2) Do not apply to overseeded turf within 6U days after

germination dates.

10014

rate as given in the following table.

bluegrass (**Poa annua**) are desirable species.

is related to the rate applied.

 \mathbf{X}

46.25" Web

Width

۷

prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed weed control in turt grasses will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation applied per year is given for each turt grass species in the Maximum Annual Rates section of this label. Most effective grass and broadleat weeds in established turt grasses and lawns. The maximum amount of this product that may be ESTABLISHED TURF: This product is a selective preemergence herbicide that, when properly applied, will control certain

Apply uniformly with suitable, calibrated application equipment.

SNOITDERIGN DIRECTIONS

upplication rates recommended for each turf species. All applications must be made prior to germination of the weed seeds. ssuppression only. Sequential application may be made so long as the total amount of product applied does not exceed the maximusu

sApplication for this weed should be made in late summer, fall, or winter prior to germination.

woled elder turt for turt grass species listed in the Maximum Matter Table below. after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Hate Table. Do not exceed the In many areas a single application of 3.9 to 9.0 lbs/1000 sq. ft. of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 3.9 to 6.0 lbs/1000 sq. ft. followed

Knotweed	Woodsorrel, yellow (from seed)
Junglerice	Witchgrass
Johnsongrass (from seed)	Spurge, prostrate
ltchgrass	Sprangletop
tidnəH	Speedwell, Persian
Goosegrass¹	Signalgrass, Broadleaf
Foxtails, Annual	Shepherd's purse ²
Cupgrass, Woolly	Rescuegrass ³
Crowfootgrass	Pusley, Florida
Crabgrass (large, smooth)	Pursiane, common

Chickweed, mouse ear (from seed)	Pigweed
Chickweed, common	Panicum (Texas, Fall, Bro
Carpetweed	Fovegrass
Bluegrass, Annual (Poa annua)	Lambsquarter, common
Barnyardgrass	Kochia

When used in accordance with this label this product will provide control of the following weeds:

MEEDS CONTROLLED

СОИТROL, СROP INJURY, ОR ILLEGAL RESIDUES.

purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research

control will be obtained when it is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1-2 inches) incorpora-This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product is a selective preemergence

back onto the treated area of the lawn or garden will help prevent run off to water bodies or drainage systems.

DIRECTIONS FOR USE

or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street,

surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind adjacent sites. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or

the level of water solubility, it may be toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations above

ZQAAZAH JATNAMNOAIVNA

Call a poison control center or doctor for treatment advice. Rinse skin immediately with plenty of water for 15 to 20 minutes.

Call a poison control center or doctor for treatment advice.

Do not give anything by mouth to an unconscious person.

Have person sip a glass of water if able to swallow.

• Call a poison control center, or doctor, or when going for treatment advice.

Remove contact lenses, it present, after 5 minutes, then continue rinsing.

Hold eye open and rinse slowly and gently with water for 15-20 minutes.

Call a poison control center or doctor immediately for treatment advice.

DIA TZAIA

or clothing, Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact

STATEMENTS YAANOTUADAA Slamina Ditsemod GNA Snamuh ot Sqaasah

CENTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes,

Do not induce vomiting unless told to do so by a Poison Control Center or doctor.

If person is not breathing, call 911 or an ambulance, then give artificial respiration,

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECUNIONS ON THIS LABEL MAY RESULT IN POOR WEED

herbicide that provides residual control of many grass and broadleaf weeds in established lawns.

You may also call 1-800-858-7378 for emergency medical treatment advice.

Move person to fresh air.

while handling the material may cause allergic reactions in some individuals.

Take off contaminated clothing.

preferably mouth to mouth if possible.

:bəledni ti

:pnidtol0 r0

lt On Skin

lt In Eyes:

:bewollsw2

tion, prior to weed seed germination and within 14 days following application.

(dotuw

Do not apply this product through any type of irrigation system. Do not apply serially. Do not apply to golf course putting parks, or on golf courses or lawns and grounds. Do not graze or feed livestock forage cut from areas treated with this product.

4tuo2	Transition	Иоци	and Weed Preventer (Ibs./1000 sq. ft.)
4	4	4	3.0
4	4	G	3.9
G	S	9	4°2
9	9	-	4.8
L	L	-	0.8
6	-	-	8.8

This product will inhibit the germination of turt species if overseeded too soon after application. Follow rates and intervals in the

desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to

• Use higher rates of this product to achieve higher levels of felility and longer periods of weed control for each turf type, but do not exceed the maximum application rates

rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass, and Rescuegrass due to reduced product rates used in sprigging situations. y be see on the second or plugged or plugged or plugged served 3.0 lbs/1001 sq. ft. of this product. Newly sprigged or plugged bernudagrass stores and second or solved for the second or plugged or

WHEN TO OVERSEED AFTER APPLICATION

ARUT DNIDERSAEVO AETER OVERSEEDING TURE

(1) For All Seasons Lawn Food Plus Crabgrass and Weed Preventer BY TURF GRASS SPECIES (1)

SPREADER GUIDE

SPREADER

0.6

8.7

For All Seasons Lawn Food Plus Crabgrass

following table below for best overseeding/reseeding results.

• Do not apply more than 17.2 lbs./1000 sq. ft. per calendar year of this product.

I nese are the maximum rates per calendar year by species limitations.

thatch plus soil.

pisyo2

specified in the Maximum Annual Rates Table

Tall Fescue (including turf-type)

St. Augustinegrass

Seashore Paspalum

Centipedegrass

Rermudagrass⁽²⁾

Perennial Ryegrass

Kentucky Bluegrass

Creeping Red Fescue

Turf Species:

Creeping Bentgrass

Bahiagrass

Buffalograss

the applicator walks, spreader condition and the texture of the turf will alter the rate of These settings can be used only as a guide, they are not absolute. The speed at which

coverage, spread remainder over same area going at right angles (See Diagram).



15

٩0

SPREADER SETTINGS

%9L

application. If product remains in spreader after application to recommended square foot

Months Before Overseeding

0.6

0.0

6.4

3.9

Lbs. Products/1000 sq. ft.