

Kills Norway Rats, Roof Rats, House Mice and Voles. First dead rats, mice and voles will appear 4 to 5 days after feeding begins.











Warfarin: 3-(alpha-Acetonylbenzyl-

EPA Reg. No. 72500-6 EPA Est. 72500-CO-1





New Solutions to Old Problems

Keep Out of Reach of Children CAUTION

Harmful if swallowed.

See back panel for additional precautionary statements.

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals CAUTION

Keep away from humans, domestic animals and pets Harmful if swallowed or absorbed through the skin because this material may reduce the clotting ability of blood and cause bleeding. Do not get in eyes, on skin or clothing. Wash arms, hands and face with soap and water after applying and before eating or smoking.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

If swallowed, call a poison control center, doctor or 1-800-858-7378 for immediate treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor.

Notice to Physician or Veterinarian

This material may reduce the clotting ability of blood and cause bleeding. If ingested, administer Vitamin K-1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Do not apply directly to water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. IMPORTANT: Do not expose children, pets or other non-target animals to rodenticides. To help prevent

accidents:

1. Store product not in use in a location out of reach of children and pets.

2. Apply bait in locations out of reach of children, pets, domestic animals and non-target widdlife. OR in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and children under six year of age, and must be used in a manner that prevents such children from reachinto bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals or in areas prone to vandalism.

Dispose of product container, unused, spoiled and unconsumed bait as specified on this label.
 Do not apply this product for any pests or sites or by any application method not specified on this

RATS AND MICE

Use Restrictions: This product may be used to control Norway rats, roof rats and house mice at the sites in "Urban Areas" and "Non-urban Areas" that are indicated below. Do not place bat in areas where there is a possibility of contaminating food or surfaces that come into contact with food. Do not use this product to control Peromyscus spp. mice or within the occupied habitats of threatened or endangered Peromyscus spp. mice or other types of threatened or endangered redorms. Wear gloves

when handling bait or animal carcasses. Do not broadcast bait.

URBAN AREAS: To control Norway rats, roof rats and house mice this product may be used in and around homes and industrial, commercial and public buildings; in transport vehicles (ships, trains and aircraft) and in and around related port or terminal buildings. Applications "around" buildings must be made in tamper-resistant bait stations situated along walls or other external parts of buildings (e.g., doorways, ramps and loading docks) where rats or mice might seek to gain entrance.

NON-URBAN AREAS: To control Norway rats, roof rats and house mice this product may be used in and around homes and agricultural buildings.

SELECTION OF TREATMENT AREAS: Determine areas where rodents will most likely find and consume bait. Generally, these areas will be along walls, by gnawed openings, in corners and concealed places, between floors and walls, beside burrow openings and other locations where signs of rodents have been seen. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

Norway Rats and Roof Rats: Apply 2 to 8 placepacks per bait placement. Space bait placements at intervals of 15 to 30 feet in the infested area, at locations consistent with the requirements of this label. Maintain an uninterrupted supply of fresh bait for 10 days or until there no longer are fresh signs of rat activity or signs that task are teeding on the bait.

House Mice: Open pack and apply ¼ to ½ ounce of bait per typical placement. Larger placements – up to one unopened 2-ounce pack – may be made at points of extremely high mouse activity. Space bait placements at intervals of 8 to 12 feet in the infested area, at locations consistent with the requirements of this label. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are fresh signs of house mouse activity or signs that house mice are feeding on the bait.

Follow-up (Rats & Mies): Replace contaminated or spoiled bait immediately. Collect and properly dispose of all visible dead animals and lettover bait. To discourage reinfestation, limit sources of rodent food, water and harborage as much as possible. If reinfestation occurs, repeat treatment. Where there is a continuous source of reinfestation, establish permanent bait stations and replenish bait as repeated.

VOI F

Use Restrictions: This product may be used to control meadow voles (Microtus pennsylvanicus), pine voles (M. pinetus) and montane voles (M. montanus) around (usually within 15 feet of) homes and industrial, commercial, agricultural and public buildings. Do not use this product within the habitat of threatened or endangered Microtus spp. voles, threatened or endangered Microtus spp. mice, or other threatened or endangered rodents. Do not expose bait on bare ground. Unless children, pets, birds and other wildlife are excluded from the treated areas, all applications for controlling voles must be made in tamper-resistant bait stations. All bait abolications must be made by

hand. Wear gloves when handling bait or animal carcasses. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas near buildings where voles or signs of vole activity (griding of plants, consumption of bulbs, burrow openings and surface runways) have been seen. These areas generally will be in flower beds or under shrubs, bushes or ground cover.

APPLICATION DIRECTIONS: Open pack and load one ounce of bait into each bait station. Struate bait stations near runways, burrow openings or plants that voles have been damaging, in, and only if, children pets, domestic animals and wildlife reliably are excluded from the treated area, bait may be applied in one-ounce amounts directly into vole runways or burrow openings. Cover treated runways or holes with a shingle or board, or ground vegetation, if present. Check placements daily. Remove any fouled or spoiled bait and replenish if vole activity and interest in balt persist. Maintain an uninterrupted supply of fresh bait for 15 days or until there no longer are fresh signs of vole activity and feeding on bait. Collect and properly dispose of any carcasses and leftover bait.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in a cool, dry place inaccessible to children and pets.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent) then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or incineration; or if allowed by state and local authorities, by burning; if burned, stay out of smoke

WARRAN

To the extent consistent with applicable law, Manufacturer and Seller make no warranty of merchantability, fitness for any purpose, or otherwise express or implied, concerning this product or its use which extend beyond the statements on this label.



New Solutions to Old Problems
P.O. Box 1045 Wellington, CO 80549
(970) 482-1330

Label Version 9/4/07

FOR CALIFORNIA: WARNING: THIS PRODUCT CONTAINS A CHEMICAL KNOWN TO THE STATE OF CALIFORNIA TO CAUSE BIRTH DEFECTS OR OTHER REPRODUCTIVE HARM.



8