

TRUSTED SINCE 1926 BONIDE

Controls Mosquito Larvae in 24 hrs.

Controls mosquito larvae in standing (contained) water



#### **ACTIVE INGREDIENT\*:**

Bacillus thuringiensis subsp. israelensis . . . 1.7%

(Equivalent to 100 ITU/mg) OTHER INGREDIENTS: .....

and potency measurements are not federally standardized. EPA Reg. No. 4-455 EPA Est. No. 4-NY-1

Will not harm fish or other beneficial aquatic organisms.

Keep Out Of Reach Of Children
CAUTION (See Back Panel for Additional
Precautionary Statements)

# Mosquito Beater

- Kills developing mosquitoes (mosquito larvae) before they become breeding, biting adults including those which can carry West Nile virus, Encephalitis and Equine Encephalitis.
- · Water soluble No Mixing No Mess
- One pouch treats up to 50 sq. ft. of water
- Floating pouch dissolves within minutes, releasing the granules for uniform coverage
- Works even in stagnant, polluted or algae prevalent water
- Can be used in all types of containerized standing water sites where mosquitoes breed, except finished, treated human drinking water sources (See Directions for Use for specific use sites).
- Can be used in conjunction with other Bonide® Mosquito Beater® products

# PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if inhaled or absorbed through the skin. Avoid contact with skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. ENVIRONMENTAL HAZARDS: Do not apply to treated, finished drinking water reservoirs or drinking water receptacles, when the water is intended for human consumption.

#### **FIRST AID**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

Center at 1-80	00-858-7378 for emergency medical treatment
information.	
If on Skin	Take off contaminated clothing.
or Clothing:	Rinse skin immediately with plenty of water
	for 15-20 minutes.
	Call a poison control center or doctor for
	treatment advice.
If Inhaled:	Move person to fresh air.
	If person is not breathing, call 911 or an
	ambulance, then give artificial respiration,
	preferably mouth-to-mouth if possible.
	Call a poison control center or doctor for
	further treatment advice.

## **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

BONIDE® Mosquito Beater® WSP™ is a highly selective microbial insecticide, effective against mosquito larvae in standing water. The active ingredient, Bacillus *thuringiensis* subsp. *israelensis* (or Bti) kills mosquito larvae upon ingesting. BONIDE® Mosquito Beater® WSP™ will not harm fish, amphibians or other [beneficial] aquatic organisms.

BONIDE® Mosquito Beater® WSP™ can be applied to any water site except, 1) New York State waters, 2) treated, finished water reservoirs or drinking water receptacles, when the water is intended for human consumption. Use BONIDE® Mosquito Beater® WSP™ to control mosquito larvae in standing water in irrigation or drainage ditches, pastures, livestock watering troughs/ponds/tanks, lawns, woodland pools, snow melt pools, tidal water, salt marsh, catch basins, storm water retention basins, water gardens, flower pots, rain barrels, roof gutters, bird baths, tree holes, unused swimming pools or spas, discarded automobile tires, and any other location around homes or other buildings where water can collect and form a breeding ground for mosquitoes and remain for extended periods.



BONIDE® Mosquito Beater® WSP™ is provided in a watersoluble pouch to be placed directly on the surface of the water to be treated. The pouch will float on the water surface and disolve within a few minutes, releasing granules coated with Bti. These granules then slowly release the long-term Bti larvicide.

HOW TO APPLY: [APPLICATION RATE] 0.2 oz. will treat up to 50 sq. ft. of water surface. [An area approximately 7' x 7'] When treating larger areas, distribute uniformly on the water surface to ensure adequate dispersion of the granules. Lower rate (0.2 oz/50 sq. ft.) can be used in clear water, against low larval infestations or young larvae (from egg hatch to mid-third instar). When larger larvae (late third instar to early fourth instar) predominate, larval populations are high, or water is polluted and/or algae are prevalent, use the higher rate of 0.4 oz per 50 sq. ft.

The following table can be used as a guide for determining how much **BONIDE® Mosquito Beater® WSP™** to use in a particular situation. Application rates for areas smaller than 50 sq. ft, are listed after the table.

		Water surface area to be treated	
Water Condition	Mosquito Infestation	Up to 50 sq. ft.	100 sq. ft.
Clear Water	Light infestation or	1 pouch	2 pouchs
(containing little	small larvae		
or no algae or			
other organic	Heavy infestation or	2 pouches	3 pouches
matter)	large larvae		
Polluted water,	Light infestation or	2 pouchs	3 pouchs
high organic	small larvae		
matter, or algae			
present	Heavy infestation or	3 pouches	4 pouches
	large larvae	·	

**SMALL AREA APPLICATION:** To apply this product to small, specific areas such as bird baths, tree holes, flower pots, etc... sprinkle granules onto the area to be treated, at the following rates:

AREA TO BE TREATED	QUANTITY TO USE
1 to 12½ sq. ft.	1/4 of a pouch
12½ to 25 sq. ft.	½ of a pouch
25 to 371/2 sq. ft.	¾ of a pouch

Application can be repeated every 7 to 14 days, as needed, if environmental conditions favor continuous breeding of mosquitoes. Longer periods of mosquito suppression may result where sufficient numbers of aquatic parasites or predators of mosquito larvae are present, since these are not affected by **BONIDE® Mosquito Beater® WSP™** and contribute to overall reduction in mosquito populations.

## STORAGE & DISPOSAL

Storage: Store and transport in an upright position. Store in a cool dry area inaccessible to children or pets.

Pesticide Disposal: If empty - Nonrefillable. Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled - Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

Buyer's Guarantee Limited to Label Claims.
Bonide Products, Inc.
6301 Sutliff Road
Oriskany, NY 13424
©Bonide Products, Inc. All Rights Reserved