



Southern Agricultural Insecticides, Inc. O. Box 218 Palmetto, FI 34220

LIQUID IRON MINIMUM GUARANTEED ANALYSIS:

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Avoid contact with skin and eyes. Avoid breathing spray mist. Do not take internally.

This product WILL SE-VERELY STAIN concrete, masonry and painted surfaces.

Be extremely careful not to allow spray or runoff to contact with these materials. Do not mix with other fertilizers, fungicides or insecticides unless compatibility is known.

FIRST AID

IF SWALLOWED: Drink 2 glasses of water or milk and seek medical attention. EYE OR SKIN CONTACT: Flush thoroughly with a large volume of water. If irritation persists seek medical attention.

DIRECTIONS FOR USE LAWNS

Use: 8 - 16 fl. oz. per 1,000 sq. ft. (2-3 tablespoonful per 100 sq. ft.)
Apply in a sufficient volume of water to thoroughly wet the foliage and run-off in the root zone. Hose attached sprayers provide a convenient method to treat lawns.

FLOWERS, ROSES, SHRUBS, TREES and VEGETABLES

FOLIAGE APPLICATION: Mix this material at the rate of 1 tablespoonful per gallon. Apply in a manner that will give thorough coverage to upper and lower surfaces of foliage. Use an amount sufficient to cause runoff of spray. NOTE: Deficiency symptoms in many plants are an indication of improper pH. Soil pH should be checked and adjusted if necessary.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions and precautions given herewith.

For questions concerning this product, please email us at info@southernag.com our consult our website at www.southernag.com.