

TRIMEC® 992

BROADLEAF HERBICIDE

Controls Dandelion, Clover, Knotweed, Henbit, Chickweed, Plantain, Spurge and many other listed broadleaf weeds.

ONE GALLON COVERS UP TO 2.5 ACRES

KEEP FROM FREEZING

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	30.56%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid	8.17%
Dimethylamine salt of dicamba: 3,6-dichloro-o-anisic acid	2.77%
INERT INGREDIENTS:	58.50%
TOTAL 100.00%	

THIS PRODUCT CONTAINS:

2.38 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 25.38%.
 0.63 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 6.75%
 0.21 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 2.30%.



Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI-Gordon Corporation.

KEEP OUT OF REACH OF CHILDREN DANGER-PELIGRO

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle. (If you do not understand the label, find some one to explain it to you in detail.)

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170.



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER: Corrosive. Causes irreversible eye damage. Harmful if swallowed. Harmful if absorbed through skin. Do not get in eyes, on skin, or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are natural rubber. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators and other handlers must wear*:

- protective eyewear,
- long-sleeved shirt and long pants,
- shoes and socks,
- chemical-resistant gloves and
- chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

*Applicators may choose not to wear protective eyewear ONLY with dilution rates greater (higher) than 5:1 or greater (higher) than 5 parts of water to 1 part of product.

See engineering controls for additional requirements.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid	
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or on clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

9. For Use in Non-Cropland: Brush and Weed Control

High volume foliar applications:

Apply up to 1.15 gallons of product per 100 gallons of water or apply a 1.0 to 1.15% vol/vol spray solution as a full cover spray with high volume equipment. Use the lower spray concentrations in the range for susceptible species and use the higher spray concentrations within the range for hard-to-control species, for mature plants during the late summer or under adverse environmental conditions (e.g. drought).

Spray broadleaf weeds, woody plants or mixed brush uniformly and thoroughly by wetting all leaves, stems, bark and root collars. The minimum total volume of spray solution required for adequate coverage of solid stands of mixed brush is 25 gallons of spray solution per treated acre. The spray preparation chart for applications on a spray-to-wet basis is shown below in Table 2.

Table 2. Instructions for preparing 25 to 100 gallons of spray solution at 1.0 to 1.15% spray concentration with water for high volume foliar applications.		
Spray solution per acre, Gallons	Amount of Product Needed for Spray Concentration of:	
	1.0%	1.15%
25	2.0 pt	2.3 pt
75	6.0 pt	6.9 pt
100	1.0 gal	1.15 gal

Equal measures: 1 gallon = 4 quarts = 8 pints = 128 fl.oz.

For Backpack Sprayers and Hand Pump-up Sprayers.

Table 3. Instructions for preparing 1 to 3 gallons of spray solution at 1.0 to 1.15% spray concentration with water for high volume foliar applications.		
Gallons of Water	Amount of Product Needed for Spray Concentration of:	
	1.0%	1.15%
1	1.25 fl.oz. (2.5 Tablespoons)	1.5 fl.oz. (3 Tablespoons)
2	2.5 fl.oz.	3.0 fl.oz.
3	3.8 fl.oz.	4.5 fl.oz.

Equal measures: 1 fl.oz. = 2 tablespoons (Tbs.) = 6 teaspoons (tsp.)

BRUSH:			
Ash Aspen Birch Black cherry Brambles	Cedars Elms Gooseberry Honey locust Kudzu	Multiflora rose Oak Poison ivy Poison oak Shortleaf pine	Sumac Willow and many more brush species

Table 4. Restrictions on broadcast applications to ornamental turfgrass sites, sod farms, and non-cropland.				
Use Site	Maximum Rate per Application	Maximum Number of Applications per Year	Minimum Interval Between Applications	Maximum Seasonal Rate
Ornamental turfgrass sites and sod farms	4.0 pints/A (1.19 lb 2,4-D ae/A) (0.33 lb MCPP-p ae/A) (0.11 lb dicamba ae/A)	2	30 Days	8.0 pints/A (2.37 lb 2,4-D ae/A) (0.65 lb MCPP-p ae/A) (0.22 lb dicamba ae/A)
Non-cropland	4.0 pints/A (1.19 lb 2,4-D ae/A) (0.33 lb MCPP-p ae/A) (0.11 lb dicamba ae/A)	2 (annual and perennial weeds)	30 Days	8.0 pints/A (2.37 lb 2,4-D ae/A) (0.65 lb MCPP-p ae/A) (0.22 lb dicamba ae/A)

