

ferti•lome®



Weed-Out® Broadleaf Weed Control

CONTAINS TRIMEC® HERBICIDE

- Controls: Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and Many Other Broadleaf Weeds As Listed
- For Use On Bahia, Bermuda, Centipede, Zoysia, Kentucky Bluegrass, Perennial Rye, Fescue and Bentgrass Lawns

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid.....	1.65%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid.....	0.40%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid).....	0.18%
OTHER INGREDIENTS:	97.77%
TOTAL:	100.00%

THIS PRODUCT CONTAINS:


27.4 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton or 1.37%.
6.6 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.33%.
3.0 lbs. 3,6-dichloro-o-anisic acid equivalent per ton or 0.15%.
Isomer Specific by AOAC Methods.
TRIMEC® is a registered trademark of PBI/Gordon Corporation.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional Precautionary Statements and First Aid

NET WEIGHT 10 LBS. (4.54 KG)



ferti•lome® Weed-Out® Broadleaf Weed Control

ferti•lome® Weed-Out® Broadleaf Weed Control

Covers Up To 5,000 Sq. Ft.

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If In Eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a Poison Control Center or doctor for treatment advice.
If On Skin Or On Clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a Poison Control Center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none">• Call a Poison Control Center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a Poison Control Center or doctor.• Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment.	

In the event of a medical or chemical emergency, contact ChemTel™  North America 1-800-255-3924

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Do not apply this product to St. Augustinegrass, Carpetgrass, Dichondra, ornamentals, or lawns containing desirable clovers.
- Do not exceed specified dosages for any area. Treated areas may be seeded 3 to 4 weeks after application. Application of this product to newly seeded areas should be delayed until the third mowing.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product runoff, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

WHERE TO USE

ferti-lome® WEED-OUT® BROADLEAF WEED CONTROL is a **Trimec®** complex, a combination of three proven weed killers – 2, 4-D, Mecoprop-p and Dicamba. Together, they control a wide range of lawn weeds, including Dandelion, Chickweed, Knotweed, Plantains, Henbit, Spurge and many others found in home lawns. Established turf of Kentucky Bluegrass, Bentgrass, Bahiagrass, Centipedegrass, Fescues, Bermudagrass, Zoysiagrass, and Perennial Ryegrass may be treated.

HOW MUCH TO APPLY

For Kentucky Bluegrass, Bentgrass, Bahiagrass, Centipedegrass, Fescues, Perennial Ryegrass, Bermudagrass, and Zoysiagrass: Apply 2.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 2.0 pounds of product per 1,000 square feet should be used 30 days after the first application. The maximum broadcast application rate is 2.0 pounds of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 4.0 pounds of product per 1,000 sq. ft.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

CALIBRATION INSTRUCTIONS

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage. Do not exceed the application rate for any area.

Drop spreaders and rotary spreaders should be calibrated for uniform applications of this product. **NOTE: The spreader settings listed are approximate. The speed at which you walk and the working condition of the spreader will affect the rate of application.**

SPREADER	SPREADER SETTINGS	
	Lbs. of product per 1,000 sq. ft.	
	2 lbs.	4 lbs.
Scotts®/Republic Accugreen (Drop)	2 1/4	4 1/4
Scotts®/Republic Speedy Green (Broadcast)	2 3/4	3 3/4
ferti-lome®/Earthway Ev-N-Spred (Broadcast)	12	14

THE FOLLOWING BROADLEAF WEEDS ARE CONTROLLED BY THIS PRODUCT:

Annual Yellow Sweetclover Aster Austrian Fieldcross Bedstraw Beggarticks Betony, Florida Bindweed, Field Bird Vetch Bitter Wintercress Bittercress, Hairy Bitterweed Black-eyed Susan Black Medic Black Mustard Blackseed Plantain Blessed Thistle Bloodflower Milkweed Blue Lettuce Blue Vervain Bracted Plantain Bristly Oxtongue Broadleaf Dock Broadleaf Plantain Broomweed Buckhorn Buckhorn Plantain Bulbous Buttercup Bull Thistle Bullnettle	Burclover Burdock Burning Nettle Burweed Buttercup Buttonweed Canada Thistle Carolina Geranium Carpetweed Catchweed Bedstraw Catnip Catsear Chickweed, Common Chickweed, Mouseear Chicory Cinquefoil Clover, Crimson Clover, Hop Clover, Red Clover, Strawberry Clover, Sweet Clover, White Cockle Cocklebur Common Mullein Creeping Jenny Cudweed Curly Dock Daisy, English Daisy, Fleabane	Daisy, Oxeye Dandelion Dichondra Dogfennel Dollarweed False Dandelion Falseflax False Sunflower Fiddleneck Florida Pusley Frenchweed Galinsoga Goldenrod Ground Ivy Gumweed Hairy Fleabane Hawkweed Healall Heartleaf Drymary Heath Aster Hedge Bindweed Hedge Mustard Henbit Hoary Cress Hoary Plantain Hoary Vervain Horsenettle Jimsonweed Knapel	Knotweed Kochia Lambsquarters Lespedeza Mallow Matchweed Mexicanweed Milk Vetch Morningglory Mouseear Hawkweed Mugwort Musk Thistle Mustard Narrowleaf Plantain Narrowleaf Vetch Nettle Orange Hawkweed Oriental Cocklebur Oxalis Parsley-piert Parsnip Pearlwort Pennycress Pepperweed Pigweed Pineywoods Bedstraw Plains Coreopsis Plantain Poison Ivy	Poison Oak Pokeweed Poorjoe Prairie Sunflower Prickly Lettuce Prickly Sida Prostrate Knotweed Prostrate Pigweed Prostrate Spurge Puncturevine Purslane, Common Ragweed Red Sorrel Redroot Pigweed Redstem Filaree Rough Cinquefoil Rough Fleabane Russian Pigweed Russian Thistle Scarlet Pimpernel Scotch Thistle Sheep Sorrel Sheepspurge Slender Plantain Smallflower Galinsoga Smooth Dock Smooth Pigweed Sorrel	Sowthistle Spanishneedles Speedwell Spiny Amaranth Spiny Cocklebur Spiny Sowthistle Spotted Catsear Spotted Spurge Spurweed Stinging Nettle Strawberry, India Mock Tall Nettle Tall Vervain Tansy Ragwort Tansy Mustard Tanweed Thistle Trailing Crownvetch Tumble Mustard Tumble Pigweed Velvetleaf Venice Mallow Virginia Buttonweed Virginia Creeper Virginia Pepperweed Wavyleaf Bullthistle Western Clematis Western Salsify White Mustard	Wild Aster Wild Buckwheat Wild Carrot Wild Four-O'Clock Wild Garlic Wild Geranium Wild Lettuce Wild Marigold Wild Mustard Wild Onion Wild Parsnip Wild Radish Wild Rape Wild Strawberry Wild Sweet Potato Wild Vetch Woodsorrel Woolly Croton Woolly Morningglory Woolly Plantain Wormseed Yarrow Yellow Rocket Yellowflower Pepperweed
--	---	--	--	---	---	--

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER: FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Weed-Out® and ferti-lome® are registered trademarks of Voluntary Purchasing Groups, Inc

Manufactured By:
 **Voluntary
Purchasing
Groups, Inc.**
230 FM 87 • BONHAM, TEXAS 75418
EPA Reg. No.: 2217-657-7401
EPA Est. No.: 7401-TX-01
Visit Us At: www.fertilome.com
Product Questions? 855-270-4776

10925-0116-TP

