

# ORNAMEC®

## 170 GRASS HERBICIDE

**Kills unwanted grasses in and around ornamental plants.**  
**Spray ornamentals “over-the-top”**

**ACTIVE INGREDIENT:**  
 Fluazifop-P-butyl: Butyl(R)-2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoate . . . . . 1.70%  
**OTHER INGREDIENTS:** . . . . . 98.30%  
 TOTAL 100.00%

**THIS PRODUCT CONTAINS:**  
 0.125 lb. (+) isomer (fluazifop-P-butyl) per gallon.  
 Contains petroleum distillates.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
**SHAKE WELL BEFORE USING**



**READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

### PRECAUTIONARY STATEMENTS

**Hazards to Humans & Domestic Animals**  
**CAUTION:** Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves made of barrier laminate or viton. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

First Aid	
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>Immediately call a poison control center or doctor.</li> <li>Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>Do not give any liquid to the person. Do not give anything to an unconscious person.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice. <b>Note to Physician:</b> Contains petroleum distillates – vomiting may cause aspiration pneumonia.	

**User Safety Requirements**  
 Follow manufacturer’s instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards**  
 This product is toxic to fish. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

**Physical and Chemical Hazards**  
 Do not use or store near heat or open flame.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.  
 This labeling must be in the possession of the user at the time of application.  
 Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes and being grown in ornamental gardens or parks, or on golf courses, or lawns and grounds.

#### 1. Product Description

Ornamec® 170 Grass Herbicide is a selective postemergence herbicide for control of annual and perennial grass weeds. Ornamec 170 Grass Herbicide does not control broadleaf weeds or sedges (nutgrass).

**Control Symptoms – Ornamec 170 Grass Herbicide** is a systemic herbicide which moves from the treated foliage into the shoots, roots, rhizomes, stolons, and growing points (meristematic regions) of treated grass weeds. Excellent control of a wide range of annual and perennial grass weeds will be obtained with Ornamec 170 Grass Herbicide when applied as directed on this label.

Growth of treated grass weeds stops soon after application. Symptoms include loss of vigor, yellowing and/or reddening, and eventual death to the treated grass plant. Symptoms are generally observed within one to three weeks, depending on grass weed species and environmental conditions.

Ornamec 170 Grass Herbicide can be used to control grass weeds in many newly transplanted and established ornamentals, trees, shrubs and groundcovers. Apply to actively growing grass weeds before they exceed the recommended growth stages shown below. See the following Tables 7 through 10 for specific uses.

Make over-the-top application of Ornamec 170 Grass Herbicide to ornamentals listed in Table 7. Over-the-top applications are defined as spray patterns delivered to the upper portions of the established ornamentals and target weeds. Use sufficient spray volumes to ensure a uniform distribution and complete spray coverage of the target weeds. Make directed spray applications to plants listed in Tables 8 through 10 to prevent contact of spray with foliage. Directed spray applications are defined as spray patterns aimed or directed toward the lower portion of the established ornamentals, and this method of application will minimize the phytotoxicity of the desired species. If the growth habit of the plant or other factors do not allow a directed spray, an over-the-top application can be made to plants in Tables 6 though 8. However, there is the possibility of damage in the range listed for each table when applied at label rates. Do not apply to ornamentals that may be harvested for food within one year after application.

**NOTICE TO BUYER AND USER:** It is impossible to test every species and variety or cultivar of ornamental plants under all conditions. Plant tolerance of pesticides vary as conditions vary. Plant tolerance of Ornamec 170 Grass Herbicide at label rates has been found to be acceptable within the ranges specified for the indicated genera and species. Neither the manufacturer nor the seller has determined whether Ornamec 170 Grass Herbicide can be used on plants not specified on this label. The user should determine if Ornamec 170 Grass Herbicide can be used prior to use.

## 2. Annual and Perennial Grass Control in Ornamentals, Trees, Shrubs, and Groundcovers

**Not for use in vegetable gardens or on fruit-bearing trees and shrubs.**

Spray to obtain thorough coverage of grass foliage, but not to the point of runoff. Do not apply Ornamec 170 Grass Herbicide if rainfall is expected within 1 hour. Apply Ornamec 170 Grass Herbicide to actively growing grasses before they exceed growth stages specified in Tables 5 and 6. Apply Ornamec 170 Grass Herbicide when the first grass species in a mixed grass weed population reaches the specified growth stage for treatment.

Do not apply Ornamec 170 Grass Herbicide to grasses which are stressed due to moisture, temperature, low soil fertility, mechanical or chemical injury. Do not apply to grasses which have exceeded the specified growth stages (See Tables 5 and 6).

**BROADCAST TREATMENTS:** For annual grass species, use 6 to 9 fl.oz. of Ornamec 170 Grass Herbicide per 1,000 square feet. For perennial grass species, use 9 fl.oz. of Ornamec 170 Grass Herbicide per 1,000 square feet per application.

The maximum amount of fluzifop-R-butyl per treated area is 1.1lb. a.i./acre per year (27 fl.oz./1000 sq.ft. per year).

**SPOT TREATMENTS:** Mix Ornamec 170 Grass Herbicide with water according to the amounts shown in the following chart. Spray to obtain thorough coverage, but do not spray to the point of runoff.

To Make This Spray Volume	Add This Amount of Ornamec 170 Grass Herbicide
24 fl.oz.	3.5 tablespoons
32 fl.oz.	2.5 fl.oz. (5 tablespoons)
1 gallon	10 fl.oz. (18 tablespoons)

Equal measures: 1 fluid ounce (fl.oz.) = 2 tablespoons (Tbs) = 6 teaspoons (tsp) and 8 fl.oz. = 1 cup

## 3. Bermudagrass Control in Zoysiagrass and Tall Fescue Turf\*

Use Ornamec 170 Grass Herbicide to suppress and eventually control some undesirable grasses in turf areas. To control common bermudagrass and hybrid bermudagrass, follow the directions below:

Note: The application of this product may show slight discoloration to the desirable turfgrass. Symptoms are temporary and discoloration will subside. Do not apply to tall fescue turf unless injury can be tolerated. Weather and cultural treatments can also affect applications.

\*Not applicable in California

### 3.1 Bermudagrass Control in Zoysiagrass\*

**For spot treatments of zoysiagrass with hand-operated sprayers:** Begin applications in late spring (near June 1) as bermudagrass begins growth. Repeat applications at 30-day intervals throughout the summer until fall to continue the growth suppression of bermudagrass. Complete control may take 1 to 2 growing seasons.

Avoid applications to zoysiagrass under stress due to drought, temperature extremes, or chemical injury. Use hand-operated sprayers (backpack sprayers, compression or pump-up sprayers, or tank type sprayers). Add 1.3 fl.oz. of product per 1 gallon of water to treat 1,000 square feet of turf. Additional spray preparations for zoysiagrass are presented in the following table:

To Make This Spray Volume	Add This Amount Of Ornamec 170 Grass Herbicide
1 gallon	1.3 fl.oz. (8 tsp)
2 gallons	2.6 fl.oz. (16 tsp)
5 gallons	6.5 fl.oz. (40 tsp)

Equal measures: 1 fluid ounce (fl.oz.) = 2 tablespoons (Tbs) = 6 teaspoons (tsp) and 8 fl.oz. = 1 cup

Allow time (up to 24 hours) for this product to be absorbed and translocated before mowing the treated area.

\*Not applicable in California

### 3.2 Bermudagrass Control in Tall Fescue\*

**For spot treatments of tall fescue with hand-operated sprayers:** Make applications in the spring and fall. Start applications after green up. Take care with applications to tall fescue during the summer months (July and August). Repeat in 14 to 21 day intervals if regrowth occurs. Complete control may take 1 to 2 growing seasons.

Avoid applications to tall fescue under stress due to drought, temperature extremes, or chemical injury. Use hand-operated sprayers (backpack sprayers, compression or pump-up sprayers, or tank type sprayers). Add 2 fl.oz. of product per 1 gallon of water to treat each 1,000 square feet of turf. Additional spray preparations for tall fescue are presented in the following table:

To Make This Spray Volume	Add This Amount Of Ornamec 170 Grass Herbicide
1 gallon	2 fl.oz. (12 tsp)
2 gallons	4 fl.oz. (24 tsp)
5 gallons	10 fl.oz. (64 tsp)

Equal measures: 1 fluid ounce (fl.oz.) = 2 tablespoons (Tbs) = 6 teaspoons (tsp) and 8 fl.oz. = 1 cup

Allow time (up to 24 hours) for this product to be absorbed and translocated before mowing the treated area. Wait 14 days before re-seeding treated areas of tall fescue.

Note: The application of this product may show slight discoloration to the desirable turfgrass. Symptoms are temporary and discoloration will subside. Do not apply to tall fescue turf unless injury can be tolerated. Weather and cultural treatments can also affect applications.

\*Not applicable in California

## 4. Dallisgrass Control in Tall Fescue\*

For spot treatments of tall fescue: Make a single application in the spring. Add 2 fl.oz. of Ornamec 170 Grass Herbicide per one gallon of water to treat 1,000 sq.ft. to control dallisgrass in tall fescue turf.

Note: The application of this product to tall fescue turf may show slight discoloration. Symptoms are temporary and discoloration will subside. Do not apply to tall fescue turf unless injury can be tolerated. Weather and cultural treatments can also affect applications. Use a minimum spray volume of 30 gallons of water per acre.

Application timing is critical and is determined by Growing Degree Days (GDD). GDD units are determined by the following calculation:

$$\frac{\text{Max daily temp (in } ^\circ\text{F)} + \text{Minimum daily temp (in } ^\circ\text{F)} - 50^\circ\text{F}}{2} = \text{GDD}$$

If the calculation is zero or a negative number, it is ignored. Add each day's GDD units beginning on January 1st. When your total reaches 270 to 360 GDD, apply the Ornamec 170 Grass Herbicide. Effective dallisgrass control will only be obtained in this GDD range.

Date	Max Temp	Min Temp	Average	Subtract 50	Daily GDD	Total GDD
Day 1	75	45	60	10	10	10
Day 2	65	45	55	5	5	15
Day 3	55	45	50	0	0	15
Day 4	65	45	55	5	5	20
Day 5	50	45	47.5	-2.5	-2.5	20

\*Not applicable in California

### 5. Grass Species Controlled

Table 5: Annual Grasses	
Grass Species	Growth Stage
Barnyardgrass (Echinochloa crus-galli)	2 to 8 inches tall, before tillering and/or heading.
Broadleaf signalgrass (Brachiaria platyphylla)	2 to 8 inches tall, before tillering and/or heading.
Crabgrass, large (Digitaria sanguinalis)	2 to 8 inches tall, before tillering and/or heading.
Crabgrass, smooth (Digitaria ischaemum)	2 to 8 inches tall, before tillering and/or heading.
Crabgrass, southern (Digitaria ciliaris)	2 to 8 inches tall, before tillering and/or heading.
Crabgrass, tropical (Digitaria bicornis)	2 to 8 inches tall, before tillering and/or heading.
Downy brome (Bromus tectorum)	2 to 8 inches tall, before tillering and/or heading.
Fall panicum (Panicum dichotomiflorum)	2 to 8 inches tall, before tillering and/or heading.
Field sandbur (Cenchrus pauciflorus)	2 to 8 inches tall, before tillering and/or heading.
Foxtail, giant (Setaria faberi)	2 to 8 inches tall, before tillering and/or heading.
Foxtail, green (Setaria viridis)	2 to 8 inches tall, before tillering and/or heading.
Foxtail, yellow (Setaria lutescens)	2 to 8 inches tall, before tillering and/or heading.
Goosegrass (Eleusine indica)	2 to 8 inches tall, before tillering and/or heading.
Italian Ryegrass (Lolium multiflorum)	2 to 8 inches tall, before tillering and/or heading.
Itchgrass (Rottboellia exaltata)	2 to 8 inches tall, before tillering and/or heading.
Johnsongrass, seedling (Sorghum halepense)	2 to 8 inches tall, before tillering and/or heading.
Junglerice (Echinochloa contracta)	2 to 8 inches tall, before tillering and/or heading.
Kikuyugrass* (Pennisetum clandestinum)	4 to 8 inches tall, before tillering and/or heading.
Prairie cupgrass (Eriochloa contracta)	4 to 8 inches tall, before tillering and/or heading.
Rabbitfootgrass (Polypogon monspeliensis)	2 to 8 inches tall, before tillering and/or heading.
Red rice (Oryza sativa)	2 to 8 inches tall, before tillering and/or heading.
Shattercane/Wildcane/Volunteer milo (Sorghum bicolor)	2 to 8 inches tall, before tillering and/or heading.
Sorghum alnum (Sorghum alnum)	2 to 8 inches tall, before tillering and/or heading.
Southern sandbur (Cenchrus echinatus)	2 to 8 inches tall, before tillering and/or heading.
Southwestern cupgrass (Eriochloa gracilis)	2 to 8 inches tall, before tillering and/or heading.
Texas panicum (Panicum texanum)	2 to 8 inches tall, before tillering and/or heading.
Volunteer barley (Hordeum vulgare)	2 to 8 inches tall, before tillering and/or heading.
Volunteer corn (Zea mays)	2 to 8 inches tall, before tillering and/or heading.
Volunteer oats (Avena fatua)	2 to 8 inches tall, before tillering and/or heading.
Volunteer rye (Secale cereale)	2 to 8 inches tall, before tillering and/or heading.
Volunteer wheat (Triticum aestivum)	2 to 8 inches tall, before tillering and/or heading.
Wild oats (Avena fatua)	2 to 8 inches tall, before tillering and/or heading.
Wild proso millet (Panicum milliaceum)	2 to 8 inches tall, before tillering and/or heading.
Witchgrass (Panicum capillare)	2 to 8 inches tall, before tillering and/or heading.
Woolly cupgrass (Eriochloa villosa)	2 to 8 inches tall, before tillering and/or heading.

\*Not for use in California

Table 6: Perennial Grasses	
Grass Species	Growth Stage
Bermudagrass (Cynodon dactylon)	4 to 8 inch runners
Dallisgrass** (Paspalum dilatatum)	270 to 360 GDD**
Guineagrass (Panicum maximum)	6 to 12 inches tall before seedhead initiation
Quackgrass (Agropyron repens)	6 to 10 inches
Rhizome johnsongrass (Sorghum halepense)	8 to 18 inches tall and before boot stage
Torpedograss*	3 to 10 inches tall (first application)
Torpedograss*	2 to 3 inches tall (subsequent application)
Wirestem muhly (Muhlenbergia frondosa)	4 to 12 inches tall before seedhead initiation

\*For control of Torpedograss, use 9 fluid ounces/1000sq.ft./application. 3 applications will be needed.  
 \*\*See Section 4. Dallisgrass Control in Tall Fescue for detailed instructions.

### 6. Ornamentals

Table 7. Over-The-Top Applications may be applied to the following ornamentals (Use only a nonionic surfactant).	
Common Name/Variety	Scientific Name
Abelia, Glossy	<i>Abelia grandiflora</i>
Acacia, Jim wheat	<i>Acacia schafnerii</i>
Acacia, Ongerops	<i>Acacia redolens</i>
Acacia, Shoe-string	<i>Acacia stenophylla</i>
Acacia, Willow	<i>Acacia saligna</i>
Acacia, Willow-leaved	<i>Acacia salicina</i>
Ageratum, sp.	<i>Ageratum sp.</i>
Almond, Flowering	<i>Prunus triloba</i>
Aloe vera	<i>Aloe vera</i>
Aloe zanzibarica	<i>Aloe zanzibarica</i>
Aloe, Barbados	<i>Aloe barbadensis</i>
Alyssum sp.	<i>Alyssum sp.</i>
Ash, American Mountain*	<i>Sorbus americana*</i>
Ash, Arizona	<i>Fraxinus velutina</i>
Ash, Green*	<i>Fraxinus pensylvanica*</i>
Ash, White*	<i>Fraxinus Americana*</i>
Asparagus, Myres; Asparagus, Sprenger, Sprengeri	<i>Asparagus densiflorus</i>
Aucuba	<i>Aucuba japonica</i>
Aucuba japonica variegata	<i>Aucuba japonica variegata</i>
Aurea	<i>Philadelphus coronarius</i>
Australian bush cherry	<i>Syzgium paniculatum</i>
Australian tea tree	<i>Leptospermum laevigatum</i>
Banana, Ethiopia	<i>Musa maurelli</i>
Banksia	<i>Rosa banksiae</i>
Barberry, Mentor	<i>Berberis mentorensis</i>
Barberry, Redleaf Japanese; Pygmy, Crimson*	<i>Berberis thunbergii*</i>
Bearberry, Red	<i>Arctostaphylos uva-ursi</i>
Begonia, Scarletta*	<i>Begonia Sempreflorens cultoreum*</i>
Bellflower	<i>Campanula carpatica</i>
Berkman's	<i>Thuja orientalis</i>
Birch, Eastern white*	<i>Betula pendula*</i>
Bird of Paradise	<i>Caesalpinia gilliesii</i>
Bird of Paradise	<i>Strelitzia reginae</i>
Bird of Paradise, Giant	<i>Strelitzia nicolai</i>
Bittle bush	<i>Encelia farinose</i>

(cont. on next page)

**Table 7 (cont).** Over-The-Top Applications may be applied to the following ornamentals (Use only a nonionic surfactant).

Common Name/Variety	Scientific Name
Blaauw's pink, Boule de neige, Delaware Valley white, Delaware Valley white, Fashio, Gerard's Rose, Gibraltar, Gloria, Greeting, Gumpo pink, Gumpo white, H.H. Hume, Hahm red, Herbert, Hino red, Kaempo, Kluis sensation, Masasoit, Mother's day, Pericat, Pink pearl, President Lincoln, Prize, Purple gem, Red ruffle, Red wing, Road runner, Rosebud, Royalty, Rutherfordiana Constances, Salmon spray, Snow, Stewartstonian, Sweetheart, Tabor, Tradition, White cascade	<i>Rhododendron sp.</i>
Blue point, Blue vase juniper, Gold coast juniper, Gold tip, Hetzi, Hollywood, Juniper, Keteleeri, Nicks compact juniper, Parsoni, Sargent juniper, Torulosa	<i>Juniperus chinensis</i>
Blue Star Creeper	<i>Isotoma sp.</i>
Bottle-brush	<i>Callistemon lanceolatus</i>
Bougainvillea sp.	<i>Bougainvillea sp.</i>
Boxwood, Common; Welleri	<i>Buxus sempervirens</i>
Boxwood, Japanese	<i>Buxus microphylla var. japonica</i>
Boxwood, Korean	<i>Buxus microphylla var. koreana</i>
Brown bean	<i>Sedum guatemalense</i>
Brush cherry	<i>Eugenia myrtifolia</i>
Buckthorn, Tallhedge	<i>Rhamnus frangula</i>
Burningbush, Compact	<i>Kochia scoparia f. trychophylla</i>
Cactus, Barrel	<i>Ferocactus sp.</i>
Cactus, Cholla	<i>Opuntia Cholla</i>
Cactus, Hedgehog	<i>Echinocactus sp.</i>
Cactus, Saguaro	<i>Carnegiea gigantea</i>
California pepper tree	<i>Schinus molle</i>
Camellia	<i>Camellia japonica</i>
Camellia, Sasanqua	<i>Camellia sasanqua</i>
Candelabra plant	<i>Euphorbia lacteal</i>
Cape weed	<i>Arctotheca calendula</i>
Caricature plant	<i>Graptophyllum pictum</i>
Carissa tuttlei	<i>Carissa tuttlei</i>
Carolina cherry	<i>Prunus caroliniana compacta</i>
Cascalote	<i>Caesalpinia cacalaco</i>
Cassia, African	<i>Cassia didymobotrya</i>
Cassia, Feathery	<i>Cassia artemisiodes</i>
Centaurea, Dusty miller	<i>Centaurea cineraria</i>
Century plant	<i>Agave americana</i>
Cerastium, Snow in Summer	<i>Cerastium tomentosum</i>
Ceratonia, Carob tree	<i>Ceratonia siliqua</i>
Cercis, Red bud	<i>Cercis canadensis</i>
Chionoides, Elizabeth Gable, Less dark purple, Purple elegans, Purple spendor, Rose Greeley, Roseum elegans, Roseum superbum, White catawba	<i>Rhododendron catawbiense</i>
Chives	<i>Allium schoenoprasum</i>
Cleyera	<i>Cleyera spp.</i>
Cleyera	<i>Ternstroemia gymnanthera</i>
Clover, Pink	<i>Polygonum capitatum</i>
Coffee	<i>Coffea Arabica</i>
Coleus, Jade wizard*	<i>Coleus x hybridus*</i>
Coolibah, Gum-barked	<i>Eucalyptus microtheca</i>
Coreopsis, Threadleaf	<i>Coreopsis verticillata</i>
Cotoneaster	<i>Cotoneaster microphyllus</i>
Cotoneaster	<i>Cotoneaster repens</i>
Cotoneaster apiculata	<i>Cotoneaster apiculata</i>
Cotoneaster, Coral beauty; Royal beauty	<i>Cotoneaster dammeri</i>
Cotoneaster, Spreading	<i>Cotoneaster divaricatus</i>
Cotoneaster, Willowleaf	<i>Cotoneaster salicifolius franch</i>

(cont. on next column)

**Table 7 (cont).** Over-The-Top Applications may be applied to the following ornamentals (Use only a nonionic surfactant).

Common Name/Variety	Scientific Name
Crabapple, Showy	<i>Malus floribunda</i>
Cranesbill	<i>Geranium pratense</i>
Creeping Charlie	<i>Pilea nummularifolia</i>
Crossandra	<i>Crossandra nilotica</i>
Croton	<i>Codiaeum variegatum</i>
Crown Vetch	<i>Vicia sp.</i>
Cypress, Allum lawson	<i>Chamaecyparis lawsoniana</i>
Cypress, Cripps hinoki false	<i>Chamaecyparis obtusa</i>
Cypress, Italian	<i>Cupressus sempervirens</i>
Daisy, Shasta	<i>Chrysanthemum x superbum</i>
Daisy, White africans	<i>Osteospermum fruticosum alba</i>
Daylily	<i>Hemerocallis hybrids</i>
Deutzia, Slender	<i>Deutzia gracilis</i>
Dianthus, Sweet William	<i>Dianthus barbatus</i>
Dogwood, Cornelia cherry	<i>Cornus mas</i>
Dogwood, Flaviramea; Red twig	<i>Cornus sericea</i>
Dogwood, Flowering	<i>Cornus florida</i>
Dumbcane, Giant	<i>Dieffenbachia amoena</i>
Emerald green, Globosa, Pyramidalis, Techny, Techny American arborvitae, white cedar, Woodwardii	<i>Thuja occidentalis</i>
Emerald mound	<i>Lonicera xylosteum</i>
Eranthemum, Purple false	<i>Pseuderanthemum atropurpureum</i>
Erythrina, Fastadiata; Swamp immortalia	<i>Erythrina fusca</i>
Escallonia fradessii	<i>Escallonia fradessii</i>
Escallonia rubra	<i>Escallonia rubra</i>
Euonymus fortunei	<i>Euonymus fortunei</i>
Euonymus, Siebold	<i>Euonymus alata</i>
Euonymus, Silver King	<i>Euonymus japonica</i>
Euonymus, Spreading	<i>Euonymus kiautschovicus</i>
Euryops	<i>Euryops pectinatus</i>
Evergreen, Fransher; Silver queen; Treubii ribbon	<i>Aglanoema commutatum</i>
Evergreen, Painted	<i>Aglanoema crispum</i>
Fatshedera	<i>Fatshedera lizei</i>
Fern, Desert tree	<i>Lysiloma thornberii</i>
Fern, Leatherleaf	<i>Runohra adiantiformis</i>
Fern, Shield	<i>Polystichum capense</i>
Fern, Sword	<i>Nephrolepis exaltata</i>
Fig, Creeping	<i>Ficus repens</i>
Fig, Exotica weeping	<i>Ficus benjamina</i>
Fig, Trailing hottentot*	<i>Carpobrotus chilensis*</i>
Fir, Balsam*	<i>Abies balsamea*</i>
Fir, Concolor	<i>Abies concolor</i>
Fir, Douglas	<i>Pseudotsuga menziesii</i>
Fir, Noble	<i>Abies procera</i>
Firethorn	<i>Pyracantha graberi</i>
Firethorn, Mojave	<i>Pyracantha koidzumii x coccinea</i>
Firethorn, Scarlet; Lalandei	<i>Pyracantha coccinea</i>
Firethorn, Variegated	<i>Pyracantha angustifolia</i>
Firewheel tree	<i>Stenocarpus sinuatus</i>
Forsythia intermedia	<i>Forsythia intermedia</i>
Forsythia sp.	<i>Forsythia sp.</i>
Forsythia, Weeping	<i>Forsythia suspensa</i>
Gable Hybrid	<i>Rhododendron "Gable Hybrid"</i>
Gardenia, Dwarf	<i>Gardenia jasminoides</i>
Gardenia, Tahitian	<i>Gardenia taitensis</i>
Gay feather	<i>Liatris spicata</i>
Gazania gold rush	<i>Gazania splendens</i>
Gazania uniflora leucoleana	<i>Gazania uniflora leucolaena</i>
Geranium	<i>Pelargonium domesticum</i>
Geranium, Ivy	<i>Pelargonium peltatum</i>
Geranium, Smash hit red*	<i>Pelargonium x hortorum*</i>
Gimlet, Narrow-leaf	<i>Eucalyptus spathulata</i>

(cont. on next page)







