



For Outdoor Use Around
the Home Only

Spectracide®



TRIAZICIDE® INSECT KILLER

ONCE & DONE!™

Insecticida concentrado₂

CONCENTRATE₂

Kills 180+ Listed Insects by Contact

Kills Above & Below Ground

Fast-Acting Formula



Active Ingredient
Gamma-Cyhalothrin 0.08%
Other Ingredients 99.92%

†For terms of guarantee, see back panel.

*Argentine, southern, field, Allegheny
mound, Florida carpenter, cornfield,
honey, pavement, nuisance.

For lawns & landscapes

KEEP OUT OF REACH OF CHILDREN

CAUTION See back booklet for additional
precautionary statements.

Net 32 fl oz (1 qt/946 mL)

MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS

PRECAUCIÓN Consulte el folleto en la parte posterior para
conocer las declaraciones preventivas adicionales.

17-13877

mosca de la nuez (adulta), hormigas, chinche, chinche hedionda, chinches, insecto occidental de la semilla de la conifera (leaffooted bug), áfido del nogal, hickory shuckworm, barrenador de la nuez pecana, pecan weevil, pulgón amarillo del pacanero, pecan spittlebug, filoxera del pacanero.

CANTIDAD DE DÍAS QUE SE DEBEN ESPERAR PARA LA ÚLTIMA APLICACIÓN Y LA COSECHA	
ÁRBOLES FRUTALES Y DE NUECES	IPC
Manzano, Manzano silvestre, Nispero, Mayhaw, Peral asiático, Peral, Membrito	21
Chabacano, Cerezo dulce, Cerezo agrio, Nectarina, Durazno, Ciruelo (Chickasaw, Damson, Japonés), Ciruela y Ciruelas pasa	14
Almendra, Nuez de haya, Nuez de Brasil, Nogal blanco, Nuez de la India, Castaña, Chinquapín, Avellana (hazelnut), Nogal americano, Macadamia, Nogal (negro e inglés), y Nuez pecana	14
IPC = Intervalo previo a la cosecha	

ALMACENAMIENTO Y ELIMINACIÓN

Almacenamiento del pesticida: Guarde en un lugar fresco y seco, lejos del calor y del fuego.

Eliminación del pesticida: Si está vacío: Envase no rellenable. No vuelva a usar ni rellenar este envase. Ofrezcalo para reciclar, si existe esa opción. Si está parcialmente vacío: Llame a su servicio de eliminación de desechos más cercano para obtener instrucciones de eliminación. Nunca tire el producto restante por ningún drenaje interno ni externo.

DECLARACIONES PREVENTIVAS

Riesgos para los seres humanos y los animales domésticos

PRECAUCIÓN. Nocivo si se ingiere. Lávese muy bien las manos con agua y jabón después de manipularlo y antes de comer, beber, masticar chicle, usar tabaco o usar el baño.

PRIMEROS AUXILIOS

Si se ingiere: Llame inmediatamente a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento. Si la persona puede tragar, haga que beba poco a poco un vaso de agua. No induzca el vómito a menos que así se lo indique un centro de control de envenenamientos o un médico. No administre nada por la boca a una persona que haya perdido el conocimiento.

Cuando llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento, tenga a la mano el envase o la etiqueta del producto.

Riesgos para el medio ambiente

Este producto es extremadamente tóxico para los peces. Para proteger el medio ambiente, no permita que el pesticida entre o escorra en alcantarías pluviales, zanjas de drenaje, cunetas ni agua de superficie. Aplicar este producto cuando el clima está tranquilo y no se pronostica lluvia dentro de las siguientes 24 horas asegurará que el pesticida no vuele con el viento o se lave con la lluvia del área tratada. Enjuagar el equipo de aplicación sobre el área tratada ayudará a evitar el escurrimiento a masas de agua o sistemas de drenaje. La deriva y el escurrimiento desde las áreas tratadas pueden resultar peligrosos para los organismos acuáticos de las zonas vecinas.

Este producto es extremadamente tóxico para las abejas que estén expuestas al tratamiento directo o a los residuos que queden en cultivos o malezas en flor. No aplique este producto ni permita que pase a los cultivos o malezas en flor en caso que haya abejas volando por el área tratada.

Spectracide



READ ENTIRE LABEL
BEFORE USE.

PEEL HERE
RESEALABLE LABEL

ALTO. LEA TODA LA
ETIQUETA ANTES DE
USAR.

ABRA AQUÍ
ETIQUETA
RESELLABLE

TRIAZICIDE® INSECT KILLER

— **ONCE&DONE!®** —

CONCENTRATE₂

Insecticida concentrado



Covers up to
5,120 sq ft of Lawn



Non-Staining*



Protects Lawns, Vegetables, Fruit & Nut
Trees, Roses, Flowers, Trees, Shrubs

PRODUCT FACTS

WHAT IT DOES:

Kills grubs, ants*, fleas, chinch bugs, mosquitoes, sod webworms, aphids, beetles, whiteflies, mites, deer ticks, Japanese beetles, European crane flies and other listed insects.

*Argentine, southern, field, Allegheny mound, Florida carpenter, cornfield, honey, pavement, nuisance.

WHERE TO USE:

Outdoors

QUESTIONS & COMMENTS:

Call 1-800-917-5438 or visit our website
at www.spectracide.com

*NOTE: This product is non-staining to most home siding depending on age and cleanliness. However, before using in areas where the spray may contact home siding (vinyl siding in particular), test in an inconspicuous area and recheck in a few hours. Do not use if any staining is observed.

'GUARANTEED RESULTS OR YOUR MONEY BACK'

If you are not satisfied with this product for any reason, send your written request for a refund to the address below, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

NOTA: Hasta el grado que sea compatible con la legislación pertinente, el comprador asume toda la responsabilidad para seguridad y uso que no esté de acuerdo con las instrucciones.

Distributed by Spectrum Group
Div. of United Industries Corp.
PO Box 142642, St. Louis, MO 63114-0642

EPA Reg. No. 9688-277-8845

EPA Est. Nos. 9688-MO-1 (U), 39609-MO-1 (BM)

Circled letter is first letter of lot number.

17-13299

© 2009 UIC

PROOF OF PURCHASE

FPO UPC

0 71121 95829 7

STOP. READ ENTIRE LABEL BEFORE USE.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



Do not allow children or pets into the treated area until dry.

- Do not water the treated area to the point of runoff.
- Do not make applications during rain.
- All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:
 - Treatment to soil or vegetation around structures
 - Applications to lawns, turf and other vegetation
 - Applications to building foundations, up to a maximum height of 3 ft. Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors and eaves) are limited to spot and crack-and-crevice applications only.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

In New York State, this product may not be applied to lawns within 100 ft of a coastal marsh, or stream that drains directly into a coastal marsh.

HOW TO MIX AND APPLY

- STEP 1: **Determine size of spraying job.** For lawns and vegetables, measure the area to be sprayed. For trees, shrubs and flowers, apply as a thorough cover spray. Make applications when insects first appear. Applications should be repeated only as directed to maintain control.
- STEP 2: **Use a clean sprayer.** Carefully measure and mix the amount of product and water as indicated in the directions. It is always a good idea to wear gloves when handling pesticides. Kitchen utensils such as measuring cups and measuring spoons should not be used for food purposes after use with pesticides.
- STEP 3: **Spray as directed.** Thorough coverage is important.
- STEP 4: Flush sprayer with clean water after each use.

See directions for list of use sites, plants and insects controlled. Use the rate given in the directions for the type of insect you want to control on the site or plant you are spraying.

MEASUREMENT CONVERSIONS

- 1 Tbsp = 3 tsp
- 1 fl oz = 6 tsp or 2 Tbsp
- 6 fl oz = ¾ cup
- 1½ Tbsp in 3 gallons of water = 1½ tsp in 1 gallon of water
- 2 Tbsp in 3 gallons of water = 2 tsp in 1 gallon of water

*NOTE: This product is non-staining to most home siding depending on age and cleanliness. However, before using in areas where the spray may contact home siding (vinyl siding in particular), test in an inconspicuous area and recheck in a few hours. Do not use if any staining is observed.

LAWNS

UNDERGROUND INSECTS/THATCH-INFESTING INSECTS including blue grass billbug (adult), black turgrass *Ataenius* (adult), chiggers, chinch bugs, white grubs (Japanese beetle, European chaffer, southern chaffer), dung beetle and hyperodes weevils (adult), European crane flies and mole crickets (nymphs and young adults). Mix 1.5 fl oz (3 Tbsp) in 1 gallon of water to treat 120 sq ft. For subsurface insect control (mole crickets, grubs), water treated area an additional ¼ to ½ inch of water for optimum results. Do not apply more than seven times per year per location.

SURFACE INSECTS including ants*, armyworms, cockroaches, crickets, cutworms, earwigs, fleas, lawn moths (sod webworms), millipedes, mosquitoes, palmto bugs, sowbugs, mites, spittlebugs and waterbugs. Thoroughly wet down grass a few hours before applying. Mix 1.5 fl oz (3 Tbsp) in 1 gallon of water to cover 240 sq ft. Delay additional watering or mowing for 24 hours after application for optimum control of surface insects. For heavy infestations or to kill lone star ticks, brown dog ticks, American dog ticks and deer ticks (which may transmit Lyme

Disease) use rate can be increased to 6 fl oz/1,000 sq ft. Do not apply more than 14 times per year per location. **WHEN TO TREAT:** Treat the following pests during the months indicated or when first noticed.

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
			Billbugs				Grubs				
			Dung beetle grubs								
			Hyperodes weevils								
Treat ants* and other insects when they first appear or you notice lawn damage.											

*Argentine, southern, field, Allegheny mound, Florida carpenter, cornfield, honey, pavement, nuisance.

PESTS OUTSIDE THE HOME

Mix 1.5 fl oz (3 Tbsp) in 1 gallon of water and apply as a continuous band of insecticide around building foundation and around windows, doors, eaves, vents and other areas to greatly reduce the potential for entry by crawling pests. For easier application, remove debris and leaf litter from next to the foundation, cut back vegetation and branches that touch the foundation, and move or rake back rocks, dead mulch or other potential pest harborage next to the foundation. Apply a 3-foot band around the structure and upward along the foundation to 3 feet and around windows, doors and roof overhangs. Apply as a coarse spray to thoroughly and uniformly wet the foundation and/or band area so that the insecticide will reach the soil or thatch level where pests may be active. One gallon will treat 240 sq ft of surface area. Repeat application every eight weeks.

OUTDOOR SURFACE USE: For control of ants*, centipedes, cockroaches, crickets, fleas, millipedes, mosquitoes, palmto bugs, scorpions, sowbugs, pillbugs, spiders, deer ticks and waterbugs. Mix 1.5 fl oz (3 Tbsp) in 1 gallon of water and apply as a residual treatment to ornamental plants next to foundations of buildings and to surfaces of buildings, porches, screens, window frames, eaves, patios, garages and other similar areas where these pests are active. Repeat application every eight weeks.

*Argentine, southern, field, Allegheny mound, Florida carpenter, cornfield, honey, pavement, nuisance.

ORNAMENTAL TREES, SHRUBS AND FLOWERS

For use on ornamental trees, shrubs and flowers such as but not limited to: Ageratum, Arborvitae, Arizona Cypress, Ash, Aster, Azaleas, Begonia, Birch, Boxwood, Box Elder, Camellias, Camellians, Cherry (ornamental), Chrysanthemums, Non-bearing Citrus, Coleus, Common Ninebark, Conifers, Douglas Fir, Elm, English Ivy, Euonymus, Exacum, Weeping Fig, Fir, Gladioli, Gold Belis, Hawthorn, Holly, Honeysuckle, Hypoestes, Ivy, Juniper, Lilac, Locust, Maple, Marigold, Mimosa, Mock Orange, Nannyberry, Oak, Orchid, Palm, Pansy, Pea Shrub, Petunia, Philodendron, Pine, Ornamental Plum, Poinsettia, Poplar, Tulip Poplar, Rhododendron, Roses, Snapdragon, Snowberry, Spruce, Taxus, Willow, Zinnia.

To control ants*, aphids, armyworms, azalea caterpillars, bagworms, black vine weevil (adult), box elder bugs, budworms, California oakworm, cankerworms, cockroaches, crickets, cutworms, eastern tent caterpillar, elm leaf beetles, European sawfly, leaf webworm, flea beetles, forest tent caterpillar, gypsy moth larvae, Japanese beetles (adults), June beetles (adults), lace bugs, leaf-feeding caterpillars, leafhoppers, leafminers (adults), leaf rollers, leaf skeletonizers, midges, oleander moth larvae, pillbug, pine sawfly, pine shoot beetles, pinetip moths, plant bugs, root weevil, sawfly, scale insects (crawlers), spiders, spittlebugs, striped beetles, striped oakworm, thrips, tip moth, tussock moth larvae: Mix 0.75 fl oz (1½ Tbsp) in 1 gallon of water.

To control broadmites, brown softscale, California redscale (crawler), clover mites, mealybugs, pineneedle scale (crawler), spider mites and whiteflies: Mix 1.5 fl oz (3 Tbsp) in 1 gallon of water.

*Argentine, southern, field, Allegheny mound, Florida carpenter, cornfield, honey, pavement, nuisance.

USE TIPS

- For best results, apply early before insect populations are high.
- For scale control, treat entire plant, including stem, trunks and twigs.
- For bagworm control, apply when bagworm larvae begin to hatch.
- For best results, apply to both sides of plant leaves and all plant surfaces where insects are present.

VEGETABLES

Mix 1.5 fl oz (3 Tbsp) in 1 gallon of water and apply to 300 sq ft. Apply when insects appear or when damage occurs. Spray to wet all infested plant surfaces.

Do not apply more than five times per growing season to corn. For all other crops, do not apply more than nine times per growing season.

For use on Broccoli, Brussels Sprouts, Cabbage, Cavalo Broccoli, Cauliflower, Chinese Broccoli, Chinese Cabbage, Chinese Mustard and Kohlrabi to control: alfalfa looper, cabbage looper, imported cabbageworm, southern cabbageworm, cutworm, cabbage webworm, diamondback moth, armyworm, beet armyworm, fall armyworm, yellow-striped armyworm, corn earworm, flea beetle, Japanese beetle (adult), grasshopper, leafhopper, plant bug, including lygus bug, stink bug, meadow spittlebug, aphid, whitefly, thrips, spider mite.

For use on Corn (sweet and pop) to control: cutworm, western bean cutworm, corn earworm, green cloverworm, meadow spittlebug, tobacco budworm, European corn borer, southwestern corn borer, stalk borer, hop vine borer, armyworm, beet armyworm, western corn rootworm (adult), northern corn rootworm beetle (adult), southern corn rootworm beetle (adult), Mexican corn rootworm beetle (adult), bean leaf beetle, cereal leaf beetle, Japanese beetle (adult), sap beetle (adult), stink bug, grasshopper, corn leaf aphid, English grain aphid, chinch bug, green bug. Sweet Corn only: southern armyworm, western bean cutworm, tarnished plant bug, aster leafhopper, aphid, spider mite, corn silkfly (adult).

For use on Onion (bulb) and Garlic to control: cutworm, seedcorn maggot (adult), onion maggot (adult), leafminer (adult), armyworm, onion thrips, tobacco thrips, western flower thrips, flower thrips, aphid, plant bug, stink bug.

For use on Tomato, Tomatillo, Peppers (bell and nonbell), Eggplant, Ground Cherry and Pepino to control: cabbage looper, cutworm spp., hornworm, tomato fruitworm, tobacco budworm, tomato pinworm, beet armyworm, southern armyworm, yellow-striped armyworm, fall armyworm, European corn borer, leafminer, Colorado potato beetle, flea beetle, grasshopper, leafhopper, aphid, whitefly, meadow spittlebug, stink bug, plant bug, stalk borer, blister beetle, Japanese beetle (adult), vegetable weevil, cucumber beetle (adult).

For use on Asparagus Beans, Snap Beans, Wax Beans, Black-Eyed Peas, Garden Peas, Green Peas and Sugar Snap Peas to control: cutworm spp., green cloverworm, imported cabbageworm, saltmarsh caterpillar, velvetleaf caterpillar, Mexican bean beetle, corn earworm, painted lady butterfly (larvae), European corn borer, looper spp., western bean cutworm, armyworm, yellow-striped armyworm, western yellow-striped armyworm, bean leafskeletonizer, webworm spp., leafminer spp., alfalfa caterpillar, stalk borer, cucumber beetle (adult), corn rootworm beetle (adult), flea beetle (adult), blister beetle, bean leaf beetle, Japanese beetle (adult), leafhopper spp., fleas hoppers spp., three-cornered alfalfa hoppers, meadow spittlebug, stink bug spp., plant bug spp., grasshopper, thrips spp., aphid spp.

For use on Lettuce (head and leaf) to control: alfalfa looper, cabbage looper, imported cabbageworm, cutworm spp., saltmarsh caterpillar, green cloverworm, armyworm, beet armyworm, fall armyworm, corn earworm, European corn borer, flea beetle spp., Japanese beetle (adult), vegetable beetle (adult), grasshopper, leafhopper, plant bug, stink bug, meadow spittlebug.

NUMBER OF DAYS TO WAIT BETWEEN THE LAST APPLICATION AND HARVEST		
VEGETABLES		PHI
Broccoli, Brussels Sprouts, Cabbage, Cavalo Broccoli, Cauliflower, Chinese Broccoli, Chinese Cabbage, Chinese Mustard, Kohlrabi, Lettuce (head and leaf)		1
Tomato, Tomatillo, Peppers (bell and nonbell), Eggplant, Ground Cherry and Pepino		5
Onions (bulb), Garlic		14
Corn (sweet and pop)		21
Asparagus Beans, Snap Beans, Wax Beans, Black-Eyed Peas, Garden Peas, Green Peas and Sugar Snap Peas		7
Lettuce (leaf and head)		1
PHI = Pre-Harvest Interval		

FRUIT AND NUT TREES

Mix 1.5 fl oz (3 Tbsp) in 1 gallon of water and apply to 300 sq ft. Apply when insects appear or when damage occurs. Spray to wet all infested plant surfaces.

For fruit trees do not apply more than nine times per year, and only apply five times after blooms

have formed. For nut trees do not apply more than six times per year, and only apply five times after blooms have formed.

For use on Apple, Crabapple, Loquat, Mayhaw, Oriental Pear, Pear and Quince to control: **leafhopper spp.**, **codling moth**, **tufted apple budworm**, **Oriental fruit borer**, **lesser appleworm**, **green fruitworm**, **tent caterpillar spp.**, **tentiform leaf miner spp.**, **apple maggot (adult)**, **pear sawfly**, **plum curculio**, **Japanese beetle**, **plant bug**, **stink bug**, **leafhopper**, **periodical cicada**, **apple aphid**, **rosy apple aphid**, **pear psylla**, **San Jose scale (on fruit only)**.

For use on Apricot, Sweet Cherry, Tart Cherry, Nectarine, Peach, Plum (Chickasaw, Damson, Japanese), Plumcot and Prune to control: **leafhopper spp.**, **Oriental fruit moth**, **Oriental fruit moth**, **peachtree borer spp.**, **green fruitworm**, **tent caterpillar spp.**, **American plum borer**, **cherry fruit fly spp. (adult)**, **plum curculio**, **rose chafer**, **Japanese beetle**, **plant bug**, **stink bug**, **leafhopper spp.**, **periodical cicada**, **black cherry aphid**.

For use on Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Macadamia Nut, Walnut (Black and English) and Pecan to control: **leafroller spp.**, **navel orangeworm**, **codling moth**, **filbertworm**, **peach twig borer**, **walnut husk fly spp. (adult)**, **ants**, **plant bug**, **stink bug**, **chinch bug**, **leaffooted bug**, **walnut aphid**, **hickory shuckworm**, **pecan casebearer spp.**, **pear weevil**, **pecan aphid spp.**, **pecan spittlebug**, **pecan phylloxera spp.**

NUMBER OF DAYS TO WAIT BETWEEN THE LAST APPLICATION AND HARVEST	
FRUIT AND NUT TREES	PHI
Apple, Crabapple, Loquat, Mayhaw, Oriental Pear, Pear, Quince	21
Apricot, Sweet Cherry, Tart Cherry, Nectarine, Peach, Plum (Chickasaw, Damson, Japanese), Plumcot and Prune	14
Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Macadamia Nut, Walnut (Black and English) and Pecan	14
PHI = Pre-Harvest Interval	

STORAGE AND DISPOSAL

Pesticide Storage: Store in a cool dry area away from heat or open flame.

Pesticide Recycling: If empty. Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If **partly filled**: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If **Swallowed**: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person. Have the product container or label with you when calling a Poison Control Center or doctor for treatment advice.

Environmental Hazards

This product is extremely toxic to fish. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

ALTO. LEA TODA LA ETIQUETA ANTES DE USAR.

MODO DE EMPLEO

Se considera una violación a la ley federal usar este producto de una manera que no sea la indicada en la etiqueta.



No permita que los niños ni las mascotas entren en el área tratada hasta que se seque.

- No riegue el área tratada hasta el punto de escurrimiento.
- No realice aplicaciones durante lluvias.
- Todas las aplicaciones en espacios exteriores deben limitarse a tratamientos localizados o para tratar grietas y rajaduras exclusivamente, salvo los siguientes usos permitidos:
 - Tratamiento del suelo o vegetación alrededor de estructuras
 - Aplicaciones a céspedes y otra vegetación
 - Aplicaciones a techos de edificios, hasta una altura máxima de 3 pies. Aparte de las aplicaciones a techos de edificios, todas las aplicaciones en superficies exteriores a superficies impermeables como aceras, caminos de entrada de automóviles, patios, porches y superficies estructurales (como ventanas, puertas y aleros) se limitan a aplicaciones localizadas y para el tratamiento de grietas y rajaduras exclusivamente.
- Se prohíbe la aplicación directa a alcantarillas o drenajes, así como a cualquier área similar a una cuneta en la que pueda producirse el drenaje a alcantarillas, alcantarillas pluviales, masas de agua o hábitats acuáticos. No permita que el producto entre en ningún drenaje durante o después de su aplicación.

En el Estado de Nueva York, no se podrá aplicar este producto en áreas de césped ubicadas a una distancia de hasta 100 pies de un pantano costero o de un arroyo que desemboque directamente en un pantano costero.

CÓMO MEZCLAR Y APLICAR

PASO 1: **Determine el tamaño de la tarea de rocío a realizar.** Para céspedes y hortalizas, mida el área que se va a rociar. En árboles, arbustos y flores, aplique como un rocío que cubra por completo. Realice aplicaciones en cuanto aparezcan los insectos por primera vez. Sólo se debe repetir el tratamiento de acuerdo a las indicaciones, con la finalidad de mantener la plaga bajo control.

PASO 2: **Use un rociador limpio.** Mida cuidadosamente y mezcle la cantidad de producto y de agua como se indica en las instrucciones. Es siempre recomendable usar guantes al manipular pesticidas. **No deben utilizarse utensilios de cocina tales como tazas o cucharas de medida para preparar alimentos después de usarlos con pesticidas.**

PASO 3: **Rocíe** como se indica. Es importante que la cobertura sea completa.

PASO 4: Enjuague bien el rociador con agua limpia después de cada uso.

Consulte las instrucciones para obtener una lista de los lugares y las plantas donde se usa, así como de los insectos controlados. Use la dosis que se indica en las instrucciones para el tipo de insecto que desea controlar en el lugar o planta que va a rociar.

CONVERSIONES DE MEDIDAS

1 cda. soperas = 3 ctdas.

1 onza líquida = 6 ctdas. ó 2 cdas. soperas

6 onzas líquidas = $\frac{3}{4}$ tda.

1 $\frac{1}{2}$ cda. soperas en 3 galones de agua = 1 $\frac{1}{2}$ ctdas. en 1 galón de agua

2 cdas. soperas en 3 galones de agua = 2 ctdas. en 1 galón de agua

NOTA: Este producto en general no mancha la mayoría de los revestimientos del hogar, dependiendo de la antigüedad y limpieza de los mismos. No obstante, antes de utilizarlo en áreas dónde el rocío pueda entrar en contacto con el revestimiento del hogar (en particular con revestimientos vinílicos), pruébelo en un lugar que no sea muy visible y verifique su efecto unas horas más tarde. No lo utilice si observa manchas.

CÉSPEDES

INSECTOS SUBTERRÁNEOS/INSECTOS QUE INFECTAN EL QUINCHADO entre los que se incluyen gorgojos de pasto azul anual (adultos), escarabajo negro del césped del género *Ataenius* (adulto), ácaros del género *Tribolium*, chinches, larvas blancas (escarabajo japonés, gusano blanco europeo (European chafer), escarabajo marrón (Southern chafer)), escarabajos del estiércol y gorgojos

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If Swallowed: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person. Have the product container or label with you when calling a Poison Control Center or doctor for treatment advice.

Environmental Hazards

This product is extremely toxic to fish. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

Questions & Comments? Call 1-800-917-5438 or visit our website at www.spectracide.com

DECLARACIONES PREVENTIVAS

Riesgos para los seres humanos y los animales domésticos

PRECAUCIÓN. Nocivo si se ingiere. Lávese muy bien las manos con agua y jabón después de manipularlo y antes de comer, beber, masticar chicle, usar tabaco o usar el baño.

PRIMEROS AUXILIOS

Si se ingiere: Llame inmediatamente a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento. Si la persona puede tragar, haga que beba poco a poco un vaso de agua. No induzca el vómito a menos que así se lo indique un centro de control de envenenamientos o un médico. No administre nada por la boca a una persona que haya perdido el conocimiento.

Cuando llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento, tenga a la mano el envase o la etiqueta del producto.

Riesgos para el medio ambiente

Este producto es extremadamente tóxico para los peces. Para proteger el medio ambiente, no permita que el pesticida entre o escurra en alcantarillas pluviales, zanjas de drenaje, cunetas ni agua de superficie. Aplicar este producto cuando el clima está tranquilo y no se pronostique lluvia dentro de las siguientes 24 horas asegurará que el pesticida no vuele con el viento o se lave con la lluvia del área tratada. Enjuagar el equipo de aplicación sobre el área tratada ayudará a evitar el escurrimiento a masas de agua o sistemas de drenaje. La deriva y el escurrimiento desde las áreas tratadas pueden resultar peligrosos para los organismos acuáticos de las zonas vecinas.

Este producto es extremadamente tóxico para las abejas que estén expuestas al tratamiento directo o a los residuos que queden en cultivos o malezas en flor. No aplique este producto ni permita que pase a los cultivos o malezas en flor en caso que haya abejas volando por el área tratada.

NOTA: Hasta el grado que sea compatible con la legislación pertinente, el comprador asume toda la responsabilidad para seguridad y uso que no esté de acuerdo con las instrucciones.

¿Preguntas o comentarios? Llame al 1-800-917-5438.

Distributed by Spectrum Group
Division of United Industries Corporation
PO Box 142642, St. Louis, MO 63114-0642

EPA Reg. No. 9688-277-8845

Circled letter is first letter of lot number.

EPA Est. Nos. 9688-MO-1 (U), 39609-MO-1 (BM)

17-13299

© 2009 UIC