



ferti·lome®



Weed-Out® Lawn Weed Killer

Contains TRIMEC® Herbicide

- Kills troublesome broadleaf weeds:
Dandelion, Chickweed, Clover, Henbit & many others

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

SEE BACK PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS
AND FIRST AID

Net Contents One Pint (16 FL. OZS.)

ACTIVE INGREDIENTS:

| | |
|---|---------|
| Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid | 5.45% |
| Dimethylamine salt of 2,4-dichlorophenoxyacetic acid | 5.88% |
| Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid) | 1.21% |
| OTHER INGREDIENTS: | 87.46% |
| TOTAL: | 100.00% |

See back panel for substatement of ingredient statement.

ferti•lome® WeedOut Lawn Weed Killer

15M-25-911FK

FIRST AID

| | |
|--------------------------------|---|
| If In Eyes: | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a Poison Control Center or doctor for treatment advice. |
| If Swallowed: | <ul style="list-style-type: none">• Call a Poison Control Center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a Poison Control Center or doctor.• Do not give anything by mouth to an unconscious person. |
| If On Skin Or Clothing: | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a Poison Control Center or doctor for treatment advice. |
| If Inhaled: | <ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a Poison Control Center or doctor for further treatment advice. |

Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment.

In the event of a medical or chemical emergency contact 
North America 1-800- 255-3924 or Worldwide Intl. + 01- 813-248-0585

Manufactured By:



230 FM 87

BONHAM, TEXAS 75418

EPA Reg. No. 2217-597-7401

EPA Est. No. 7401-TX-001

Visit Us At:

www.fertilome.com



PEEL FROM CORNER OF BOOKLET



STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Harmful if swallowed. Avoid contact with skin, eyes or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear long-sleeved shirt, long pants, socks, shoes and gloves. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drain systems.


This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift.

- 
- Keep people and pets out of the area during application.
 - Do not allow people or pets to enter the treated area until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Keep from freezing. Store in original container and place in locked storage area when not using.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down and indoor or outdoor drain.

WHEN TO USE

Spray at any time when Weeds are growing. For best results, apply in Spring when Weeds are small and growing rapidly. Apply in Fall to remove late germinating Weeds.

HOW TO USE

Hose-End Sprayer: Use 1 gallon of water for every 250 square feet to be sprayed. For concentrate sprayers, add only the required amount of **ferti-lome® Weed Out Lawn Weed Killer** listed in the rate chart depending on type of Grass and number of square feet to be treated. Do not add water to concentrate sprayers. For dilution sprayers, add the required amount of **ferti-lome® Weed Out Lawn Weed Killer** and fill sprayer with water to proper level. Spray uniformly.

Pressure-Sprayer: Use 1/2 gallon of water for every 250 square feet to be sprayed. Add the required amount of **ferti-lome® Weed Out Lawn Weed Killer** listed in the rate chart for the type of Grass to be treated and fill

sprayer with water to the proper level. Adjust spray nozzle to deliver coarse spray. Spray uniformly.

Spot-Treatment: If hardy Weeds persist in Sensitive Grasses such as St. Augustine, Centipedegrass, Bermudagrass, Buffalograss or Bentgrass, spot treat individual Weeds using 1 teaspoon of **ferti-lome® Weed Out Lawn Weed Killer** in 1 quart of water (equivalent to 4 teaspoons per gallon of water.) Minimize contact of Grass.

**RATES OF ferti-lome® Weed Out Lawn Weed Killer
FOR VARIOUS LAWN GRASSES**

| Gallons Of Water Spray | | Area to Treat (Sq. Ft.) | St. Augustine- grass, Centipedegrass (fl. oz.) | Bermudagrass, Buffalograss Bentgrass (fl. oz) | Bluegrass Zoysia- grass, Fescue (fl. oz) |
|---------------------------|---------------------|----------------------------------|---|--|--|
| Hose End Sprayer | Pressure Sprayer | | | | |
| 1 | 1/2 | 250 | 1/2 | 2/3 | 2/3 to 1 |
| 4 | 2 | 1,000 | 2 | 2 2/3 | 2 2/3 to 4 |
| 8 | 4 | 2,000 | 4 | 5 1/3 | 5 1/3 to 8 |
| 20 | 10 | 5,000 | 10 | 13 1/3 | 13 1/3 to 20 |
| 32 | 16 | 8,000 | 16 | 21 1/3 | 21 1/3 to 32 |

EQUAL MEASURES

- 1 fluid ounce (fl. oz.) = 2 tablespoons (Tbs.) or 6 teaspoons (tsp.)
2 fluid ounces = 4 tablespoons 4 fluid ounces = 1/2 cup
8 fluid ounces = 1 cup 16 fluid ounces = 1 pint (pt.)
32 fluid ounces = 1 quart (qt.)


HOW TO USE IN THE LAWN WEED KILLER APPLICATOR

1. Connect sprayer nozzle to garden hose; remove safety cap from container and insert sprayer unit, securing tightly.
2. Turn on water; place finger over hole on right side of nozzle and product will automatically be dispersed.

3. Spray evenly and uniformly over area requiring application. (Check application guide to determine the correct area to be covered per ounce.)

| APPLICATION GUIDE | | | |
|--------------------------|--|--|---|
| Fluid Ounces | Recommended Coverage Area Per Ounce | | |
| | Sensitive Grasses Such As Centipedegrass And St. Augustinegrass (sq. ft.) | Bermudagrass, Bentgrass, And Buffalograss (sq. ft.) | Bluegrass, Fescue, And Zoysiagrass (sq. ft.) |
| 1 | 500 | 375 | 250 |
| 2 | 1,000 | 750 | 500 |
| 3 | 1,500 | 1,125 | 750 |
| 4 | 2,000 | 1,500 | 1,000 |
| 5 | 2,500 | 1,875 | 1,250 |
| 6 | 3,000 | 2,250 | 1,500 |
| 7 | 3,500 | 2,525 | 1,750 |
| 8 | 4,000 | 3,000 | 2,000 |
| 9 | 4,500 | 3,375 | 2,250 |
| 10 | 5,000 | 3,750 | 2,500 |
| 11 | 5,500 | 4,125 | 2,750 |
| 12 | 6,000 | 4,500 | 3,000 |
| 13 | 6,500 | 4,875 | 3,250 |
| 14 | 7,000 | 5,250 | 3,500 |
| 15 | 7,500 | 5,625 | 3,750 |
| 16 | 8,000 | 6,000 | 4,000 |

Use Precautions: Do not apply this product through any type of irrigation system. Use only lawn sprayer that delivers coarse spray to reduce drift. Avoid spray drift to Vegetables, Flowers, Ornamental Plants. Do not spray roots of Trees and Ornamentals. Do not exceed specified dosages for any area; be particularly careful within dripline of Trees and other Ornamental species. Do not spray Carpetgrass, Dichondra, or Lawn where desirable Clovers are present. Do not apply this product to "Floritam" St. Augustine Grass in Florida. Do not apply to newly seeded Lawn Grasses until well established. Do not spray when air temperatures exceed 85°F. Seed can be



sown 3 to 4 weeks after application. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides. Failure to follow above precautions may result in injury.

Other Use Precautions:

- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays and oceans. Do not apply when windy. To prevent run-off, do not over water the treat area(s) to the point of run-off or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.

PRODUCT INFORMATION

Ferti-lome® Weed Out Lawn Weed Killer controls troublesome common Lawn Weeds, including Poison Ivy, in both Northern and Southern Grasses.

- BROADLEAF WEEDS -

| | | |
|---------------------------|---------------------|-------------------|
| Annual Yellow Sweetclover | Bull Thistle | Cudweed |
| Aster | Bullnettle | Curly Dock |
| Austrian Fieldcress | Burclover | Daisy, English |
| Bedstraw | Burdock | Daisy Fleabane |
| Beggarticks | Burning Nettle | Daisy, Oxeye |
| Betony, Florida | Burweed | Dandelion |
| Bindweed, Field | Buttercup | Dichondra |
| Bird Vetch | Buttonweed | Dogfennel |
| Bitter Wintercress | Canada Thistle | Dollarweed |
| Bittercress, Hairy | Carolina Geranium | False Dandelion |
| Bitterweed | Carpetweed | Falseflax |
| Black-Eyed Susan | Catchweed Bedstraw | False Sunflower |
| Black Medic | Catnip | Fiddleneck |
| Black Mustard | Catsear | Florida Pusley |
| Blackseed Plantain | Chickweed, Common | Frenchweed |
| Blessed Thistle | Chickweed, Mouseear | Galinsoga |
| Bloodflower Milkweed | Chicory | Goldenrod |
| Blue Lettuce | Cinquefoil | Ground Ivy |
| Blue Vervain | Clover, Crimson | Gumweed |
| Bracted Plantain | Clover, Hop | Hairy Fleabane |
| Bristly Oxtongue | Clover, Red | Hawkweed |
| Broadleaf Dock | Clover, Strawberry | Healall |
| Broadleaf Plantain | Clover, Sweet | Heartleaf Drymary |
| Broomweed | Clover, White | Heathaster |
| Buckhorn | Cockle | Hedge Bindweed |
| Buckhorn Plantain | Cocklebur | Hedge Mustard |
| Bulbous Buttercup | Common Mullein | Henbit |
| | Creeping Jenny | Hoary Cress |

- BROADLEAF WEEDS - *(Continued)*

| | | |
|---------------------|---------------------|-----------------------|
| Hoary Plantain | Pennywort | Scotch Thistle |
| Hoary Vervain | Pepperweed | Sheep Sorrel |
| Horsenettle | Pigweed | Sherherdspurse |
| Jimsonweed | Pineywoods Bedstraw | Slender Plantain |
| Knawel | Plains Coreopsis | Smallflower Galinsoga |
| Knotweed | Plantain | Smooth Dock |
| Kochia | Poison Ivy | Smooth Pigweed |
| Lambsquarters | Poison Oak | Sorrel |
| Lespedeza | Pokeweed | Sowthistle |
| Mallow | Poorjoe | Spanishneedles |
| Matchweed | Prairie Sunflower | Speedwell |
| Mexicanweed | Prickly Lettuce | Spiny Amaranth |
| Milk Vetch | Prickly Sida | Spiny Cocklebur |
| Morningglory | Prostrate Knotweed | Spiny Sowthistle |
| Mouseear Hawkweed | Prostrate Pigweed | Spotted Catsear |
| Mugwort | Prostrate Spurge | Spotted Spurge |
| Musk Thistle | Prostrate Vervain | Spurweed |
| Mustard | Puncturevine | Stinging Nettle |
| Narrowleaf Plantain | Purslane, Common | Strawberry, India |
| Narrowleaf Vetch | Ragweed | Mock |
| Nettle | Red Sorrel | Tall Nettle |
| Orange Hawkweed | Redroot Pigweed | Tall Vervain |
| Oriental Cocklebur | Redstem Filaree | Tansy Ragwort |
| Oxalis | Rough Cinquefoil | Tansy Mustard |
| Parsley-Piert | Rough Fleabane | Tanweed |
| Parsnip | Russian Pigweed | Thistle |
| Pearlwort | Russian Thistle | Trailing Crownvetch |
| Pennycress | Scarlet Pimpernel | Tumble Mustard |



- BROADLEAF WEEDS - (Continued)

| | | |
|----------------------|-------------------|---------------------|
| Tumble Pigweed | Wild Four-O'Clock | Woodsorrel |
| Velvetleaf | Wild Garlic | Woolly Croton |
| Venice Mallow | Wild Geranium | Woolly Morningglory |
| Virginia Buttonweed | Wild Lettuce | Woolly Plantain |
| Virginia Creeper | Wild Marigold | Wormseed |
| Virginia Pepperweed | Wild Mustard | Yarrow |
| Wavyleaf Bullthistle | Wild Onion | Yellow Rocket |
| Western Clematis | Wild Parsnip | Yellowflower |
| Western Salsify | Wild Radish | Pepperweed |
| White Mustard | Wild Rape | and other Broadleaf |
| Wild Aster | Wild Strawberry | Weeds |
| Wild Buckwheat | Wild Sweet Potato | |
| Wild Carrot | Wild Vetch | |

LIMITED WARRANTY AND DISCLAIMER

The Manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This Limited Warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the Manufacturer or the Seller. Any damages arising from a breach of the Manufacturer's Warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss

or profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the Seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

This page left blank intentionally.

This page left blank intentionally.

THIS PRODUCT CONTAINS:

0.39 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 4.50%.

0.42 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 4.88%.

0.09 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 1.00%.

Isomer Specific by AOAC Methods.

Contains the single isomer form of Mecoprop-p.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

Manufactured By:



230 FM 87 • BONHAM, TX 75418

EPA Reg. No. 2217-597-7401

EPA Est. No. 7401-TX-001

Visit Us At: www.fertilome.com



Get the free mobile app at
<http://gettag.mobi>