



**ferti·lome®**



**BROWN PATCH**

**SUMMER PATCH**

**OPTIMUM  
DISEASE  
CONTROL**

**PINK SNOW MOLD**

**DOLLAR SPOT**

# **F-STOP™**

## **For Picture-Perfect, Disease-Free Turfgrass**

Provides systemic prevention and control of major lawn diseases including: anthracnose, brown patch, dollar spot, fusarium patch (pink snow mold), powdery mildew, red thread, rust, summer patch and other listed lawn diseases.

- For use on all types of home lawns.
- One application protects for up to 4 weeks.

**ACTIVE INGREDIENT:**  
myclobutanil: a-butyl-a-(4-chlorophenyl)-1-H-1,2,4 triazole-propanenitrile:..... 0.39%  
**OTHER INGREDIENTS:**..... 99.61%  
**Total:**..... 100.00%

This product contains 0.195 lb. of myclobutanil per 50 lb. bag.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL  
NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

**Net Weight 8 LBS. (3.63 kg)**

**Covers Up To  
2,000 Sq. Ft.**

**F-STOP™ 10770**



ferti·lome®



# F-STOP™

## For Picture-Perfect, Disease-Free Turfgrass

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long-sleeved shirt, long pants, and shoes and socks. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

**Notice:** Read the entire label. Use only according to label directions. Before using this product, read "Warranty Disclaimer," "Inherent Risks of Use," and "Limitation of Remedies" at end of Directions for Use. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994

**Agricultural Chemical:** Do not ship or store with food, feeds, drugs or clothing.

### FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice. Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. You may also contact the 1-800-992-5994 for emergency treatment information.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

**Not for use on turfgrass being grown for sale or other commercial use as sod, or for commercial seed productions, or for research purposes.**

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

**PESTICIDE STORAGE:** Keep away from fire and sparks. Store in a cool, dry, well-ventilated area.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container.

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

### HOW THIS PRODUCT WORKS

Ferti-lome® F-STOP™ is a systemic, protectant and curative Fungicide for the control of listed Diseases in established home lawns and ornamental turfgrass. Optimum disease control is achieved when this product is applied to established turfgrass in a regularly scheduled preventative program. Use this product in conjunction with turf management practices that promote good plant health and optimum disease control. The key to selecting a fungicide is the proper diagnosis of the organism causing the disease. Diagnostic kits, extension experts, or other identification methods should be used when developing disease control strategies.

### WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description of the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the Inherent Risks set forth below. Seller MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

### INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. To the extent allowed by law, all such risks shall be assumed by the buyer.

### LIMITATION OF REMEDIES

### HOW TO APPLY

Apply Ferti-lome® F-STOP™ uniformly over the turfgrass area using a properly calibrated drop or rotary-type spreader designed to apply granules. Before each application, calibrate the spreader according to the equipment manufacturer's directions. Check frequently to make sure equipment is operating properly and distributing product uniformly. A more uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. Avoid skips and excessive overlaps during application. Avoid the use of spreaders that apply this product in narrow rows or concentrated bands.

### WHEN TO APPLY

Reduce the interval between applications of this product when conditions are favorable for disease development. Unless otherwise specified, when disease pressure is high or when used as a curative, use higher rates of Ferti-lome® F-STOP™ and shorter application interval. Under light to moderate disease pressure apply this product at the low use rate and/or longer application intervals. To avoid pick-up, lightly irrigate treated areas soon after application. On short cut bentgrass (1/2 inch or less) when temperatures are above 80°F; apply only to dry foliage.

### HOW MUCH TO APPLY

Optimum disease control is achieved when Ferti-lome® F-STOP™ is applied in a preventative disease control program at a rate of 4 lb per 1,000 sq. ft. See the following table for specific application rates for various diseases. Under any circumstances, do not apply more than 46 lb of this product per 1,000 sq. ft. per year.

### SPREADER GUIDE ONE BAG WILL COVER UP TO 2,000 SQUARE FEET

SPREADER	SPREADER SETTINGS
Scotts/Republic Accugreen (Drop)	4 1/4
Scotts/Republic Speedy Green (Broadcast)	3 3/4
ferti-lome®/EarthWay Ev-N-Spred (Broadcast)	14

### TERMS AND CONDITION OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the Seller for a full refund of purchase price paid. Otherwise, use by the Buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

To the extent consistent with applicable law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by Buyer or user for product bought, or

2. Replacement of amount of product used.

To the extent allowed by law, Seller shall not be liable for the losses of damages resulting from the handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Seller or the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner..

Disease	ferti-lome® F-STOP® (lb/1,000 sq ft)	Application Interval/ Timing (Days)	Use Recommendations	Restrictions
Anthrachnose Red Thread Septoria Leaf Spot	4	14 - 21	Apply when conditions are favorable for Disease development.	Do not apply more than 46 pounds of product per 1,000 sq. ft. per year.  For Nassau and Suffolk Counties in New York State, use of this product is limited to 11.5 pounds per 1,000 sq. ft. per year (1.95 lb. active ingredient per acre).
Brown Patch		14	Begin applications when conditions are favorable for Disease development and before Disease symptoms are apparent.	
Copper Spot Zonate Leaf Spot			Apply when conditions are favorable for Disease development.	
Crown Rot Leaf Spot Melting-Out			Apply when conditions are favorable for Disease development. For crown rot, water in with 3 to 4 gallons of water per 1,000 sq. ft. to increase penetration to crown and roots.	
Dollar Spot		14 - 28	Apply when conditions are favorable for Disease development. Make no more than 3 consecutive applications for Dollar Spot control before rotating to a registered Fungicide with a different mode of action.	
Fusarium Blight		14 - 21	Apply when conditions are favorable for Disease development.	
Fusarium Patch (Pink Snow Mold)		Fall-Winter	Apply prior to snow cover.	
Gray Leafspot		14	Apply when conditions are favorable for disease development.	
Leaf Smuts			Apply in the fall after turfgrass enters dormancy and/or in the Spring prior to the initiation of growth.	
Necrotic Ring Spot		Spring: 28	Make applications on a preventative basis in early to mid-Spring.	
Necrotic Ring Spot		Fall: 28	Make 2 applications beginning in August before the turfgrass goes dormant.	
Powdery Mildew Rusts		14 - 28	Apply when conditions are favorable for Disease development.	
Summer Patch			Begin applications in the Spring when conditions are favorable for Disease development. Make 2 to 4 applications depending on recommendations from local Turfgrass Extension Experts. Water in with at least 3 to 4 gallons of water per 1,000 sq. ft. to increase penetration to crown and roots.	



Get the free mobile app at  
<http://gettag.mobi>

Manufactured by:



230 FM 87 • BONHAM, TEXAS 75418

EPA Reg. No. 62719-461-7401 EPA Est. No. 7401-TX-01

Visit Us At: [www.fertilome.com](http://www.fertilome.com)

