



Count on Patriot® for Broadleaf Weed Control.

Patriot® Selective Herbicide controls tough broadleaf weeds such as Russian thistle and mustards in wheat, barley, fallow, pastures and rangeland. It provides an economical post-patent alternative. With contact and residual control, Patriot burns down existing weeds and keeps late flushes down for four to six weeks. And its wide window of application gives you flexibility if application is delayed.

KEY PERFORMANCE BENEFITS

- Excellent post-emerge weed control
- Offers excellent crop safety
- Provides residual control for up to six weeks
- Wide application window
- Excellent tank-mix partner

EPA REG. NO.	228-391
ACTIVE INGR.	metsulfuron-methyl (60%)
FORMULATION	water dispersible granule
CHEM. FAMILY	sulfonylurea
HRAC NO.	2
MODE OF ACT.	ALS inhibitor
REI	4 hours
SIGNAL WORD	caution
RESTRICTED	no
PACKAGE SIZE	8 x 16 wt oz



Grow a better tomorrow.



KEY USES

Barley
Conservation reserve program (CRP) land
Fallow
Grain sorghum
Pastures
Rangeland
Triticale
Wheat

KEY WEEDS CONTROLLED

(See product label for complete list)

Annual marshelder
Bitter sneezeweed
Blackeyed-Susan
Broomweed, common
Bur buttercup
Carolina geranium
Chamomile; false, mayweed
Chickweed, common
Coast fiddleneck (*tarweed*)
Conical catchfly
Cowcockle
Curly dock
Dandelion
Dogfennel
Field pennycress (*fanweed*)
Filaree
Flixweed
Groundsel
Henbit
Kochia
Lambsquarters
Lettuce; miners, prickley
Marestail
Mustard; tansy, treacle,
tumble, wild
Pigweeds
Plains coreopsis
Purslane, common
Shepherd's-purse
Smallseed falseflax
Smartweeds
Sunflower, volunteer
Thistle, Russian
Waterpod
Woolly croton
Yarrow

USE SUMMARY

APPLICATION TIMING AND RATES

Wheat and Barley: Apply 1/10 oz/A. To avoid crop injury, do not apply during boot or early heading. For dryland wheat and barley (except durum or wampum variety), apply after the crop is in the 2-leaf stage, but before boot. For durum and wampum variety spring wheat, make applications after tillering, but before boot. Applications to durum and wampum varieties should be made in combination with 2,4-D. For irrigated wheat and barley, apply after the crop begins tillering, but before boot. First post-treatment irrigation should be delayed for at least 3 days after treatment and should not exceed one inch of water.

Pasture and Rangeland: Apply 1/10 to 4/10 oz/A as a broadcast spray. For spot treatments, use up to 1 oz per 100 gal of water.

Harvest Aid: Apply 1/10 oz/A in combination with 2,4-D (or Credit® 41 Extra / Credit® Xtreme / Razor® Pro) to aid in dry down of many broadleaf weeds. Make applications after the crop has reached the hard dough stage, but no later than 10 days before harvest.

Fallow: Patriot® may be used as a fallow treatment in the spring or fall when the majority of weeds have emerged and are actively growing. Apply at 1/10 oz/A.

Grain Sorghum: Apply 1/20 oz/A plus 1/4 lb active ingredient of 2,4-D amine per acre. Do not use a surfactant or crop oil.

See label for complete directions for use.



NUFARMPPOINT.COM