33030

# **Turf & Ornamental** Weed & Grass Stopper **Containing Dimension®**

Turt & Ornamental Weed & Grass Stopper



eed & Grass Stopper urf & Ornamental W

For Control Of Crabgrass And Control Or Suppression Of Many Other Annual Grass And Broadleaf Weeds In Ornamental Landscape Areas, **Established Lawns And Ornamental Turf** 

• Can Be Applied Up To 4 Weeks Later Than Other Preemergence **Crabgrass Preventers** 

**ACTIVE INGREDIENT:** 

OTHER INGREDIENTS: ...... 99.875% 

This product contains 0.0625 pound of the active ingredient dithiopyr per 50 pound bag.

# **KEEP OUT OF REACH OF CHILDREN** CAUTION

See Back/Side Panel For Additional **Precautionary Statements.** 

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes

- Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes
- For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns

In the State of New York, do not exceed 4.6 lb. per 1,000 sq. ft. per year of this product (equivalent to 0.25 lb. of active ingredient per acre). In the State of Florida, do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/ garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden.

# **Turf & Ornamental Weed & Grass Stopper**



NET WEIGHT 12 LBS. (5.44 KG)



Product Questions? 855-270-4776 moo.bleiv-id.www :tA eU tieiV FPA Est. No. 7401-TX-01 F047-E84-91723.00.62719-483-7401 230 FM 87 • BONHAM, TEXAS 75418



Manufactured By:

## :sətst2 bətinU ənt nintiw "snoipəA" əsU

Crabgrass control. ינ**קצצ צרמטלופ**ע וס קלאנא וע אסמן וסכפמסט מקצפת סע תופ מפצוגים ובולותו סו אפפת כסנתיסו סע נופפת וסג לסכנונופו

Suppression only

information. Hot Line Number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment

#### **SQRASAH JATNBMNORIVNB**

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. To protect

the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

unacceptable, return at once unopened. Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are

DIRECTIONS FOR USE

## It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**JASO9SID DNA 3DAROTS** 

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are

unacceptable, return at once unopened.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing. In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

Do not contaminate water, foodstuffs, feed or seed by storage and disposal.

Container Handling: Nonrefillable Container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partially filled: Call you local solid waste agency or 1-800-CLEANUP for Pesticide Storage: Store this product only in its original container in a dry, cool, secured storage area.

disposal instructions. Never place unused product down any indoor or outdoor drain.

#### USE DIRECTIONS FOR TURF

This turf product, containing Dimension® herbicide, can be applied up to 4 weeks later than other Crabgrass preventers and provides up to 4 months of Crabgrass control.

Hi-Vield® Turk & Ornamental Weed & Grass Stopper provides pre-emergent control of Crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seedling Crabgrass after germination and emergence. Postemergence control is limited to Crabgrass only, but is effective up to 4 weeks after Crabgrass has germinated. This generally corresponds to the time when Crabgrass selfer germination and emergence. Postemergence to not apply this product provides the crabgrass selfer germination and emergence. Postemergence control apply that product the time when Crabgrass selfer germination and the established turt grass. Do not

Florida Applications: Do not exceed 4 applications per year on St. Augustinegrass in central Florida. Do not apply more than 9.2 Ib of this product per 1,000 af t per application and no more than 27.6 Ib of this product/1000 sd ft per year (equivalent to 0.5 and 1.5 Ib of active ingredient per acre, respectively). apply this product later than 4 weeks after Crabgrass germination.

Apply this product only to turtgrass species listed on this label. The following turtgrass species are tolerant to this product when applied as directed.

| Centipedegrass<br>St. Augustinegrass<br>Zoysiagrass | Bahiagrass<br>Bermudagrass<br>Bahiagrass | Ryegrass, Perennial | Bentgrass, Creeping<br>Bluegrass, Kentucky<br>Fescue, Tail<br>Fescue, Tail |
|---|--|---------------------|--|
| SESSARD NOSAES-MRAW                                 |  | COOL-SEASON GRASSES |  |

#### **USE PRECAUTIONS FOR TURF**

This product should be applied only to lawns and turtgrasses that are well established. For newly established lawns, initial application of this product should be made only after turtgrasses have developed a vigorous root system, uniform stand, and received at least two mowings following seding, sodding, or sprigging. Exception: applied two weeks after the first sign of germination.

Use of this product on lawns and turfigrasses that are not well established, or weakened by stress due to mechanical operations, pests, excessive temperatures, or lack of moisture may result in turfigrasses injury.

- For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro aerification, and verticutting, should be done before application of this product. Do not apply this product until turgrass has recovered from the mechanical injury caused by these cultural practices.

Reseeding, Overseeding, or Sprigging: Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a single application of this product or within 76 weeks after a single application of this product or within 70 weeks after a single application of this product or within 76 weeks after a single application of this product or within 76 weeks after a single application of this product or within 76 weeks after a single application of this product or within 76 weeks after a single application of this product or within 76 weeks after a single application of the product or within 70 weeks after a single application of the product or within 70 weeks after a single application of the product or within 70 weeks after a single application of the product or within 70 weeks after a single application of the product or within 70 weeks after a single application of the product or within 70 weeks after a single application of the product or within 70 weeks after a single application of the product of the product or the product or within 70 weeks after a single application of the product of the product of the product or the product or within the product of the product or the product of t

#### OTHER USE PRECAUTIONS

• Do not allow people or pets to enter treated area until dust has settled Do not use clippings from treated turf for mulching around vegetables or fruit trees

#### **APPLICATION TIMING**

#### Spring Application

Control of Crabgrass: Hi-Yield® Turf & Ornamental Weed & Grass Stopper provides pre-emergent control of Crabgrass and control or suppression of other listed weeds when applied prior to their germination. In addition, this product provides postemergence control of seediing Crabgrass after germination and emergence. Postemergence corresponds to the time when Crabgrass arealing a real the 3-leaf stage of growth and first become visible in the established turf. This product will not control crabgrass at later growth stages when Crabgrass plants begin to tiller establings. Do not apply this product later than 4 weeks after germination and first become visible in the (spread). Properly timed applications will control existing Crabgrass and prevent further emergence of Crabgrass seedings. Do not apply this product later than 4 weeks after Crabgrass germination.

and broadleat weeds, this product must be applied as a preventative treatment prior to their germination and emergence. Only Crabgrass will be controlled with applications made up to 4 weeks after weed germination. Control or Suppression of Other Grass and Broadleaf Weeds: For control or suppression of other grass

#### Fall Application

applied in the fall, a sequential or second application of this product must be made the following spring to provide season-long control of Crabgrass. If this product is applied in the fall, do not reseed, overseed or sprig turgsrasses within 12 weeks after application. This product can also be applied from late summer through early fall (late August through November). When

#### Tips for Improved Weed Control

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.
- The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or
- irrigation. Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.

#### 21 91 **۲**۲ fertielome® /Earthway Ev-N-Spred (Broadcast) Scotts®/Republic Speedy Green (Broadcast) \$\t b ∜ε 8 %ε Scotts®/Republic Accugreen (Drop) %⊅ ₺ ₺ Þ 3.2 lbs. .2dl 7.4 SQI 1 Spreader Lbs. of product per 1,000 sq. ft.

usually be schieved by applying one-half of the required amount of product over the treatment area and then usually be schieved by applying one-half in a different direction (e.g., at a right angle to the previous direction). Avoid application rate under actual application conditions. The desired calibration setting may be marked or recorded for future reference. Apply this product uniformly over the treatment area. More uniformity of application can

best results, apply this product eventy and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules in narrow rows or concentrated bands. Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended to the manufacturer's directions. This product may be applied with drop or rotary-type spreaders designed to apply granular herbicides. For

ΑΡΡΓΙΟΑΤΙΟΝ ΙΝΑΤRUCTIONS ΑΝD ΕQUIPMENT In the state of New York, do not exceed 4.6 lb per 1000 sq. ft. per year of this product (equivalent to 0.25 Ib of active ingredient (ai) per acre).

Do not apply more than 9.2 Ib of this product per 1,000 sq. ft. per application and no more than 27.6 Ib of this product/061 ft. per acre, respectively).

(0.22-0.256 lb ai./acre)

<sup>++</sup>.ft.ps 000 f\dl 7.4 - 0.4

(0.174 lb ai/acre)

.ft .ps 000 f\dl S.S

sdfnom 4 - 6

------ Length of Crabgrass Control

Amount of Hi-Yield® Turf & Ornamental Weed & Grass Stopper to Apply (Pounds of product to apply per 1), or square feet of turigrass 1)

Recommended Use Rates

 $\triangleleft$ 

2

D

D.ª

 $^{+1}$  Use the higher rate in rate range in areas of high weed pressure or for increased length of control  $^{+1}$  Use the higher rate in rate range in areas of high weed pressure or for increased length of control  $^{+1}$ 

Spreader Settings

(0.22 - 0.5 lb ai/acre)

<sup>++</sup> .ft .ps 000 f\dl 2.e - 0.4

(0.22 - 0.5 lb ai/acre)

<sup>++</sup> .ff .ps 000 f\dl 2.e - 0.4

Control

Early Postemergence

**Y**auth

North

#### **TERMS AND CONDITIONS OF USE**

permitted by law otherwise, use by the Buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies. acceptable, return unopened package at once to the Seller for a full retund of purchase price paid. To the extent If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not

#### WARRANTY DISCLAIMER

purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MARCHANTABILITY OR FITUESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the

#### INHERENT RISKS OF USE

be assumed by Buyer. abnormal conditions (such as excessive rainfall, drought, fornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. All such risks shall other unintended consequences may result because of such tactors as use of the product contrary to label instructions (including conditions noted on the label, such as untavorable temperature, soil conditions, etc.). lt is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or

#### **LIMITATION OF REMEDIES**

To the extent permitted by law the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election,

:poie of the following:

Check equipment frequently to verify calibration and proper functioning.

streaking, skips, or overlaps during application.

Seten Replication Rates:

qqnos

North

Region .S.U to

(0.192 lb ai/acre)

ft ps 000 f\dl 7.5

(0.128 lb ai/acre)

ft ps 000 f\dl 4.2

2 - 3 months

1. Refund of purchase price paid by Buyer or user for product bought, or

2. Replacement of amount of product used.

To the extent permitted by law seller shall not be liable for losses or damages resulting from handling

or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law in no case shall Seller be liable for consequential or incitental changes or the extent permitted by law in the case shall Seller be liable for consequential

or incidental damages or losses. The terms of the Warranty Discialmer and Inherent Risks of Use above and this Limitation of Remedies cannot be

varied by any written or verbal statements or agreements. No employee or sales agent of Seller or the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Hi-Yield® is a registered trademark of Voluntary Purchasing Groups, Inc.

# Turf & Ornamental Weed & Grass Stopper

#### • Not for use on furt being grown for sale or other commercial use as sod, or for commercial seed production,

- for research purposes. or for research purposes. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or
- gardens or parks, or on golf courses or lawns. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental

of active ingredient (ai) per acre). In the state of New York, do not exceed 4.6 lb. per 1,000 sq. ft. per year of this product (equivalent to 0.25 lb.

is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back onto your lawn/garden. In the state of Florida, do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain

# PRECENTIONARY STATEMENTS PRECENTIONARY STATEMENTS PROFESSION ON SUMMUH OT SURASAH

#### CAUTION: Causes eye irritation • Harmful if inhaled

contact while handling this material may cause allergic reaction in some individuals. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin

#### **DIA TERIA**

for treatment advice. If In Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor

preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial

### Weeds Controlled or Suppressed

Hi-Yield® Turf & Ornamental Weed & Grass Stopper will provide control of Crabgrass and control or

|  | Pigweed, Redroot<br>Purslay-piett<br>Purslay-piett<br>Spurge, Common<br>Speedwall, Com<br>Spurge, Garden<br>Spurge, Garden<br>Spurge, Garden<br>Spurge, Prostrate<br>Spurge, Prostrate | Britercress<br>Britercress<br>Mustard<br>Medic, Black<br>Medic, Black | Goosegrass<br>Foxtall, Green<br>Foxtall, Yellow<br>Mikuyugrassf<br>& Perennial)<br>& Perennial)<br>Sandbur<br>Smutgrass | Barley<br>Bluegrass, Annual<br>Bluegrass, Amodh<br>Crabgrass, Large<br>Crabgrass, Large<br>Crabgrass, Large<br>Dallisgrass<br>Dallisgrass |
|--|---|---|---|---|
|  | BROADLEAF WEEDS   |   | SESSARD   |   |

#### Length of Control

your location. Use the following table to determine the amount of Hi-Yield® Turf & Ornamental Weed & applied according to label directions. The actual length of control will vary depending on use rate, weather conditions, condition of the turgitass, intensity of weed pressure, and the location or "Region" within the United States where this product is being applied. Refer to the map to determine the region of the U.S. that includes Hi-Vield® Turf & Ornamental Weed & Grass Stopper will provide up to 4 months of Crabgrass control when applied according to table directions. The scutal length of control will vary depending on use rate, weather the provided according to the scuta scuta scuta in the scuta sc