

**READY-TO-USE** 

# ATGEA RICK Corumes Granues

- Kills Insects in Lawns & Landscapes
- Kills Surface & Sub-Surface Insects



**NET WEIGHT 10 LBS (4.54 kg)** 

#### DEEP REACHING FORMULA

- Kills Grubs & Mole Crickets
- Vegetable Garden Insect Control

Treats up to 8,300 sq. ft.



NOOKUUT

## Ant, Flea & Tick Killer" Granules

342031

Subscript used is first letter of run code on container. 44616-MO-1C, EPA EST. NO. 59144-GA-1A, 67572-GA-1B, EPA REG. NO. 228-494-59144

P. O. Box 290 Madison GA 30650 Distributed by Gro Tec, Inc. KnockOut<sup>TM</sup> is a registered trademark of Gro Tec, Inc.

Do not contaminate water, food or feed by storage or disposal.

Call CHEMTREC (800) 424-9300. For Chemical Spill, Leak, Fire, or Exposure, For Medical Emergencies Only, Call (877) 325-1840.

any indoor or outdoor drain. available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down PESTICIDE CONTAINER DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if partial use, reseal open bag by folding top down and securing. place and avoid excess heat. Carefully open containers. After partial use replace lids and close tightly. For bags after PESTICIDE STORAGE: Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry

STORAGE AND DISPOSAL

When using other spreaders, calibrate the spreader settings based on the manufacturer's recommendations.    Days to wait				
Pests	to harvest	səldatapəV		
Ants, cutworms, crickets, earwigs, weevils grasshoppers, grubs, seed maggots, weevils and wireworms.	3 Days	Green Peas, Edible Pods (Sugar Snap Peas, Snow Peas) Green Beans (Wax Beans, Snap Beans) Dry Beans (Black-eyed Peas, Cowpea)		
Ants, armyworm, cutworms, crickets, earwigs, European corn borer, fall armyworm, grasshoppers, southwestern corn borer and weevils	1 Day	moO təəw2		
Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms.	sybū 7	Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Cavalo Broccoli, Chinese Broccoli, Chinese Cabbage, Kohlrabi		
Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms.	syad &	Cucurbit Vegetables Chayote, Citron melon, Cucumber, Edible Gourd Muskmelon (Cantaloupe, casaba, Crenshaw melon, Golden melon, Pineapple melon, Santa Claus melon, Snake melon) Pumpkin, Summer squash (Crookneck, Scallop, Straightneck, Zucchini) Winter squash (Butternut, Calabaza, Hubbard including Acorn and Spaghetti) Watermelon		
Ants, cutworms, crickets, earwigs, weevils gnasshoppers, grubs, seed maggots, weevils and wireworms.	sysd 7	Peppers		
Ants, cutworms, crickets, earwigs, grasshoppers, grubs, seed maggots, weevils and wireworms.	eyad 7	Eggplant 		
Ants, cutworms, crickets, earwigs, weevils grasshoppers, grubs, seed maggots, weevils and wireworms	sysū 7	Незд Гейпсе		

steaq tisw ot synd		səldstəpəV		
When using other spreaders, calibrate the spreader settings based on the manufacturer's recommendations.				
some overlapping occurs.				
ss about five feet from the previous one, so	шяке еяср ра		Rotary	
Shut off spreader anytime you stop. For complete coverage		2	Scotts SpeedyGreen®	
when you have reached the header strips or stopped.				
wheel marks to ensure uniform feeding. Shut off spreader			Drop Spreader	
Operate spreader the longest way of the garden area. Overlap		3-1/4	Scotts AccuGreen®	
SHOWS THE	-1	product for 500 sq. f	ianpa ido	

### HOME VEGETABLE GARDEN SPREADER TIPS

Vegetable Garden Insect Control - How to Apply A single application may be applied just prior to planting or after plants emerge and worked into the top 4 to 6 inches of soil. Use 5 cups (1.2 lbs.) of product for a 500 sq. ft. area.					
Shut off spreader anytime you stop. For complete coverage make each pass about five feet from the previous one, so some overlapping occurs. Avoid applying this product on flower, vegetable and shrub plantings.	Yotato?				
Mow lawn first if grass is tall enough to touch the bottom of the spreader. Operate spreader the longest way of the lawn. Overlap wheel to ensure uniform feeding. Shut off spreader when you have reached the header strips or stopped.	Orop Spreader				
	190 Joborg To sbnuog 11 .ps 000, t\000 8.p\p. S p. S\2.t S. t\2.0				
Directions	Spreader Settings				
SPREADER TIPS					

SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT. BUYER OR USER BY WAY OF INDEMINIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WASHINGS OR INSTRUCTIONS FOR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR FIRST OF THE GOOODS SOLD BY MANUFACTURER OR SELLER TO CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MENUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DEMAGES, OR FOR

ISE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUBRANTEES, OR REPRESENATIONS OF ANY KIND TO BUYER, OR USER, EITHER The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED KAIS" (2 ft. x 250 ft.) If necessary, repeat applications every 2 weeks only if there are signs of renewed insect activity.

PERPESEUTATIONS AND AGREEMENTS BETWEEN MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY

How to Apply: Evenly apply a 2-foot band of granules next to the house foundation. Use 1.2 lb. for each 500 sq. ft. **Pests Controlled:** Ants, chiggers, crickets, earwigs, grasshoppers, spiders and ticks. Home Invading Insects: Barrier Treatment Around House Foundation (Outdoors) there are signs of renewed insect activity. Repeat application should be limited to no more than once per seven

How to Apply: Evenly apply 1.2 lb. of product for every 500 sq. ft. area. Repeat application as necessary only if Pests Controlled: Ants, crickets, earwigs, grasshoppers, spiders and ticks. Ornamental Flower Beds (Outdoors) \*\*Food utensils such as tablespoons/teaspoons must not be used for food purposes after use in measuring pesticides. 3) freat new mounds as they appear and 4) thoroughly water the treated area immediately after application. best results, 1) apply in cool weather, 65 to 80° F, or in early morning or late evening hours. 2) Avoid disturbing ants,

To control Fire Ants, uniformly apply 1 tablespoon or 3 teaspoons\*\* of this product to the surface of each mound. **For Fire Ant Mounds** treated area immediately after application. Uniformly apply 2.4 to 4.8 pounds over a 1,000 sq. ft. area. For best results: treat in late spring, and water the

Mole Crickets (including Tawny Mole Crickets) watered immediately after application.

Uniformly apply 4.8 pounds over a 1,000 sq. ft. area. Best results are obtained if the treated area is thoroughly Centipedes, Millipedes, Pavement Ants watered immediately after application.

Uniformly apply 2.4 pounds over a 1,000 sq. ft. area. Best results are obtained if the treated area is thoroughly (including Brown Recluse Spiders) and Ticks (including Brown Dog Ticks) Argentine Ants, Black Turf Ants, Black Turfgrass Ataenius, Fleas (including Cat Fleas), Leafhoppers, Spiders

application should be limited to no more than once per seven days. retreat as necessary. Repeat application is necessary only if there are signs of renewed insect activity. Repeat materials. This 10 pound bag will treat a 4,150 to 8,300 square feet area. Treat when insects first appear and area by using any hand- or power-operated gravity or rotary type spreader designed for the application of granular Moth), and Weevils (including Hyperodes and Bluegrass). Uniformly apply 1.2 to 2.4 pounds over a 1,000 square feet Bugs, Crickets, Cutworms (including Black Cutworms), Earwigs, Grasshoppers, Mealybugs, Sod Webworms (Lawn Foraging Fire Ants, Armyworms (including Fall Armyworms), Billbug (including Bluegrass Billbug), Chiggers, Chinch Use this product for the control of ants, including Carpenter, Red Harvester, Odorous Pyramid, Pharoah (Sugar),

Surface-Feeding Insects Do not allow children or pets to enter the area until dry. Water thorougly after application.

may increase absorption and so should be avoided.

OR CLOTHING:

IE ON SKIN

LAWN, HOME PERIMETER, ORNAMENTAL & FLOWER GARDENS Do not allow children or pets to enter the area until dry.

but not sooner than two weeks after the first application. In New York State, do make a single repeat application of this product if there are signs of renewed insect activity,

(lake, pond, river, stream, wetland, or drainage ditch). In New York State, this product may NOT be applied to any grass or turt area (lawns) within 100 feet of a water body Please read the following "Directions for Use" carefully and fully before applying the product.

watering by 0.1 to 0.5 inches immediately after application generally improves the effectiveness of this product. lawns, home perimeter, ornamental & flower gardens. See specific pests for recommended level. Irrigating or This product is designed for use as a ready-to-use granular insecticide to control various insects and spiders on

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. DIRECTIONS FOR USE

contaminate water when disposing of equipment washwaters or rinsate.

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water. Do not **EUVIRONMENTAL HAZARDS** 

This product is a pyrethroid. If large amounts have been ingested, milk, cream or other digestivle fats and oils **NAISIRAM** OT **3TON** 

Emergencies Only, Call (877) 325-1840. For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Medical HOT LINE NUMBER

 Call a poison control center or doctor for treatment advice. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. IE IN EXES: Call a poison control center or doctor for treatment advice.

> Rinse skin immediately with plenty of water for 15 to 20 minutes. Take off contaminated clothing. **GIA T2811**

chewing gum or using tobacco. Remove and wash contaminated clothing before reuse. clothing. Wash thoroughly with soap and water after handling and before eating, drinking, Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or CAUTION: HAZARDS TO HUMANS AND DOMESTIC ANIMALS PRECAUTIONARY STATEMENTS

Nockon The Best Willer Granies