(O)ARDEN 7 **191**

Distributed by: TechPac, LLC. P.O. Box 24830 Lexington, KY 40524







NET WT 5 LBS. (2.27 kg)

TO CLOSE —— FOLD TOP OF BAG, SECURE WITH A CLIP

Ready-To-Use Sevin® 5% Dust

For Homeowner (Residential) Use Only DIRECTIONS FOR USE I law to use this product in a manner inconsistent with its labeling. General Directions

Crop	Preharvest Interval (Days)	Crop	Preharvest Interval (Days)
Sweet corn	2	Dried beans	21
Garden beets (roots), radishes, rutabagas, turnips (roots), carrots, potato, blackberries, boy- senberries, dewberries, loganberries, raspberries, blüeberries, grapes and strawberries	7	Fresh beans, broccoli, Brussels sprouts, cabbage, cauliflower, kohlrabi, cantaloupe, cucumbers, melons, pumpkins, squash, eggplant, pepper and tomato	3
Collards, garden beets (tops), kale, lettuce (head and leaf), mustard greens, spinach and turnin (tops)	14		

VEGETABLES

Plant	Insect Pests	Rate (Pounds) Per 1000 Sq. Ft.	Comments
Beans	Bean leaf beetle, cucumber beetles, corn earworm, flea beetles, Japanese Beetles, leafhoppers, Mexican bean beetle	0.5	Where Mexican bean beetle is the only problem a rate of 0.25 pound per 1000 sq. ft. is sufficient.
	Lygus bug, Lima bean pod borer, stink bugs	1.0	
Broccoli, Brussels sprouts, cabbage, cauliflower, kohlrabi	Flea beetles, harlequin bugs	0.25-0.5	
	Armyworms, corn earworm, diamond- back moth larvae, imported cabbageworm	0.5-1.0	Will not control cabbage loopers.
Collards, garden beets, kale, lettuce, mustard greens, radishes, rutabagas, spinach, turnips	Flea beetles, harlequin bug, leafhoppers	0.25-0.5	
	Armyworms, corn earworm, diamond- back moth larvae, imported cabbageworm, stink bugs, tarnished plant bug	0.5-1.0	For best control of corr earworm after lettuce heads begin to form, use 1.0 lb. per 1000 sq. ft. Will not control cabbage loopers.
Cantaloupe, cucumbers, melons, pumpkins, squash	Cucumber beetles, flea beetles, melonworms, pickelworm, squash bug	0.5	
Com	Corn earworm, European corn borer, fall armyworm, Japanese beetles, sap beetles	0.75-1.0	For budworm damage in the whorl, apply to entire plant. For insects attacking silks and ears make first application when first silks appear and continue until silks begin to dry. Three or more applications may be required depending upon seventy of infestation.
Carrots	Flea beetles, leaf- hoppers, lygus bugs, spittlebugs, stink bugs	0.5-1.0	
Eggplant, pepper, tomato	Fall armyworm, flea beetles, leafhoppers, tomato fruitworm	0.5	
	European corn borer, stink bugs, tomato hornworm	0.5-1.0	
Potato	Colorado potato beetle	0.25	
	Flea Beetles, leafhoppers	0.5	
	Lygus bugs	1.0	

Plant	Insect Pests	Rate (Pounds) Per 1000 Sq. Ft.	Comments
Blackberries, boysenberries, dewberries, loganberries, raspberries	Japanese beetles, leafhoppers, leafrollers, raspberry aphid	1.0	
Blueberries	Blueberry maggot, cherry fruitworms, cranberry fruitworms, Japanese beetles	0.75-1.0	Apply 3 weeks before harvest and repeat 10 days later if necessary.
Grapes	Grape leaffolder, grape leafhopper, Grape leaf skeletonizer	0.5-1.0	Apply just before first brood leaffolder larvae emerge from rolls and as needed for leafhoppers.
	Grape berry moths, Japanese beetle, June beetle, red-banded leafroller	1.0	
Strawberries	Meadow spittlebug, strawberry leafroller, strawberry weevil	0.5-1.0	

Storage: Store pesticide in original container and place in a locked storage area.

Disposal: If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed or inhaled. Avoid contact with

٦	This produ	First Aid This product is a N-methyl carbamate and is a cholinesterase inhibitor.					
1 25 ft.	If Swallowed	Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swaflow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.					
	If Inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a polson control center or doctor for further treatment advice.					
-	If in Eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then confinue rinsing eye. Call a poison control center or doctor for treatment advice.					
+	If on Skin or Clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.					