



3^{IN}1

WEED & FEED

for SOUTHERN LAWNS


MATA LAS MALEZAS*; PREVIENE LAS MALEZAS* POR 6 MESES; Y FERTILIZA

1 KILLS
WEEDS* INCLUDING DOLLARWEED, DANDELION & CLOVER

2 FEEDS
GREENS & STRENGTHENS YOUR LAWN

3 PREVENTS
BROADLEAF & GRASSY
WEEDS* PLUS CRABGRASS UP TO **6** MONTHS

TREATS UP TO
5,000 Sq Ft
NET WT 12.5 Lbs (5.6 kg)



Active Ingredients:
Dicamba acid 0.088%
Penoxsulam 0.041%
Indaziflam 0.027%
Other Ingredients 99.844%
Total 100.000%

*LISTED WEEDS

KEEP OUT OF REACH OF CHILDREN
CAUTION See back panel for additional
precautionary statements

35-0-3

FOR USE ON
SOUTHERN TURFS
BERMUDAGRASS, ST AUGUSTINEGRASS,
CENTIPEDEGRASS & ZOYSIA

BIOADVANCED
SCIENCE-BASED SOLUTIONS

quick FACTS

- KILLS BROADLEAF WEEDS* INCLUDING DOLLARWEED, DANDELION, CLOVER, CHICKWEED & HENBIT**
- FEEDS & STRENGTHENS THE LAWN. CONTAINS SLOW RELEASE FERTILIZER FOR LONG LASTING FEEDING.**
- PREVENTS GRASSY AND BROADLEAF WEEDS* SUCH AS CRABGRASS, ANNUAL BLUEGRASS, COMMON DANDELION, GOOSEGRASS & SPURGE UP TO 6 MONTHS**

WHEN TO APPLY

Apply anytime in early spring or fall.

WHERE TO USE

Use only on established lawns of the turf types listed below:

Bermudagrass, Centipedegrass, all varieties of St. Augustinegrass (including Floratam) and Zoysia

DO NOT USE ON ANY OTHER TURF TYPES

including: Bahiagrass, Bentgrass, Ormond variety of Bermudagrass, Buffalograss, Carpetgrass, Dichondra, Fescue, Kikuyugrass, Ryegrass, or Seashore Paspalum lawns.

COVERAGE

This bag treats 5,000 sq. ft.



Questions or comments, call toll-free (877) 229-3724
www.BioAdvanced.com



For the full Spanish translation go to www.BioAdvanced.com

****Money Back Guarantee:** If you are not satisfied with this product, we will gladly refund your original purchase price.

***LISTED WEEDS**

KILLS

Beggarweed, Creeping Beggarweed, Dwarf Belfony, Florida Burdock
Buttercup, Hairy Buttonweed, Virginia[†] Clover, Hop Clover, Red Cocklebur, Common Coffeeweed,
(Common Chickory) Daisy, English (Lawn Daisy) Dandelion, False

Dollarweed (Pennywort) Goldenrod Hawkweed Heallall Ivy, Ground (Creeping Charlie) Knotweed Kyllinga, Green[†] Lespedeza, Common Mallow, Roundleaf Matchweed Nutsedge, Purple[†] Nutsedge, Yellow[†] Parsley-piert

Peppergrass Pepperweed, Virginia Plantain, Broadleaf Smartweed Sorrel, Sheep Speedwell, Common Spotted Catsear Spurweed Thistle Vetch, Common Wild Violet Woodsorrel, Creeping
[†]Suppression

PREVENTS

Ageratum, Tropic Barley, Mouse Barnyardgrass Bitweed, Smooth Bitterress Black Nightshade, Hairy Bluegrass, Annual (Poa annua) Bromo, Foxtail Buckwheat, Wild Burweed, Lawn California Burclover Chamomile, False Cheatgrass Crabgrass, Henry

Crabgrass, Large (Hairy Crabgrass) Crabgrass, Smooth Cudweed, Linear Leaf (Purple Cudweed) Dandelion, Cat's Ear Doveweed Evening Primrose, Common Evening Primrose, Cutleaf Flatsedge, Rice Fleabane, Hairy

Foxtail, Giant Foxtail, Green Foxtail, Yellow Filaree, Redstem Goosegrass Groundsel, Common Guineagrass Kochia Kyllinga, Fragrant Lovegrass, Tufted Mallow, Little Maretail Mulberry Weed

Mustard, Black Mustard, Wild Nightshade, American Black Nightshade, Hairy Parthenium Phyllanthus, Longstalked Pigweed, Prostrate Pigweed, Redroot Plantain, Paleseed Puncturevine Redmaids Rocket, London Ryegrass, Italian

Ryegrass, Perennial Sandbur Sedge, Annual Sesbania, Hemp Sida, Prickly (Teaweed) Southern Brassbuttons Sowthistle, Annual Speedwell, Corn Spurge, Prostrate Sunflower, Common Swinecress Tassel Flower

Vetivert Vetch, Purple Wild Carrot Willowherb Woodsorrel (Oxalis)

KILLS AND PREVENTS

Burnweed, American Chickweed, Common Chickweed, Mouseear Clover, White Dandelion, Common Dock, Curly Eclipta Henbit Knotweed, Prostrate Kyllinga, Annual[†] Lambsquarters Medic, Black Plantain, Buckhorn Purslane, Common Pusley, Florida Ragweed, Common Shepardspurse Spurge, Spotted Thistle, Canada Woodsorrel, Yellow
[†]Suppression

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEFORE YOU USE

• Use this product only on established Southern lawns listed on this label – new lawns must be at least 3 months old if sodded, 16 months old if sprigged & 16 months old if seeded. • Do not apply this product to water-saturated soils. • Apply only to actively growing turf. • Apply in spring after the lawn is actively growing and has been mowed 2 times. Do not apply while turfgrass is just turning green in early spring. • Apply in fall no closer than 6 weeks before the first historical frost date for your area. • Do not apply to Centipedegrass after October 1. • Do not apply this product if the lawn appears unhealthy e.g., due to lack of water (drought, watering restrictions), turf disease (dollarspot, large patch, grey leaf spot, take-all root rot, etc.), parasitic microscopic worms (nematodes), turf insects (chinch bugs, etc.), heavy shade, recent aeration or verticutting, or if the lawn is prone to winter injury. • Do not allow granules to come into contact with flowers, ornamental shrubs or trees, or other desirable broadleaf plants, as serious plant injury may occur. • Apply when the air is calm to avoid drift onto desirable plants.

HOW TO USE

• Use rotary or drop type spreader to apply granules evenly over the lawn. To be sure you cover lawn areas uniformly, first treat the border of the lawn then fill in the center. Make parallel passes, walking at a steady pace to ensure even distribution of the granules. • Sweep, brush, or blow any accidentally applied material off driveways, sidewalks etc. back onto the lawn. • Wait 24 hours after applying then activate residual barrier by lightly watering in. • Do not overlap treated areas or make repeat applications within 90 days as over application could damage the lawn. • Do not make more than two applications per site in any 12 month period. • Heavy raking will disturb the weed preventative barrier and reduce the effectiveness of this product.

FOR BEST RESULTS

WHEN TO APPLY • Apply when - weeds are small and actively growing - temperatures are between 50–90°F. • May be applied to dry or (for best results) to dew-wetted lawn. • To prevent Crabgrass and Goosegrass apply early spring before the temperatures are regularly in the lower 80°F range. • To prevent Annual Bluegrass (Poa annua) apply August - December. • Lawns may be seeded 12 months, sodded 6 months, and sprigged 12 months after applying this product.

MOWING

• If the lawn needs mowing, do it 1 to 2 days before (for best results) or 1 day after applying this product. • Clippings from the first three mowings should be left on the treated area. Do not use the clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

WATERING

• If lawn is dry, water 1 to 2 days before using this product. • Do not water for at least 24 hours after application or apply if rain is expected within 24 hours.

HOW IT WORKS


This product contains a proven blend of weed killers to kill existing weeds* plus a residual herbicide to keep them from coming back for up to 6 months. This product starts working immediately to stop weed growth until weeds* are dead within weeks. Plus, it feeds your lawn for quick greening and long lasting feeding.



Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

SPREADER SETTINGS

If your spreader is not listed, please call our Consumer Service Hotline toll-free (877) 229-3724 for help.



- Settings are approximate. The condition of your spreader, the speed at which you walk and uneven terrain may affect the rate of application. Calibrate your spreader according to manufacturer's directions.
- Tip: 1,000 sq ft = 20 ft x 50 ft
- Do not use a spreader that applies granules in narrow, concentrated or uneven bands. Be sure the spreader works properly and spreads granules evenly.

Broadcast Spreader

Drop Spreader

SETTING TO DELIVER 2.5lb/1000 sq. ft.

Broadcast Spreader	SETTING
Earthway Ev-N-Spread	15
Sta-Green	5
Scotts 2000	4 1/2
Scotts Mini	4 3/4
Scotts Deluxe Edge Guard	4 3/4
Scotts Edge Guard Pro	4 1/4

Drop Spreader

Scotts Accugreen 1000	5 3/4
Scotts Accugreen 2000	6 1/2
Scotts Accugreen 3000	4 1/4

ENVIRONMENTAL HAZARDS

- This product is toxic to fish, aquatic invertebrates, and plants.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.
- **Ground Water Advisory:** This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.
- **Surface Water Advisory:** This pesticide may impact water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application.

RIESGOS AMBIENTALES • Este producto es tóxico para los peces, organismos invertebrados acuáticos y las plantas. • Para proteger al medio ambiente, no permitir que el pesticida ingrese o corra por los drenajes de tormenta, drenaje de aguas pluviales, canaletas o aguas superficiales. Aplicar este producto en clima calmo cuando no haya pronóstico de lluvia en las próximas 24 horas para asegurar que el viento o la lluvia no soplen o laven el pesticida fuera del área de tratamiento. Barrer todo producto que quede en caminos, aceras o calle, de nuevo hacia el área tratada del césped o jardín ayudará a evitar que el producto ingrese en cuerpos de agua o sistemas de drenaje. • **Asesoramiento sobre aguas subterráneas:** Este pesticida tiene propiedades y características asociadas con químicos detectados en aguas subterráneas. Este químico puede escurrirse a las aguas subterráneas si se usa en suelos donde sea permeable especialmente donde la mesa de agua sea poco profunda. • **Asesoramiento sobre aguas superficiales:** Este pesticida puede impactar sobre la calidad de agua si drena al agua de lluvia. Este es real para suelos con mal drenaje y suelos con aguas subterráneas poco profundas. Este producto está clasificado como que tiene un alto potencial de alcance al agua de la superficie a través del drenaje durante varios meses o más después de su aplicación.

FIRST AID



NOTE: When calling poison control center, have this product label accessible. If seeking treatment at an emergency room or doctor's office, bring this product label to show medical personnel. You may call toll-free 1-877-229-3763 for medical emergency information. Active Ingredients: 0.088% Dicamba, (CAS # 1918-00-9), 0.041% Penoxsulam (CAS # 219714-96-2) and 0.027% Indaziflam (CAS # 950782-86-2).

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 – 20 minutes.
- Call a poison control center or doctor for further treatment advice.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15 – 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for further treatment advice.

STORAGE & DISPOSAL



PESTICIDE STORAGE Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.
PESTICIDE DISPOSAL & CONTAINER HANDLING If Empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.
If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

Do not contaminate water, food, or feed by storage and disposal


PRIMEROS AUXILIOS

EN CASO DE INGESTIÓN • Llamar inmediatamente a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento. • Hacer que la persona beba tragos de agua, si puede tragar. • No inducir el vómito salvo que así lo indique el médico o el centro de control de envenenamiento. • No administrar nada por vía oral a una persona inconsciente. **EN CASO DE CONTACTO CON LA PIEL O LA ROPA** • Quitar la ropa contaminada. • Enjuagar la piel inmediatamente con abundante agua durante 15 a 20 minutos. • Llamar a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento. **EN CASO DE CONTACTO CON LOS OJOS** • Mantener abiertos los ojos y enjuagar de forma lenta y suave con agua durante entre 15 y 20 minutos. • Retirar los lentes de contacto, si los hubiera, después de los primeros 5 minutos y luego continuar enjuagando los ojos. • Llamar a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento.

NOTA: Tener la etiqueta del producto a mano al llamar a un centro de control de envenenamiento. Si busca tratamiento en una sala de emergencias o en el consultorio de un médico, lleve esta etiqueta del producto para mostrársela al personal médico. También puede llamar al número gratuito 1-877-229-3763 para solicitar información médica de emergencia. Ingredientes activos: 0.088% de dicamba, (CAS n.º 1918-00-9), 0.041% de penoxsulam (CAS n.º 219714-96-2) y 0.027% de indaziflam (CAS n.º 950782-86-2).

ALMACENAMIENTO Y ELIMINACIÓN

No contaminar el agua o los alimentos al almacenar o eliminar
ALMACENAMIENTO DE PESTICIDAS Almacenar en el contenedor original en un lugar fresco y seco, fuera del alcance de los niños, preferentemente en un gabinete bajo llave. Proteger del congelamiento.
ELIMINACIÓN DEL PESTICIDAS Y MANIPULACIÓN DE ENVASES Si está vacío: Envase no rellenable. No rellenar ni volver a utilizar este envase. Colocar el envase vacío en la basura o entregarlo para reciclar si fuera posible. Si está parcialmente lleno: Llamar a la agencia local de residuos sólidos o al número gratuito 1-877-229-3724 para obtener indicaciones para la eliminación. Nunca colocar el producto sin utilizar debajo de algún desagüe interno o externo.



EPA Reg. No. 92564-61

EPA Est. No. indicated by 3rd and 4th digits of the batch number on this package.
(26) = 82757-FL-1 (57) = 071106-GA-003

BioAdvanced™ Science-Based Solutions
3-in-1 Weed & Feed for Southern Lawns
35-0-3

GUARANTEED ANALYSIS

Total Nitrogen (N) 35.0%
35% Urea Nitrogen[†]
Soluble Potash (K₂O) 3.0%
Sulfur (S) Total 4.6%
3.5% Free Sulfur (S)
1.1% Combined Sulfur (S) **F2414**

Derived From: Urea, Polymer Coated Sulfur Coated Urea, & Potassium Sulfate
[†]Contains 17.5% slowly available Nitrogen from Polymer Coated Sulfur Coated Urea
Information regarding the contents and levels of metals in this product are available on the Internet at <http://www.aapfo.org/metals.html>
Check with your county or city government to determine if there are local regulations for fertilizer use.

Distributed by:
SBM Life Science Corp.
1001 Winstead Drive, Suite 500
Cary, NC 27513

BioAdvanced is a trademark of
SBM Life Science Corp.

40224346 17652171