

A Liquid Topical Flea Product for Cats

READ THE ENTIRE LABEL BEFORE EACH USE ACTIVE INGREDIENTS:

lmidacloprid:	9.10%
Pyriproxyfen:	0.46%
OTHER INGREDIENTS:	<u>90.44%</u>
TOTAL:	100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS. CAUTION. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and warm water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Keep out of reach of children. Do not contaminate feed or food. HAZARDS TO DOMESTIC ANIMALS. For external use only. Do not apply to cats or kittens under 8 weeks of age or weighing less than 2 lbs. As with any product, consult your veterinarian before using this product on debilitated, aged, pregnant or nursing cats. Individual sensitivities, while rare, may occur after using ANY pesticide products for cats. If signs persist, or become more severe, consult a veterinarian immediately. If your cat is on medication, consult your veterinarian before using this or any other product. **RESTRICTIONS:** Do not apply to cats or kittens less than 8 weeks old and weighing less than 2 lbs. Do not use on other animals. Do not apply more than one tube per treatment, even for larger cats. Do not have contact or allow children to have contact with treated area until completely dry. Do not treat more than 1 cat per tube. Do not exceed labeled dosage amount for cats.

FIRST AID		
If Swallowed:	Call a poison control center or a doctor immediately for treatment advice. Do not give any liquid to the person. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.	
If In Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.		
lf On Skin:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment or advice.	
HOT LINE NUMBER		
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergencies call 1-844-685-9173.		
NOTE TO PHYSICIAN		

Treat the patient symptomatically.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. DO NOT ALLOW CHILDREN TO APPLY PRODUCT. To prevent harm to you and your cat, read the entire label and package insert before each use. Follow all directions and precautionary statements carefully. Use ONLY on Cats 8 Weeks and Older and Weighing Over 2 lbs.

Side Effects: Monitor your cat after application. Side effects, although very rare, may include signs of skin irritations such as redness, scratching, or other signs of discomfort. Gastrointestinal signs such as hypersalivation, vomiting or diarrhea have also been reported. If these or other side effects (such as lethargy) occur, consult your veterinarian or call 1-844-685-9173.

If you have questions or comments about this product, please write: Promika, LLC 501 Cascade Pointe Lane, Suite 102, Cary, NC 27513

©2016 Promika. Important: Always read and follow label instructions. Adventure™ Plus is a trademark of Promika, LLC. Advantage[®] II is a registered trademark of Bayer Healthcare LLC Animal Health Division. Adventure™ Plus For Cats is not manufactured, or distributed by Bayer Healthcare LLC Animal Health Division, seller of Advantage[®] II For Cats.



CONTAINS IMIDACLOPRID AND PYRIPROXYFEN, THE ACTIVE INGREDIENTS USED IN ADVANTAGE® II FOR CATS

SMALL For Use ONLY on Cats 8 Weeks and Older and Weighing 2 to 5 lbs.

4-Pack of 0.0078 fl oz Net Contents: 0.0312 fl oz

MEDIUM For Use ONLY on Cats 8 Weeks and Older and Weighing 5 to 9 lbs.

4-Pack of 0.014 fl oz Net Contents: 0.056 fl oz

LARGE For Use ONLY on Cats 8 Weeks and Older and Weighing Over 9 Ibs.

4-Pack of 0.027 fl oz Net Contents: 0.108 fl oz



DIRECTIONS FOR USE

Separate paper from corner of blister package. Peel back paper and take out the tube. Remove one applicator tube from the package. Hold applicator tube in an upright position facing away from you and your pet's face and eyes. Pull cap off tube. Turn the cap around and place other end of cap back on tube. Twist cap to break seal, then remove cap from tube. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to expel the entire contents directly on the skin. Do not get this product in your cat's eyes, or allow your cat to ingest this product. The product is bitter tasting and salivation may occur for a short time if the cat licks the product immediately after treatment. Treatment at the base of the skull will minimize the opportunity for the cat to lick the product. Do not allow the product to run off. Discard empty tube as described in the Storage and Disposal section. Under normal conditions this product is effective for a month. However, in case of severe flea infestation, retreatment may be necessary earlier than four weeks. On cats, do not retreat more often than once every seven days. After flea control is attained, return to a monthly retreatment schedule.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Pesticides Storage: Store in a cool, dry place inaccessible to children and pets.

Pesticide Disposal and Container Handling: Nonrefillable container. If empty: Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

> In Case of Emergency, Call: 1-844-685-9173