

APPLICATION GUIDELINES



Turf and Lawn Care

For existing homelawns, apply 3 lb/1,000 sq ft. Apply once a year for normal lawns, or in the spring and fall for optimal results. For new lawns, apply at 5 lb / 1,000 sq ft and till or water/irrigate into soil before planting seed or laying sod. For professional turfgrass, apply 2 lbs/1,000 sq ft twice per year in the early spring and fall which should be broadcast prior to a rain or watered-in. This rate can be split and applied up to 4 times per year.

Orchards > Citrus > Nut Trees > Tree Fruits

For small or home orchards apply annually to trees on ground surface under the drip line at the rate of 1 to 5 pounds/tree, depending on the tree size. Double the application for decline or blight stricken trees. For commercial orchards rates are generally 150 lbs/acre broadcast, 100 lbs/acre banded, 50 lbs/acre for row or trickle irrigation application. Soil and tissue testing should be used to help verify proper rates. Double the broadcast rate for new orchard plantings.

Arboriculture

Apply annually to all trees on ground surface under the drip line. Apply a light coating from the trunk out to the drip line. Recommended application rate for trees:

Tree Base Diameter	Rate
< 10"	1 - 5 lbs.
10" - 20"	5 - 15 lbs.
> 20"	15 - 40 lbs.

Compost

Add 50 lbs of AZOMITE® per ton of compost, preferably at the beginning of the process.

