














# LONG-LASTING WEED CONTROL WITH SEMERA™



Semera 51.0% WDG is a fast-acting contact herbicide that controls selected submersed, emergent and floating aquatic weeds. It is most effective when applied to young, actively growing weeds in water with a pH of less than 8.5. Application to public aquatic areas may require special approval and/or permits. Consult with local state agencies.

## **Outstanding Pre-emerge Performance:**

- Fast-acting, residual control
- Broad-spectrum control of tough weeds
- · Flexible application options
- Unique mode of action
- Excellent tank-mix partner
- Exceptional value
- Doesn't volatilize and move off target

## **Key Uses**

- · Container and field-grown conifers and deciduous trees
- · Around established ornamentals in landscapes
- To maintain bare ground in non-crop areas
- Dormant warm-season turfgrass grown on residential sites, golf courses, sod production and similar sites
- · Quiescent or slow moving bodies of water

Refer to label for complete list of approved uses.



# PRODUCT NOTES

**EPA REGISTRATION NUMBER** 59639-120-91234

**ACTIVE INGREDIENT** flumioxazin 51%

**FORMULATION**Water dispersible granules

**HRAC NUMBER** 14

SIGNAL WORD
Caution

PACKAGE SIZE 4 x 5 lb







### **Use Precautions and Restrictions:**

- Do not apply in enclosed greenhouse structures if plants are present.
- · Do not move plants for 24 hours into enclosed greenhouses until the Semera 51.0% WDG Herbicide treated area has been watered.
- Do not apply when weather conditions favor spray drift from treated areas.
- Do not graze treated fields or hay to livestock.
- Do not incorporate into soil after application.
- Do not apply this product through any type of irrigation system.
- Do not apply when plants are under stress from insects, diseases, animals or winter injury, planting shock or any other stresses.
- Only apply to healthy established trees and ornamentals.
- Do not apply more than 12 oz of Semera 51.0% WDG Herbicide per acre per application.
- Do not apply more than 24 oz of Semera 51.0% WDG Herbicide per acre per year.
- Do not apply to intertidal or estuarine areas.
- In areas with dense weed vegetation only treat 1/2 the water body at one time and wait 10-14 days before treating the remaining area. Do not re-treat the same section of water within 28 days of application.
- · Do not use in water utilized for crawfish farming.
- Do not re-treat the same section of water more than 6 times per year.

Refer to product label for application and mixing instructions.

©2016 Atticus, Important; Always read and follow label instructions before buying or using this product. Semera™ is a trademark of Atticus, LLC



#### **KEY WEEDS CONTROLLED**

American burnweed

Annual bluegrass

Annual sowthistle

Barnyardgrass '

Birdseye pearlwort\*

Bristly starbur\*

California lovegrass\*

California burclover Canada thistle\*

Carolina geranium

Carpetweed

Chamberbitter

Chickweed (common, mouseear)

Coffee senna

Common groundsel

Common lambsquarters

Common purslane

Crabgrass\* (large, smooth, Southern

crabgrass) Dandelion\*

Dogfennel

Doveweed

Eclipta

Field pennycress

Florida beggarweed

Florida pusley

Foxtail\* (bristly, giant, green, yellow)

Goosegrass\*

Green kyllinga\* Ground ivy\* Groundsel tree

Hairy bittercress

Hairy galinsoga

Hairy indigo Hemp sesbania

Henbit

Hoary alyssum

Horseweed\*

**Jimsonweed** Kochia

Ladysthumb

Liverwort

Longstalked Phyllanthus

Mallow (common, little, Venice)

Marsh parsley

Marsh yellowgrass

Mayweed\*

Morningglory (entireleaf, ivyleaf, red/scarlet,

smallflower, tall)

Moss

Mulberry weed Mustard (tumble, wild)

Nightshade (black, Eastern black, hairy)

Northern willow herb

Palmer amaranth

Panicum\* (fall, Texas)

Parsley-piert

Pennsylvania smartweed

Pigweed (prostrate, redroot, smooth,

Pineapple-weed\*

Plantain\* (broadleaf, buckhorn)

Prickly sida (teaweed)

Puncturevine

Ragweed (common, giant)

Redmaids

Redstem filaree\*

Redweed

Russian thistle

Shepherd's-purse

Signalgrass' Spiny amaranth

Spurge (petty, prostrate, spotted)
Tassel-flower

Thickhead

Tropic croton

Tropical spiderwort

Waterhemp (common, tall)

Wild poinsettia

Yellow rocket

Yellow woodsorrel\*

\*Pre-emergence control only