

ACTIVE INGREDIENT Azoxystrobin

% ACTIVE INGREDIENT 22.9%

FORMULATIONSuspended Concentrate

CHEMICAL FAMILY
Strobilurin/Qol inhibitors

BRAND ALTERNATIVE Heritage TL®

MODE OF ACTION
Acropetal Penetrant

RESTRICTED USENone

SIGNAL WORD
Caution

PACKAGING INFORMATION 4x1 gal. 12x1 pt.

AZOXY 2SC SELECT

FUNGICIDE

Long Term Protection in a Convenient Liquid Formulation



Azoxy 2SC Select ™ is the equivalent liquid formulation to the popular WDG Azoxystrobin products. The convenient Suspended Concentrate formulation allows for easier tank mixing and application. Total protection is achieved through a combination of foliar and root absorption. Highly systemic and with a higher than average degree of root uptake, Azoxy 2SC Select ™ provides continuous protection to the whole plant for up to 28 days.

- Convenient liquid formulation for easy tank mixing
- · Defends against both soil borne and foliar diseases
- Provides continuous protection for up to 28 days
- Complete root-to-leaf distribution
- · Constant delivery for even distribution to new growth
- Greater degree of root uptake when compared to other strobilurins
- · Prevents spore germination, thereby preventing many fungal infections
- Does not impact honey bees and other beneficials
- Low use rate





FUNGICIDE

Application Information:

Azoxy 2SC Select ™ may be applied as a foliar spray, in alternating spray programs, or in tank mixes with other registered pesticides. Azoxy 2SC Select is compatible with many commonly used fungicides, liquid fertilizers, herbicides, insecticides, and biological control products. If tank mixes are desired, observe all directions, precautions, and limitations on labeling of all products used.



Azoxy 2SC Select is recommended for control of anthracnose, brown patch, cool weather brown patch (yellow patch), Fusarium patch, gray leaf spot, gray snow mold (Typhula blight), leafspot, melting out, necrotic ring spot, pink patch, pink snow mold, Pythium blight, Pythium root rot, red thread, Rhizoctonia large patch, southern blight, spring dead spot, summer patch, take-all patch, and Zoysia patch on golf courses, lawns and landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas and athletic fields.

Azoxy 2SC Select is most effective when applied prior to disease development. Mix Azoxy 2SC Select with the required amount of water and apply as a dilute spray application in 2-4 gallons of water per 1000 square feet (87-174 gallons per acre). Repeat applications at specified intervals for as long as required. For spot treatments, use 0.4 oz. Azoxy 2SC Select per 1 to 2 gallons of water. Do not apply more than 9.6 quarts product/acre/year (7.1 fl. oz. product/1000 square feet/year). Make applications by ground only.

LABELED DISEASES

Anthracnose Brown Patch Cool Weather Brown Patch Fusarium Patch Gray Leaf Spot Gray Snow Mold Leaf Spot Melting Out Necrotic Ring Spot Pink Patch Pink Snow Mold Pythium Blight Pythium Root Rot Red Thread Rhizoctonia Large Patch Spring Dead Spot Southern Blight Summer Patch Take-All Patch Typhula Blight Yellow Patch Zoysia Patch

LABELED USES

Residential Turf Commercial Turf Parks/Rec Athletic Fields Golf Turf

Download a specimen label at

Prime-SourceLLC.com

