



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

ABSOLUTE® MAXX

Use on Wheat – Reduced Application Rate for Control of Rust Pathogens and Early Season Disease Management

EPA Reg. No. 264-849

FIFRA 2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF COLORADO, KANSAS, MONTANA, NEBRASKA, OKLAHOMA, SOUTH DAKOTA, AND TEXAS

This recommendation is valid until December 31, 2019

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

| WHEAT | | | |
|--|----------------------------|--|--|
| Diseases Controlled | Rate fl oz/Acre | Application Timing | Notes |
| Rusts (Puccinia spp.) | 4.0 – 5.0 | Begin applications preventatively when conditions are favorable for disease development. | For longer residual activity against a wider spectrum of diseases, use the 5.0 fl oz/Acre rate of Absolute Maxx. Absolute Maxx may be applied by ground, aerially, or chemigation. |
| Leaf Blight (Septoria tritici) Powdery Mildew (Blumeria graminis f. sp. tritici) Tan Spot (Pyrenophora tritici-repentis) | 2.5 | Apply early in the spring for control of early developing foliar diseases. Treatments may be included in a tank-mix with a herbicide application. | This treatment alone is not intended to provide season-long protection for the crop. To protect the flag leaf an application of Prostaro 421 SC Fungicide (EPA Reg. No. 264-862) at 5 fl oz/Acre may be made after flag-leaf emergence. Absolute Maxx may be applied by ground, aerial or chemigation. |
| Restrictions | | | |
| <ul style="list-style-type: none"> Do not apply more than one application of Absolute Maxx per season. Do not apply within 35 days of harvest. For optimum disease control a non-ionic surfactant (NIS) may be tank-mixed with Absolute Maxx. For resistance management, do not apply more than 2 consecutive applications of Group 11-containing fungicides per acre per year. Do not allow livestock to graze within the treated area or harvest the treated crop for forage within 30 days after application, or for hay and straw within 45 days after application. Restricted-entry interval (REI) = 12 hours | | | |

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.
 For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.
 For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*