



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

ADMIRE® PRO Systemic Protectant
EPA Reg. No. 264-827

Approval: October 21, 2005

**Label Revisions: Use directions for Pome,
Tropical, and Stone Fruit.**

Supplemental Label

ADMIRE® PRO Systemic Protectant

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

POME FRUIT

Crops Of Crop Group 11 Including: Apple, Crabapple, Loquat, Mayhaw, Pear (including Oriental pear), Quince

Pests Controlled	Rate fluid ounces/Acre
Aphids (including woolly apple aphid) Leafhoppers	7.0 – 10.5
Notes and Restrictions	
Pre-Harvest Interval (PHI): 21 days	
Maximum ADMIRE PRO allowed per season: 10.5 fluid ounces/Acre (0.38 lb AI/Acre)	
Do not apply pre-bloom or during bloom or when bees are actively foraging	
Applications	
Apply specified dosage in the following method:	
1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.	

TROPICAL FRUIT

Including: Acerola, Avocado, Black sapote, Canistel, Feijoa, Jaboticaba, Guava, Longan, Lychee, Mamey sapote, Mango, Papaya, Passionfruit, Persimmon, Pulasan, Rambutan, Sapodilla, Spanish lime, Star apple, Starfruit, Wax jambu

Pests Controlled	Rate fluid ounces/Acre
Aphids Avocado lace bug Leafhoppers Whiteflies	10.5 – 14.0
Pests Suppressed	
Scales Thrips (foliage feeding thrips only)	14.0
Notes and Restrictions	
Pre-Harvest Interval (PHI): 6 days	
Maximum ADMIRE PRO allowed per season: 14.0 fluid ounces/Acre (0.5 lb AI/A).	
Do not apply pre-bloom or during bloom or when bees are actively foraging	
Applications	
Apply specified dosage in the following method:	
1. Chemigation through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.	

STONE FRUIT

Crops Of Crop Group 12 Including: Apricot, Cherry (including sweet and tart), Nectarine, Peach, Plum (including Chickasaw, Damson and Japanese), Plumcot, Prune (fresh and dried)

In-field, Soil Application	
Pests Controlled	Rate fluid ounces/Acre
Aphids (including woolly apple aphid) Leafhoppers	7.0 – 10.5
Notes and Restrictions Pre-Harvest Interval (PHI): 21 days Maximum ADMIRE PRO allowed per season: 10.5 fluid ounces/Acre (0.38 lb AI/Acre) Do not apply pre-bloom or during bloom or when bees are actively foraging	
Applications Apply specified dosage in the following method: 1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.	
Pre-plant, Root Dip Application	
Pest Controlled	Rate fluid ounces/10 gallons root-dip solution
Black peach aphid (infesting roots)	0.87 (26 mLs)
Mix ADMIRE PRO at 0.87 fluid ounces (26 mLs) per 10 gallons of water. Thoroughly wet bare-root transplant to slightly above the graft union by soaking roots in the ADMIRE PRO solution for up to 5 minutes. Allow solution to dry on roots and transplant trees as soon as possible following treatment.	

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).