



COMPLETE BRAND INSECT DUST

For GARDENS Ready-To-Use

This product delivers effective long lasting control of Aphids, Armyworms, Beetles, Caterpillars, Japanese Beetles plus other listed pests. Protecting your garden from pests has never been easier.

READ THE LABEL FIRST!

QUICK FACTS

- Kills Over 60 Types of Insects
- Effective Long Lasting Control
- Use on Fruit Trees, Flowers, Fruits and Vegetables

KILLS

Aphids, Armyworms, Beetles, Japanese Beetles, Cabbageworms, Caterpillars, Hornworms, Fruit Flies, and other listed pests.

WHERE TO USE

Use on Home Garden Fruits, Roses, Flowers and Vegetables



For questions or comments, call toll-free 1-877-BayerAG, (1-877-229-3724), or visit us on-line at www.BayerAdvanced.com



Para preguntas, comentarios o solicitar la etiqueta completa en español llame sin cargo al 1-877-BAYERAG (1-877-229-3724) o visite nuestra página en www.BayerAdvanced.com

ACTIVE INGREDIENT: Permethrin* 0.25%
OTHER INGREDIENTS 99.75%
 *Cis/trans ratio: 100.00%
 Min. 35% (±) cis, Max. 65% (±) trans

EPA Reg. No. 28293-126-72155
 EPA Est. No. 58996-MO-1A 432-TX-1B
 264-MO-003C 065387-AR-002D
 67572-GA-1E 011623-GA-003F

Subscript used is first letter of run code on container.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read Entire Label Before Each Use.

HOME GARDENS

For use on home gardens. Not for use on plants being grown for sale or other commercial use. Do not water the treated area to the point of run-off. Do not make applications during rain

HOW TO APPLY

Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. (Many insects hide on the underside of the leaf).

Apply when air is calm to avoid drift and contact with eyes and skin.

WHEN TO APPLY

Use at first signs of insect infestation, preferably before damage from feeding occurs. Note the pre-harvest (time between use and harvest) interval with each crop listed.

INSECTS CONTROLLED

On Fruit and Nut Trees: Apple aphid, caterpillars, codling moth, filbertworm, fruit fly, green fruitworm, leafhoppers, leafrollers, lepidopterous larvae, lesser peach tree borer, mirid bug, navel orangeworm, oriental fruit moth, peach twig borer, pear psylla, plum curculio, redbanded leafroller, rose chafer, spotted tentiform leafminer, tarnished plant bug, thrips, walnut husk fly, and white apple leafhopper.

(cont'd)

On Vegetables & Melons: Aphids, armyworms, artichoke plume moth, asparagus beetle, aster leafhopper, cabbage aphid, cabbage looper, Colorado potato beetle, corn earworm, corn rootworm (adult), cucumber beetle (adult), cutworm complex, cutworms, diamondback moth, European corn borer, fall armyworm, flea beetles, granulate cutworm, hornworms, imported cabbageworm, leafhoppers, leafminers, melonworm, onion maggot (adult), onion thrips, pepper weevil, pickleworm, plant bugs (including lygus, squash, and stink bugs), potato aphid, potato flea beetle, potato leafhopper, potato psyllid, potato tuberworm, rindworm, squash vine borer, tomato fruitworm, tomato pinworm, and vegetable leafminer.

On Ornamentals (Including Roses): Asparagus beetle, beet armyworm, cabbage looper, Japanese beetle, lygus bug, and omnivorous leafroller.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry place. Do not contaminate feed or foodstuffs by storage or disposal.

PESTICIDE DISPOSAL: If Partly Filled: • Call your local solid waste agency or 1-877-229-3724 for disposal instructions. • Never place unused product down any indoor or outdoor drain.

CONTAINER DISPOSAL: If Empty: • Nonrefillable container. Do not reuse or refill this container. • Place in trash or offer for recycling if available.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Humans: Harmful if swallowed, inhaled. Avoid breathing dust. Avoid contact with eyes and skin. Wash thoroughly with soap and water after handling. Avoid contamination of feed and foodstuffs. **Animals:** Harmful if swallowed, inhaled. Consult a veterinarian before using this product on debilitated, aged, pregnant, or nursing animals or animals on medication. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

FIRST AID +

IF IN EYES • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

IF SWALLOWED • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.

IF INHALED • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1 (877) BAYERAG [1 (877) 229-3724].

Crop - Dust Lightly	Applications Per Season (Max)	Pre-harvest interval
Cole Crops – Broccoli, Brussel Sprouts, and Cauliflower.	8	1 day
Cole Crops – Cabbage, and Chinese Cabbage (tight-heading varieties only)	5	1 day
Cucurbit vegetables including cucumbers, gourds, honeydew melons, pumpkins, cantaloupe, zucchini, squash, and watermelon (including hybrids).	8	May be applied until harvest
Eggplant	10	3 days
Bell Peppers – including green peppers	8	3 days
Tomatoes	6	May be applied until harvest
Leafy Vegetables (except brassica) including arugula, celery, lettuce, parsley, radicchio, rhubarb, spinach, and Swiss chard.	5	1 day
Onions (dry bulb and garlic)	6	1 day
Potatoes	8	14 days
Sweet Corn	6, 8 in Florida	1 day
Artichokes	5	May be applied until harvest
Asparagus	4	1 day
Tree Crops - Dust Lightly, using a dust gun or rotary duster	Applications Per Season (Max)	Pre-harvest interval/Remarks
Apples		Do not apply after petal fall
Cherries - In the East	6	No more than 4 applications after petal fall
Cherries - In the West	4	No more than 3 applications after petal fall
Peaches	5	14 days
Pears (dormant)	2	May be applied during the dormant through the delayed dormant growth periods only.
Pears (pre-bloom)	2	May be applied from dormant through bud burst.
Walnuts	8	1 day

Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

Distributed by: Bayer Advanced
 A Business Unit of Bayer CropScience LP
 P.O. Box 12014, 2 T.W. Alexander Drive
 Research Triangle Park, NC 27709

Bayer (reg'd), Bayer Cross (reg'd) and Bayer Advanced™ are trademarks of Bayer.

79540396 R.2



6 87073 00065 8