



BRUSH KILLER

Controls Woody Plants and Broadleaf Weeds
Contains TRICLOPYR

Active Ingredient(s):

Triclopyr (3,5,6-trichloro-2-pyridinyloxyacetic acid), as the triethylamine salt* 8.8%
Inert Ingredients 91.2%

Contains 0.6 pounds of active ingredient per gallon.

*Equal to 6.36% Triclopyr acid.

Keep out of reach of children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Eye Irritation * Harmful If Swallowed, Inhaled Or Absorbed Through The Skin. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Do not allow children or pets onto treated areas until sprays have dried. Remove contaminated clothing and wash before reuse.

First Aid

If in Eyes: Immediately flush eyes with plenty of water for 15 minutes. Call a physician if irritation persists.

If on Skin: Remove contaminated clothing and wash skin with soap and water. If irritation continues, get medical attention.

If Swallowed: Call a physician. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

EPA EST.No.829-FL-1

EPA Reg. No. 62719-226-829

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USING.

Read entire label before use. Use strictly in accordance with label precautionary statements and directions.

General Information

SA-50 brand BRUSH KILLER is recommended for the control of unwanted woody plants, vines and broadleaf weeds around homes, cabins, buildings, trails, fences, walkways, rights-of-way, and other non-crop areas.

General Use Precautions

Do not apply this product directly to, or allow to drift to grapes, vegetable crops, flowers, ornamentals or other desirable foliage. Applications should be made when wind is low and there is little chance of spray drift. Do not spray when the wind is blowing toward desirable plants. Do not apply with nozzles that produce a fine-droplet spray. Measuring utensils and spray equipment should not be reused for purposes other than herbicide application.

Do not use on or around edible food or food crops.

Do not graze treated areas.

Woody Plants and Vines Controlled Include:

Alder, Arkansas Rose, Arrowwood, Ash, Aspen, Beech, Birch, Blackberry, Blackgum, Box Elder, California Rose, Cascara, Ceanothus, Cherry, Chinquapin, Choke Cherry, Cottonwood, Crataegus (Hawthorn), Douglas Fir, Dogwood, Elderberry, Elm, Hazel, Honeysuckle, Hornbeam, Kudzu, Locust, Madrone, Maples, Mesquite, Mimosa, Mulberry, Oaks, Persimmon, Pine, Poison Ivy, Poison Oak, Poplar, Raspberry, Salmonberry, Sassafras, Scotch Broom, Sumac, Sweetbay Magnolia, Sweetgum, Sycamore, Tanoak, Thimbleberry, Trumper Creeper, Tulip Poplar, Virginia Creeper, Western Hemlock, Willow, Winged Elm, Wild Grape, Wild Rose.

Perennial Broadleaf Weeds Controlled Include:

Bindweed, Burdock, Canada Thistle, Chickory, Clover, Curly Dock, Dandelion, Field Bindweed, Field Horsetail Rush, Henbit, Lambsquarter, Oxalis, Plantain, Ragweed, Smartweed, Spurge, Tansy Ragwort, Vetch, Wild Lettuce, Wild Violet.

Application Directions

When to Apply

For best results, apply as a full coverage foliar spray when brush and weeds are fully leafed out and actively growing. Wilting or other symptoms will occur in 1-6 weeks. Apply only when the air is calm to prevent drift to desirable vegetation. Rainfall within 24 hours may reduce effectiveness.

Application Methods

Foliage Spray

Using a hose-end or tank sprayer, mix 8-16 tablespoons (4-8 ounces) in enough water to make one gallon mixed spray. For mature or hard-to-control plants (such as ash, blackgum, choke cherry, elm, maple, oak, pine or winged elm), or during drought conditions, use the highest rate of application indicated. Resprouting may occur the year following treatment when lower rates are used. The addition of a surfactant (such as SA-50 brand Surfactant for Herbicides) enhances herbicidal action and improves distribution of the spray on hard-to-wet plant surfaces. Apply at the rate of one gallon mixed spray to approximately 500 square feet of area. Coverage may vary depending on the amount of foliage to be treated. Using coarse spray pattern and low pressure, apply as thorough foliage spray to wet all leaves. See "Use Precautions."

Stump Treatment

To kill stumps and prevent sprouting, apply the undiluted product with a paint brush or backpack sprayer to completely cover the freshly cut surface. Do not reuse paint brush. Wrap in several layers of newspaper and discard in trash.

Vine Control

When treating vines such as poison oak, poison ivy, etc. that grow on or around desirable plants, do not allow spray to contact desirable vegetation. Cut the vine first and treat

as directed for stump treatment.

STORAGE AND DISPOSAL

Storage: Store product in original container and keep out of reach of children. Store in a cool, dry place. Avoid contamination of food and foodstuffs.

Container Disposal: Do not reuse container. Rinse container thoroughly before discarding in trash. Securely wrap container in several layers of newspaper and discard in trash.

Warranty Disclaimer

Southern Agricultural Insecticides, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below.


SOUTHERN AGRICULTURAL INSECTICIDES, INC. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Southern Agricultural Insecticides, Inc. or the seller. All such risks shall be assumed by Buyer.

NOTICE: Buyer assumes all risk and liability of use, storage and /or handling of this material not in accordance with the terms of this label.

IN CASE OF EMERGENCY CALL: CHEMTREC 1-800-424-9300



Southern Agricultural Insecticides, Inc.

FL 34220

Hendersonville, NC 28793

Boone, NC 28607

revised 1998