



BurnOut II

Fast Acting Weed & Grass Killer

CONCENTRATE

*For Non-Selective Control
of Herbaceous Broadleaf
and Grass Weeds.*

MADE FROM PLANT OILS

DANGER
KEEP OUT OF REACH
OF CHILDREN

DANGER: See back panel for
additional precautionary statements.



ACTIVE INGREDIENTS:

Citric Acid24%
Clove Oil8.0%

OTHER INGREDIENTS:

Water, Lauric Acid, Sodium Caprylate,
Gum Arabic, Xanthan Gum, Sodium Acetate
Total Other 68%
Total 100%

NET CONTENTS: 2.5 US GALLON (9.46 L)

BurnOut II

Fast Acting Weed & Grass Killer CONCENTRATE

DANGER: KEEP OUT OF REACH OF CHILDREN

FIRST AID

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

IF SWALLOWED: Call a doctor or get medical attention. Do not induce vomiting. Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND ANIMALS

DANGER: Causes eye irritation. Wear safety glasses and gloves when applying. Do not get in eyes, on skin, or on clothing. Harmful if swallowed.

In case of contact, immediately flush eyes with plenty of water. Get medical attention if irritation persists. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS: Avoid spraying directly into water.

DO NOT APPLY THIS PRODUCT THROUGH ANY TYPE OF IRRIGATION SYSTEM.

PERSONAL PROTECTION EQUIPMENT (PPE):

Applicators and other handlers must wear appropriate protective eyewear, such as goggles or face shield, long sleeved shirt and long pants, waterproof gloves and shoes.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Keep from freezing. Store only in original tightly sealed container and out of reach of children.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container (bottle). Rinse thoroughly before discarding in trash.

DIRECTIONS FOR USE:

DIRECTIONS FOR USE: Mix 1 part BurnOut II Weed and Grass Killer concentrate to 3 parts water; (mixing ratio is 3:1). For older more established weeds a (2:1) mixing ratio may be required to kill target plants.

BROAD CAST APPLICATIONS: Spray BurnOut to run-off using tank sprayer; typically 35-75 gallons of spray solution per acre.

BAND APPLICATION: Using trigger sprayers, cover desired weeds and grasses as needed, typically 25-50 gallons per acre.

SHAKE WELL BEFORE USING

BurnOut Weed & Grass Killer is non-selective to both broadleaf and grassy weeds and is recommended for areas such as borders, driveways, sidewalks, around mature tree bases, around buildings, in fence rows, in greenhouses, school grounds, and around flower beds.

BurnOut Weed & Grass Killer is fast acting. Foliar contact results in rapid desiccation of plant cell tissue, and top growth reduction of herbaceous perennial weeds and grasses. Re-treatment is required for control of established perennial weeds.

FOR EARLY SEASON ANNUAL WEED CONTROL:

When weeds are small (3 to 5 leaf) and actively growing, apply BurnOut Weed & Grass Killer to the foliage of the weeds, making sure that all of the foliage is wet. Thorough coverage is necessary to achieve desirable control.

BurnOut Weed & Grass Killer is fast acting. It can be applied up to one hour before rainfall and still be effective. If it rains less than one hour after treatment, determine if the treated leaves are showing symptoms. If symptoms are not obvious, wait until the leaves dry and re-treat.

FOR CONTROL OF LARGER ANNUALS AND BURNDOWN OF PERENNIAL WEED GROWTH:

Larger annual weeds and perennials are more difficult to control and may require retreatment.

Treat initially as recommended above and repeat if new growth of leaves appear. Thorough coverage of all foliage is necessary to achieve desirable control. Best results are achieved from spring / early summer applications to actively growing young weeds. Weeds that are mature, dormant, or hardened due to moisture stress are more tolerant of herbicide treatments. Only contacted vegetation will be affected. Rainfall within one hour of application will reduce degree of control.

Avoid application to reactive metals such as aluminum, tin, iron, or items such as fencing or lawn furniture in order to prevent staining, mottling, or otherwise interfering with finished metal surfaces.

NOTE: All contacted vegetation will be affected. Avoid contact with desirable plants. Overspray or drift will injure or kill contacted vegetation.

TYPICAL WEEDS CONTROLLED

ANNUAL

Broadleaf Weeds

Black Medic
Chickweed
Cinquefoil (rough)
Lamb's-quarters
Mustard spp.
Oxalis spp.
Pigweed spp.
Ragweed spp.

Annual Grasses

Crabgrass
Foxtail spp.

PERENNIAL

Broadleaf Weeds

Cinquefoil (silvery)
Dandelion
Plantain spp.
Toadflax
Tufted vetch
Wild carrot

Perennial Grasses

Bluegrass
Quack grass

CONDITIONS OF SALE

The label instructions for the use of this product reflect the opinion of experts based on research and field use. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffective or other unintended consequences may result because of such factors as weather conditions, presence of other materials, herbicide resistant weed preparations or the use or application of the product contrary to label instructions, all of which are beyond the control of St. Gabriel Organics. ALL SUCH RISKS WILL BE ASSUMED BY THE USER. St. Gabriel Organics shall not be responsible for losses or damages resulting from the use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label. St. Gabriel Organics warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above. St. Gabriel Organics DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. ST. GABRIEL ORGANICS EXCLUSIVE REMEDY AND ST. GABRIEL ORGANICS EXCLUSIVE LIABILITY SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THIS PRODUCT.

This Product has not been registered by the United States Environmental Protection Agency.
St. Gabriel Organics Represents That this product qualifies for exemption from registration with the Environmental Protection Agency under the Federal Insecticide, Fungicide, and Rodenticide Act.

St. Gabriel Organics

14044 Litchfield Dr., Orange, VA 22960

1-800-801-0061 • www.StGabrielOrganics.com

