

Centipede Weed And Feed

10926

Covers up
to 5,000 Sq. Ft.



Slow Release Formulation



ferti·lome®



Centipede Weed And Feed

15-0-15

- ★ Slow Release Formulation.
- ★ Contains 3% Iron For Super Greening Power.
- ★ Contains *Trimec® Herbicide.



Controls Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge, And Many Other Broadleaf Weeds Which Are Listed On This Label, In St. Augustinegrass, Centipedegrass, Bahiagrass, Bermudagrass, Zoysiagrass, Bluegrass, Perennial Ryegrass, Fescue And Bentgrass Lawns.

| | |
|--|---------|
| ACTIVE INGREDIENTS: | |
| Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid..... | 0.29% |
| Dimethylamine salt of 2,4-dichlorophenoxyacetic acid..... | 0.17% |
| Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid) | 0.07% |
| INERT INGREDIENTS: | 99.47% |
| TOTAL: | 100.00% |
| <small>THIS PRODUCT CONTAINS: 4.80 lbs. or 0.24% (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton. 2.80 lbs. or 0.14% 2,4-dichlorophenoxyacetic acid equivalent per ton. 1.20 lbs. or 0.06% 3,6-dichloro-o-anisic acid equivalent per ton. Isomer Specific by AOAC Methods Contains the single isomer form of Mecoprop-p. TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.</small> | |

KEEP OUT OF REACH
OF CHILDREN
CAUTION

SEE BACK PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected.
Apply this product only to your lawn / garden, and sweep any product that lands
on the driveway, sidewalk, or street, back onto your lawn / garden.

NET WEIGHT 20 LBS. (9.07 KG)

10926

Centipede Weed And Feed

Centipede Weed And Feed

Centipede Weed And Feed



Centipede Weed And Feed

STOP! READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

FIRST AID

| | |
|---|---|
| If In Eyes: | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a Poison Control Center or doctor for treatment advice. |
| If On Skin Or Clothing: | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 - 20 minutes.• Call a Poison Control Center or doctor for treatment advice. |
| If Swallowed: | <ul style="list-style-type: none">• Call a Poison Control Center or doctor immediately for treatment advice.• Have a person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a Poison Control Center or doctor.• Do not give anything by mouth to an unconscious person. |
| Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment. | |

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

CONTAINER DISPOSAL: **If Empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If Partly Filled:** Call your Local Solid Waste Agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

USE PRECAUTIONS

Do not use this product where desirable Clovers are present. Do not apply in areas underlaid by roots of desirable Trees or Shrubs. Do not exceed specified dosages for any area. Do not apply to newly seeded Lawns. During application, close spreader openings when turning. Do not contaminate domestic or irrigation waters.

INSTRUCTIONS

ferti·lome® CENTIPEDE WEED AND FEED controls a wide range of Lawn Weeds, including Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many others commonly found in Home Lawns. In addition, a 15-0-15 fertilizer provides a supply of Plant food. Possible retreatment may be necessary if (a) Lawn is heavily overrun by Weeds or (b) if adverse low moisture conditions prevail and Weeds are in a state of poor growth. A second application may be made in approximately 30 days. Do not use on Dichondra. Under conditions of adequate moisture, high temperatures and high humidity, over-application may cause minor transitory discoloration on St. Augustinegrass. Do not use this on "Floritam" St. Augustinegrass.

The maximum application rate to turf is 0.3 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

1. Mow Lawn to normal height 1 to 2 days before application.
2. Water Lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when Weeds are young and actively growing, preferably in the morning when dew is on the Grass. If Grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
4. Do not wash from Weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

HOW MUCH TO APPLY

For Kentucky Bluegrass, Fescues, Perennial Ryegrass, Bermudagrass, Zoysiagrass, St. Augustinegrass And Centipedegrass: Apply 3.2 to 4.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of Weeds or for adverse seasonal conditions. The second application of 3.2 to 4.0 pounds of product per 1,000 square feet should be used 30 days after the first application.

The maximum application rate to turf is 0.3 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage and avoid spreading granules onto Flowers, Vegetables, and Ornamental Shrubbery. Do not exceed the application rate for any area and use care in the application of this product within the dipline of Trees, Shrubs and other Ornamentals.

KEEP OUT OF REACH OF CHILDREN

Distributed By:

VOLUNTARY PURCHASING GROUPS, INC.

**P.O. BOX 460 • HWY. 56 WEST
BONHAM, TEXAS 75418**

Visit Us At: www.fertilome.com

EPA Reg. No. 2217-603-7401 • EPA Est. No. 7401-TX-001

Information regarding the contents and levels of metals in this product
is available on the internet at: <http://www.aapfco.org/metals>

BROADLEAF WEEDS

| | | | |
|---------------------------|---------------------|--------------------|----------------------|
| Annual Yellow Sweetclover | Clover, Sweet | Narrowleaf Vetch | Spotted Spurge |
| Aster | Clover, White | Nettle | Spurweed |
| Austrian Fieldcress | Cockle | Orange Hawkweed | Stinging Nettle |
| Bedstraw | Cocklebur | Oriental Cocklebur | Strawberry, |
| Beggarticks | Common Mullein | Oxalis | India Mock |
| Betony, Florida | Creeping Jenny | Parsley-Piert | Tall Nettle |
| Bindweed, Field | Cudweed | Parsnip | Tall Vervain |
| Bird Vetch | Curly Dock | Pearlwort | Tansy Ragwort |
| Bitter Wintercress | Daisy, English | Pennycress | Tansy Mustard |
| Bittercress, Hairy | Daisy Fleabane | Pepperweed | Tanweed |
| Bitterweed | Daisy, Oxeye | Pigweed | Thistle |
| Black-Eyed Susan | Dandelion | Pineywoods | Trailing Crownvetch |
| Black Medic | Dichondra | Bedstraw | Tumble Mustard |
| Black Mustard | Dogfennel | Plains Coreopsis | Tumble Pigweed |
| Blackseed Plantain | Dollarweed | Plantain | Velvetleaf |
| Blessed Thistle | False Dandelion | Poison Ivy | Venice Mallow |
| Bloodflower | Falseflax | Poison Oak | Virginia Buttonweed |
| Milkweed | False Sunflower | Pokeweed | Virginia Creeper |
| Blue Lettuce | Fiddleneck | Poorjoe | Virginia Pepperweed |
| Blue Vervain | Florida Pusley | Prairie Sunflower | Wavyleaf Bulbthistle |
| Bracted Plantain | Frenchweed | Prickly Lettuce | Western Clematis |
| Bristy Oxtongue | Galinsoga | Prickly Sida | Western Salsify |
| Broadleaf Dock | Goldenrod | Prostrate Knotweed | White Mustard |
| Broadleaf Plantain | Ground Ivy | Prostrate Pigweed | Wild Aster |
| Broomweed | Gumweed | Prostrate Spurge | Wild Buckwheat |
| Buckhorn | Hairy Fleabane | Prostrate Vervain | Wild Carrot |
| Buckhorn Plantain | Hawkweed | Puncturevein | Wild Four-O'clock |
| Bulbous Buttercup | Healall | Purslane, | Wild Garlic |
| Bull Thistle | Heartleaf Drymary | Common | Wild Geranium |
| Bullnettle | Heath Aster | Ragweed | Wild Lettuce |
| Burclover | Hedge Bindweed | Red Sorrel | Wild Marigold |
| Burdock | Hedge Mustard | Redroot Pigweed | Wild Mustard |
| Burning Nettle | Hoary Cress | Redstem Filaree | Wild Onion |
| Burweed | Hoary Plantain | Rough Cinquefoil | Wild Parsnip |
| Buttercup | Hoary Vervain | Rough Fleabane | Wild Radish |
| Buttonweed | Horsenettle | Russian Pigweed | Wild Rape |
| Canada Thistle | Jimsonweed | Russian Thistle | Wild Strawberry |
| Carolina Geranium | Knawel | Scarlet Pimpernel | Wild Sweet Potato |
| Carpetweed | Knotweed | Scotch Thistle | Wild Vetch |
| Catchweed | Kochia | Sheep Sorrel | Woodsorrel |
| Bedstraw | Lambsquarters | Sheeperspurse | Woolly Croton |
| Catnip | Lespedeza | Slender Plantain | Woolly |
| Catsear | Mallow | Smallflower | Morningglory |
| Chickweed, | Matchweed | Galinsoga | Woolly Plantain |
| Common | Mexicanweed | Smooth Dock | Wormseed |
| Chickweed, | Milk Vetch | Smooth Pigweed | Yarrow |
| Mouseear | Morningglory | Sorrel | Yellow Rocket |
| Chicory | Mouseear | Sowthistle | Yellowflower |
| Cinquefoil | Hawkweed | Spanishneedles | Pepperweed |
| Clover, Crimson | Mugwort | Speedwell | and other |
| Clover, Hop | Musk Thistle | Spiny Amaranth | Broadleaf Weeds |
| Clover, Red | Mustard | Spiny Cocklebur | |
| Clover, Strawberry | Narrowleaf Plantain | Spiny Sowthistle | |
| | | Spotted Catsear | |

SPREADER GUIDE ONE BAG WILL COVER UP TO 5,000 SQUARE FEET

These settings can be used only as a guide, they are not absolute. The speed at which the applicator walks, spreader condition and the texture of the turf will alter the rate of application. If product remains in spreader after application to recommended square foot coverage, spread remainder over same area going at right angles (See Diagram).



| SPREADER | SPREADER SETTINGS |
|--|-------------------|
| ferti·lome® by Spyker | 3.2 |
| ferti·lome® Plus Iron Broadcast/Earthway 2850 | 14 |
| ferti·lome® Plus Iron Prof. Broadcast/Earthway 2800A | 14 |
| ferti·lome® Deluxe Broadcast/Earthway 2050 S.U. | 14 |
| Scott Broadcast/Speedy Green | 6 |
| Republic/Scotts Broadcast | 5 |
| Republic Drop | 6 |
| Precision Broadcast/Red Devil | 10 |
| Precision Drop FH-22D | 9 |

The above spreader settings are approximate. Calibrate your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

15-0-15

GUARANTEED ANALYSIS

| | |
|--|-------|
| Total Nitrogen (N) | 15% |
| 4% Water Insoluble Nitrogen* | |
| 11% Urea Nitrogen | |
| Soluble Potash (K ₂ O) | 15% |
| Boron (B) | 0.02% |
| Copper (Cu) | 0.05% |
| 0.05% Water Soluble Copper (Cu) | |
| Iron (Fe) | 3.00% |
| 0.10% Water Soluble Iron (Fe) | |
| Manganese (Mn) | 0.05% |
| 0.05% Water Soluble Manganese (Mn) | |
| Zinc (Zn) | 0.05% |
| 0.05% Water Soluble Zinc (Zn) | |
| Derived from: Urea, Urea Formaldehyde, Muriate of Potash, Sodium Borate, Copper Sulfate, Iron Sulfate, Iron Oxide, Manganese Sulfate, Zinc Sulfate. | |
| *This product contains 4% Slow Release Nitrogen from Urea Formaldehyde | |
| F370 | |

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the Manufacturer or the Seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied in any written or verbal statements or agreements. No employee or agent of the Seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.



6-08