

Home & Garden

CHELATED LIQUID IRON

FOR THE CORRECTION AND PREVENTION OF FOLIAGE YELLOWING DUE TO IRON DEFICIENCY OF FLOWERS, SHRUBS, TREES, VEGETABLES AND LAWNS.

MINIMUM GUARANTEED ANALYSIS:

Soluble Iron as (Fe)	5.00%
Chelated Iron as (Fe)	3.50%
Combined Sulfur as (S).....	3.40%

DERIVED FROM: Iron Sulfate, Iron Lignin Sulfonate.

Chelating agent is Lignin Sulfonate

Guaranteed by: Southern Agricultural Insecticides, Inc.

P.O. Box 218 Palmetto, FL 34220

F 672-841SA-50

Brand Chelated Liquid Iron is intended for use in the correction and prevention of foliage yellowing due to iron deficiency of flowers, shrubs, trees, vegetables and lawns.

USE PRECAUTIONS

THIS PRODUCT WILL SEVERELY STAIN CONCRETE, MASONRY AND PAINTED SURFACES. Be extremely careful not to allow spray or runoff to contact with these materials. Do not mix with other fertilizers, fungicides or insecticides unless compatibility is known.

CAUTION:

KEEP OUT OF REACH OF CHILDREN.

Avoid contact with skin and eyes. Avoid breathing spray mist. Do not take internally.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 2 glasses of water or milk and seek medical attention.

IF IN EYES: Flush eyes thoroughly with a large volume of water. If irritation persists seek medical attention.

IF ON SKIN: Rinse affected area with clear water and then wash thoroughly with soap and water.

DIRECTIONS FOR USE

Shake well before using.

FLOWERS, VEGETABLES, SHRUBS AND TREES

FOLIAGE APPLICATION:

Mix the material at the rate of 1 tablespoonful per gallon of water. Apply in a manner that will give thorough coverage to upper and lower surfaces of foliage. Use an amount sufficient to cause run-off of spray.

SOIL APPLICATION:

Use at the rate of 1 fluid ounce in at least 1 gallon of water per 100 sq. ft. For flower beds, treat the bed area. For shrubs and trees, treat the root area. This area usually extends out as far as the foliage.

LAWNS and TURF

Use 1 pint (16 fl. oz.) per 1,000 sq. ft. Apply in a sufficient volume of water to thoroughly wet the foliage and runoff in the root zone. Hose attached sprayers provide a convenient method to treat lawns.

Note: Iron deficiency in many plants is an indication of high pH or alkaline soil. If yellowing persists or returns rapidly after treatment, soil pH should be checked and adjusted if necessary.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions and precautions given herewith.

Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220

Hendersonville, NC 28793

Boone, NC 28607

revised 1998

DIRECTIONS FOR USE

Shake well before using.

FLOWERS, VEGETABLES, SHRUBS AND TREES

FOLIAGE APPLICATION: Mix the material at the rate of 1 tablespoonful per gallon of water. Apply in a manner that will give thorough coverage to upper and lower surfaces of foliage. Use an amount sufficient to cause run-off of spray.

SOIL APPLICATION: Use at the rate of 1 fluid ounce in at least 1 gallon of water per 100 sq. ft. For flower beds, treat the bed area. For shrubs and trees, treat the root area. This area usually extends out as far as the foliage.

LAWNS and TURF

Use 1 pint (16 fl. oz.) per 1,000 sq. ft. Apply in a sufficient volume of water to thoroughly wet the foliage and runoff in the root zone. Hose attached sprayers provide a convenient method to treat lawns.

Note: Iron deficiency in many plants is an indication of high pH or alkaline soil. If yellowing persists or returns rapidly after treatment, soil pH should be checked and adjusted if necessary.

Home & Garden CHELATED LIQUID IRON

FOR THE CORRECTION AND PREVENTION
OF FOLIAGE YELLOWING DUE TO IRON
DEFICIENCY OF FLOWERS, SHRUBS,
TREES, VEGETABLES AND LAWNS.

MINIMUM GUARANTEED ANALYSIS:

Soluble Iron as (Fe)	5.00%
Chelated Iron as (Fe)	3.50%
Combined Sulfur as (S).....	3.40%

DERIVED FROM: Iron Sulfate, Iron Lignin Sulfonate.

Chelating agent is Lignin Sulfonate

Guaranteed by: Southern Agricultural Insecticides, Inc.
P.O. Box 218 Palmetto, FL 34220

Southern Agricultural Insecticides, Inc.
Palmetto, FL Hendersonville, NC Boone, NC
F 672-841 J4 Net Wt.: 1 lbs. 4.5 Ozs.
Net Contents: 1 Pint (.474 liters)

SA-50 Brand Chelated
intended for use in the
prevention of foliage yellowing
deficiency of flowers, shrubs,
vegetables and trees.

USE PRECAUTIONS

THIS PRODUCT WILL SEVERELY DAMAGE
MASONRY AND PAINTED SURFACES.
Be careful not to allow spray or runoff on
masonry or painted materials. Do not mix with other
pesticides or insecticides unless compatibility is
established.

CAUTION

KEEP OUT OF REACH OF CHILDREN.

Avoid contact with skin.
Avoid breathing spray mist. Do not eat or drink.


STATEMENT OF PRACTICES

IF SWALLOWED: Drink 2 glasses of water.
Seek medical attention.

IF IN EYES: Flush eyes thoroughly with water.
If irritation persists seek medical attention.

IF ON SKIN: Rinse affected area with water.
Wash thoroughly with soap and water.

Buyer assumes all risks of use, storage, handling,
material not in strict accordance with directions and
cautions given herewith.



0 51538 019