For the prevention and control of turf diseases and the diseases of annual and perennial flowers, bedding plants, foliage plants, ground covers, plus deciduous and evergreen trees and shrubs.

ACTIVE INGREDIENT Thiophanate-methyl OTHER INGREDIENTS 100.00% **Keep Out Of Reach Of Children CAUTION**

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or when going for treatment. For 24-Hour Emergency Assistance Call Chemtrec 1-800-424-9300.

EPA Reg. No. 1001-70



Cleary Chemicals LLC 178 Ridge Road, Suite A, Dayton, (732) 329-8399 (800) 524-1662

NET CONTENTS: 30 lb. (13.6 kg.)





ACTIVE INGREDIENT Thiophanate-methyl	Keep Out Of Reach Of Children CAUTION	Bermudagrass Decline Gaeumannomyces graminis var. graminis Take-All-Patch Gaeumannomyces graminis var. avenae	6-9	Apply in mid-July or when disease symptoms first appear and repeat at 14 day intervals for suppression. Use higher rates under most severe disease expression. Follow proper agronomic recommendations to maintain plant vigor.
FIRST AID IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.		Cool Season Brown Patch Rhizoctonia cerealis Necrotic Ring Spot Leptosphaeria korrea Spring Dead Spot Leptosphaeria korrea	6-9	For prevention, apply in fall before turf has stopped all growth activity. Apply second application in early Spring when soil temperatures reach 55-60°F or when disease first appears. For curative action, apply when disease first appears in early spring and continue at 14 day intervals.
Have the product container or label with you when calling a poison control center or doct	or, or when going for treatment. For 24-Hour Emergency Assistance Call Chemtrec 1-800-424-9300.	Coprinus Snow Mold Coprinus psychromorbidus	6-9	Apply 2 treatments at 21 day intervals in late fall to early winter, with the last application made just prior to first permanent snow cover. Rotations with PCNB can be utilized.
EPA Reg. No. 1001-70	Cleary Chemicals LLC 178 Ridge Road, Suite A, Dayton, New Jersey 08810-1501 (732) 329-8399 (800) 524-1662	Dollar spot Moellerodiscus, Lanzia, Sclerotinia Large/Brown Patch	1.5-6	For prevention, use lowest rate and apply application before expected period of disease development. Apply when disease first appears and continue at 14 day intervals. Rotations with chlorothalonil, iprodione, or mancozeb (Protect) can be utilized.
HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION: Harmful if swallowed, absorbed through skin or inhaled. Causes moderate eye with soap and water after handling and before eating, drinking, chewing gum or using too personal protective Equipment (PPE) Some materials that are chemical-resistant to this product are made of any waterproof m selection chart.	ITIONARY STATEMENTS is irritation. Avoid contact with skin, eyes or dothing. Avoid breathing dust or spray mist. Wash thoroughly pacco. interials. If you want more options, follow the instructions for Category C on the EPA chemical resistance	Rhizoctonia solani Ascochyta Leaf Blight Ascochyta spp. Copper Spot Gloeocercospora sorghi Red Thread Laetisaria tuciformis Fusarium Patch Fusarium nivale Zoysia Patch Rhizoctonia solani		
Loaders, applicators and other handlers must wear: Long-sleeved shirt and long pants Shoes plus socks		Fusarium Blight Fusarium roseum and F. triticum Gray Leaf Spot (Blast)	6-9	Apply when disease first appears at 14 day intervals. Apply preventative application before expected period of disease development. Continue applications at 14 day intervals
 Chemical resistant gloves are required for applicators using hand held equipment. 		Pyricularia grisea	6-9	as needed.
USER SAFETY REQUIREMENTS: Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.		Leaf Spot Drechslera spp. Leaf, crown, and root diseases Bipolaris, Curvularia, and Exserohilum	6-9	Apply when disease first appears and make applications at 14 day intervals as needed. Rotations with chlorothalonil, iprodione, or mancozeb (Protect) are recommended under severe conditions.
USER SAFETY RECOMMENDATIONS: Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove clothing/PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.		Pink Snow Mold Michrodochium nivale	3-6	Apply middle-late Fall to early Winter before turf has stopped all growth activity or in spring during periods of thaw, or when disease symptoms first appear. Continue at 14 day intervals throughout anticipated disease period. Best results are achieved when first application is made before turf is completely dormant.
ENVIRONMENTAL HAZARDS		Stripe Smut Ustilago striiformis	6-9	Apply when disease first appears and make two applications at 14 day intervals. For prevention, apply in spring and fall.
Do not apply directly to water, or areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water.		Summer Patch Magnaporthe poae	6-9	For prevention, apply 3 applications starting late April or early May using 21 day intervals. Rotations may be used as part of the three application program. For suppression, apply at 14 day intervals when disease first appears.
DIRECTIONS FOR USE		Bentgrass Dead Spot Ophiosphaerella agrostis	6-9	For prevention, apply in early June or based upon local Extension Service recommendations. Apply at 14 day intervals as needed. Rotations and/or tank mix combinations may be used for season long disease prevention.
o not apply this product in a way that will contact workers or other persons, either directl pecific to your State or Tribe, consult the agency responsible for pesticides.	STENT WITH ITS LABELING. For use only by certified applicators and those under their immediate supervision. ly or through drift. Only protected handlers may be in the area during application. For any other requirements	*Observe the maximum individual application	on rates and maximi	um seasonal application limits in Table 2.
	JRAL USE REQUIREMENTS	Table 4: Guide to 3336 DG Lite Re	ecommended S	preader Settings*
forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains	ard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, requirements for training, decontamination, notification, and emergency assistance. It also contains specific active equipment (PPE), and restricted entry interval (REI). The requirements in this box apply to uses of this			itition of spreader and operator speed will cause variation in applied rates of material. Spreader calibration at the beginning of e

product that are covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours. Personal Protective Equipment (PPE) required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, · Coveralls over long-sleeved shirt and long pants

Chemical-resistant gloves made of any waterproof material Chemical-resistant footwear plus socks · Chemical-resistant headgear for over head exposur NON-AGRICULTURAL USE REQUIREMENTS The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this

Do not contaminate water, food, or feed by storage or disposal. Pesticide Storage: Store in the original container in a dry, temperature controlled area. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, contain/re-capture spillage and use in accordance with instruction below or dispose of in accordance with the Pesticide Disposal Instructions listed below. Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be used according to the label instructions, contact your state Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Shake empty product container to remove all possible material. When bags are

COMMERCIAL TURF AND ORNAMENTAL

empty, they may be disposed of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

held rotary spreader (e.g. bellygrinder). Application equipment should be properly calibrated prior to use. Application equipment should be thoroughly cleaned after use. If a rate range is given, use the nomeowner use and can only be used by individuals/firms licensed or registered by the state to apply ornamental or turf pest control products. No claim of compatibility with other pesticides is implied. Do not apply this product through any type of irrigation system.

Resistance Management: To avoid the development of tolerant strains of fungi, 3336 DG Lite should be used with fungicides of different modes of action. Cleary Chemical does not recommend the use of products containing thiabendazole in combination or rotation with 3336 DG Lite. These utilize similar chemistry and mode of action and can contribute to development of disease tolerance. If, after using 3336 DG Lite as recommended, the treatment is not effective, a tolerant strain of fungi may be present. Consult your local Cleary Chemical representative, your State Agricultural Experiment Station, or your State Cooperative Extension Service for proper disease identification and advice on the prompt use of some other suitable fungicide or disease control strategy. As long as recommended precautions are followed, 3336 DG Lite can remain useful for disease control.

COMMERCIAL TURF APPLICATIONS For use on all fine turf applications such as commercial, residential and public (such as home lawns, parks, athletic fields, schools, and day care centers), and golf courses (greens, tees, fairways and aprons) of cool and warm season grasses such as Bentgrass, Bluegrass, Bermudagrass, Fescue, Ryegrass, St. Augustinegrass, Zoysiagrass, or their mixtures. 3336 DG Lite is not phytotoxic to any of the above mentioned grasses when used in accordance with the label, 3336 DG Lite is to be used for the prevention and control of the diseases mentioned below. 3336 DG Lite may be applied to dry or moist leaf/soil conditions and should be activated with irrigation or natural rainfall within 24 hours for best results. If irrigated, application should be watered in by applicator with 0.1-0.25 inches of water to begin activation of product for disease control. When treating golf greens, treatment of aprons is recommended. Use the highest recommended rate under conditions of severe disease pressure. Do not graze animals on treated turf. Do not feed dippings to livestock or poultry. Not for use on turf being grown for sale or other commercial use as sod.

Use Site	Maximum application rate of 333	36 DG Lite Re-Trea	tment Interval	Comments				
Residential or Public Areas	131 Lbs/Acre (3 lbs / 1,000 s	sq.ft.)	4 days					
Golf CourseTees, Greens, Aprons	392 Lbs/Acre (9 lbs / 1,000	sq.ft.)	4 days					
Golf Course Fairways – except Florida	262 lbs/Acre (6 lbs / 1,000 s	sq.ft.)	4 days	Excludes Florida				
Golf Course Fairways – Florida Only	131 lbs/Acre (3 lbs / 1,000 s	sq.ft.)	4 days	Florida Only				
ble 2: Maximum Seasonal Application Rates Do not exceed the following amounts of product per Acre per season								
Use Site	Maximum Pounds 3336 DG Lite per Acre per Season	Pounds 3336 DG Lite per 1,000 sq.ft.	Comments					
Residential or Public Areas	523 lbs.	12 l bs.	4 Application	ns per year				

For prevention in historic areas of disease pressure, apply twice at 14 day intervals when soil temperature reaches 60°l For curative control, apply when disease first appears and continue at 14 day intervals. Rotations with chlorothalonil or triadimefon can be utilized. Anthracnose, foliar

The Andersons and DG Lite logo are registered trademarks of The Andersons, Inc. 3336 is a registered trademark of Cleary Chemicals, LLC.

Accepted 091113 Version 110614

BACK

Nursery, Greenhouse, Landscape & Interiorscape

For control of soil borne diseases effecting Bulb

Myrothecium, Penicillium, Ramularia, Rhizoctonia

The Andersons and DG Lite logo are registered trademarks of The Andersons, Inc.

3336 is a registered trademark of Cleary Chemicals, LLC.

Corms, Rhizomes, Roots, Crowns and Stems caused by: Botrytis, Cylindrocladium, Fusarium, Gliocladium,

Annual and Perennial Flowers, Bedding Plants, Foliage Plants, Ground Covers, plus Deciduous and Evergreen Trees and Shrubs

3336 DG Lite is a broad spectrum systemic fungicide which controls a variety of foliar, stem, and root diseases on a wide range of commercially important plants. 3336 DG Lite may be used for disease control and prevention programs on annual and perennial flowers, bedding plants, foliage plants, ground covers, and deciduous and evergreen trees and shrubs found in Nursery, Greenhouse, Interiorscape, and Landscape sites. Follow guidelines provided under general information for use of this product.

3336 DG Lite may be used as a broadcast, fly-on, surface, over the top, and media incorporation application for the diseases listed below. For surface and broadcast applications to established

plants apply at the rates below. For cut flowers do not exceed the maximum application rate 0.5 lbs ai/acre/application. 3336 DG Lite may be applied to dry or moist soil conditions and should be

ctivated with irrigation or natural rainfall within 24 hours for best results. If irrigated, application should be watered in by applicator with 0.1-0.25 inches of water to begin activation. For media corporation, mix at a rate of 2-3 lb/cu yd before seeding or transplanting of desired crop. Media incorporated with 3336 DG Lite may be stockpiled for later use without activation until needed.

Note: The "Directions For Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species

For preventative control, apply as media incorporation over the top application after seeding or transplanting

to propagation beds, containers, pots, or nursery or landscape beds. For curative control, apply after

seeding or transplanting when disease first appears

Note: 3336 DG Lite does not control Pythium or etridiazole, propamocarb, mono and dipotassium salts of phosphorous acid or fosetyl-Al are required for the control of *Pythium* and *Phytophthora*.

and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on Swedish Ivy (Plectranthus australis), Boston Fern (Nephrolepis exhalta), and Easter

risks referred to therein. To the extent consistent with applicable law, CLEARY CHEMICALS LLC makes no other expressed or implied warranty of fitness or merchantability or any other expressed or implied warranty. To the extent consistent with applicable law, neither CLEARY CHEMICAL nor seller shall be liable for consequential, special or indirect damages resulting from the use or seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized

NET CONTENTS: 30 lb. (13.6 kg.)



SEAL AREA NO COPY

Client: Andersons Cylinder: 27" **Distortion:** 98.557% **V**= -1.443 .067 Surface TW: N/A Slit Width: 13" Length: 27" Film Type: SWS Date: 07.06.11 Artist: wp PMS NUMBER'S CALLED OUT ON PROOF WILL

NOTE TO CUSTOMER - READ CAREFULLY Spiralkote Flexible Packaging, shall not be responsible for errors left uncorrected on this proof. Spiralkote n-compliance with local, state, or federal laws and egulations or for conflict or infringement o BE MATCHED ON PRESS.

DIRECT PLASTICS A Clondalkin Company If colour square targets are shown on this proof they <u>may</u> appear on the finished product

hey are required as part of our printing proce

however can be moved to a more appropriate location if requested.

Artwork Approval

Copy & Content Package Size and Dimensions Inside Bag Dimensions Seal Area / No Print Area Number of Colours & PMS #'s Vent Holes and Postioning Eyemark size and Postioning **UPC Number**

SEAL AREA NO COPY

Although every effort is made to produce the most accurate artwork possible, errors may occur. We do not assume any liabilities beyond the replacement of artwork. PLEASE CHECK CAREFULLY!