



Liquid Wormer 2X

(pyrantel pamoate)

NDC 30798-760-31
EACH 16 OUNCE BOTTLE
TREATS 960 LBS

Liquid Wormer 2X

(pyrantel pamoate)

5ML PER 10 LB BODY WEIGHT

Canine Anthelmintic Suspension
Removes Large Roundworms & Hookworms in Puppies & Dogs



FOR ANIMAL USE ONLY
KEEP OUT OF REACH OF CHILDREN
ANADA 200-007 APPROVED BY FDA
NET CONTENTS: 16 FL OZ (473 mL)

Manufactured for: Durvet, Inc.
100 S.E. Magellan Drive, Blue Springs, MO 64014
ISS12XB04



7 45801 01173 1

NDC 30798-760-31

EACH 16 OUNCE BOTTLE
TREATS 960 LBS

Liquid Wormer 2X

(pyrantel pamoate)

5ML PER 10 LB BODY WEIGHT

Canine Anthelmintic Suspension
Removes Large Roundworms & Hookworms in Puppies & Dogs



FOR ANIMAL USE ONLY
KEEP OUT OF REACH OF CHILDREN
ANADA 200-007 APPROVED BY FDA
NET CONTENTS: 16 FL OZ (473 mL)



0.00000

Liquid Wormer 2X

(pyrantel pamoate)

INDICATIONS FOR USE:

To prevent reinfestation of *Toxocara canis* in puppies and adult dogs and in lactating bitches after whelping.
For the removal of large roundworms, (*Toxocara canis* and *Toxascaris leonina*) and hookworms, (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies.
The presence of these parasites should be confirmed by laboratory fecal examination. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

TAKE TIME OBSERVE LABEL DIRECTIONS

IMPORTANT: SHAKE WELL BEFORE USE
WARNING: KEEP OUT OF THE REACH OF CHILDREN



.5" NON VARNISH AREA FOR LOT & EXP IMPRINT COPY

Liquid Wormer 2X

(pyrantel pamoate)

Canine Anthelmintic Suspension

DIRECTIONS FOR USE:

For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8 and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfestation. Administer one full teaspoonful (5 mL) for each 10 lbs. of body weight.
For the removal of large roundworms (ascarids) and hookworms. Administer one full teaspoonful (5 mL) for each 10 lbs. of body weight. It is not necessary to withhold food prior to treatment.
Dogs usually find this wormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption.
It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2 to 4 weeks after first treatment.

RECOMMENDED STORAGE:
STORE BELOW 30°C (86°F)
Read Accompanying Professional Information

