

Hookworms & Large Roundworms
Dog Dewormer Tablets for Large Dogs

50 Tablet Count

CAUTION

If your dog looks or acts sick, do not treat with this product.

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. Restricted Drug - Use only as directed. Not for human consumption.

KEEP OUT OF REACH OF CHILDREN CAUTION

For use only as directed.





Manufactured for: Farnam Pet Products A Division of Farnam Companies, Inc. P.O. Box 34820 Phoenix, AZ 85067-4820

For information or comments, call 1-800-234-2269 Visit our Web site at farnampet.com

Printed in the U.S.A.
Product Code: 3092 RM #: 2008928
©2007 Farnam Companies, Inc. 07-1448



NDC 017135-350-50



Dog Dewormer

For Removal of **Hookworms** in Puppies & Dogs

Ancylostoma caninumUncinaria stenocephala

For Removal of **Large Roundworms** (Ascarids)

- in Puppies & Dogs
 Toxocara canis
- Toxascaris leonina

To Prevent Reinfection of **Large Roundworms** in Puppies, Adult Dogs & Lactating Bitches After Whelping

Toxocara canis

50 Tablet Count

Active Ingredient: Each tablet contains 113.5 mg. Pyrantel base as Pyrantel Pamoate.

NADA #101-331 • Approved by FDA NET WEIGHT: 4.80 OZ



DIRECTIONS

For the removal of large roundworms (Ascarids) and hookworms, give 1 tablet for each 50 lbs. of body weight. Tablets may be broken in half to provide 1/2 tablet for 25 lbs. of body weight. The presence of these parasites should be confirmed by laboratory fecal examination. Do not withhold food from your dog prior to or after treatment.

Place tablet directly into back of mouth or conceal tablet in a small amount of food. A follow-up fecal examination should be conducted in 2 to 4 weeks after first treatment to determine the need for retreatment.

Since anthelmintics cannot be relied upon to prevent reinfection or to remove larvae not present in the intestinal tract at the time of initial treatment, for maximum control, it is recommended that puppies be treated at 2,3,4,6,8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs should be routinely treated at monthly intervals to protect against environmental *T. canis* reinfection. Retreatment of adult dogs may be necessary at monthly intervals as determined by laboratory fecal examinations or in animals kept in known contaminated quarters.

Hookworms & Large Roundworms
Dog Dewormer Tablets for Large Dogs

50 Tablet Count