FIRST AID Hold eye open and rinse slowly and gently with water for 15-20 minutes. If in eves Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. If on skin · Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. or clothing · Call a poison control center or doctor for treatment advice. If inhaled · Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. In case of medical emergencies or health and safety inquiries, or in case of fire, leaking or damaged containers or for

In case of medical emergencies or health and safety inquiries, or in case of fire, leaking or damaged containers or for product use information, call the National Pesticide Information Center at 1-877-757-4943.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

In the event of a medical or chemical emergency, contact Chemical North America 1-800-255-3924 or Worldwide Intl. + 01-813-248-0585

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Harmful if inhaled. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

1. WHAT IS THIS PACKAGE FOR?

Use ferti-lome® Dipel® Dust Biological Insecticide to kill Worms biologically. One bite of treated foliage and within minutes Worms stop feeding and eventually die. ferti-lome® Dipel® Dust Biological Insecticide is easy to use for killing Tomato Hornworms, Bagworms, Armyworms, Webworms, Gypsy Moth Larvae, Cankerworms, Loopers, Ticaterpillar, Tomato Fruitworms, Sod Webworms, Variegated Cutworms, Imported Cabbage Worms, Rindworms, Melonworms and other listed Worms.

2. HOW DO YOU USE IT

Simply dust this product on plants at the rates listed below. Dust to thoroughly cover all plant surfaces. Apply when pests appear and repeat as necessary. Apply recommended number of ounces (ozs.) below to 50 sq. ft. for vegetable and turf treatment.

BEANS, PEAS, CABBAGE, TURNIPS, CAULIFLOWER, COLLARDS, BROCCOLI AND MUSTARD GREENS: Use 1/2-1 oz. to kill Loopers and Webworms. For Hornworms, Cabbageworms, Diamondback Moths and Green Cloverworms, use 1/4-1/2 oz. For Armyworms use 1-2 ozs.

TOMATOES, CUCUMBERS, SQUASH, PEPPERS, MELONS, LETTUCE, CELERY, POTATOES, AND SPINACH: Use 1/2-1 oz. to kill Loopers, Tomato Fruitworms, Variegated Cutworms, Rindworms, Salt Marsh Caterpillars. For Hornworms use 1/4-1/2 oz. For Armyworms and Melonworms use 1-2 ozs.

TURF: To kill Sod Webworms use 2-4 ozs.

SHRUBS AND FLOWERS: Dust to thoroughly cover plant surfaces. Use to kill Loopers, Gypsy Moths, Cankerworms, Tent Caterpillars, Bagworms, Webworms, Red-Humped Caterpillars, Omnivorous Leafrollers, Tobacco Budworms, Spruce Budworms, Saddle Prominent Caterpillars, Fruit Tree Leafroller, Blackheaded Budworms, Saddleback Caterpillar, Western Tussock Moth, Diamondback Moth, Hornworms, California Oak Moths and Armyworms.

3. HERE ARE THE RESULTS YOU MAY EXPECT!

When used as directed, ferti-lome® Dipel® Dust Biological Insecticide will kill a wide variety of Leaf-eating Caterpillars that can severely damage and kill vegetables, shrubs, lawns and flowers.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

PESTICIDE STORAGE: Store in a cool and dry place inaccessible to children and pets.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: If empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Manufactured By:

VPG

*Voluntary

*Verbinstring

*Groups, Inc.

"the ferti-lome people"

FM 87 • BONHAM, TEXAS

"the ferti-lome people"
230 FM 87 • BONHAM, TEXAS 75418
EPA Reg. No. 7401-290
EPA Est. No. 7401-TX-01
Visit Us At: www.fertilome.com
Product Ouestions? 855-270-4776





ferti-lome

Dipel® Dust

Biological Insecticide

- For Vegetables, Shrubs, Lawns And Flowers
- Kills Tomato Hornworms, Bagworms, Armyworms, Webworms, Gypsy Moth Larvae, Cankerworms, Loopers, Tent Caterpillars And Other Listed Worms

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

ACTIVE INGREDIENT

Bacillus thuringiensis subsp. kurstaki strain ABTS-351
Fermentation solids, spores and insecticidal toxins....0.064%
OTHER INGREDIENTS: 99.936%
TOTAL: 100.000%

Home and Residential Use Only.

Potency: The percent active ingredient does not indicate product performance and potency rates are not federally standardized.

NET WEIGHT 1 LB. (0.453 KG)