



SUPPLEMENTAL LABELING

This supplemental label expires on June 1, 2015 and may not be used or distributed after this date.

GROUP 11 FUNGICIDE

DISARM® G Fungicide

For the control of foliar, stem and root diseases in turf and ornamentals. For commercial production and in landscape areas around residential, municipal and commercial properties, field grown ornamentals and ornamentals in greenhouses, interiorscapes and other enclosed structures.

ACTIVE INGREDIENT:

Fluoxastrobin: [(1E)-[2 [[6 (2 Chlorophenoxy) 5 fluoro 4 pyrimidinyl]oxy]phenyl] [5,6 dihydro 1,4,2 dioxazin 3 yl) methanone-O-methyloxime].....0.25%

OTHER INGREDIENTS:.....99.75%

TOTAL:100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions, worker protection standard requirements, and precautions on the EPA registered main label for **DISARM G FUNGICIDE EPA Reg. No. 66330-380** that is affixed to the container before applying. Use of DISARM G FUNGICIDE according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for DISARM G FUNGICIDE.

This supplemental label must be in the possession of the user at the time of application.

DIRECTIONS FOR APPLICATION TO ORNAMENTALS

DISARM G Fungicide is recommended for control of certain pathogens causing foliar, root and stem diseases of ornamentals. Applications can be made to plants growing in containers, benches, flats, plugs and beds in greenhouses, shadehouses, outdoor nurseries, field plantings, retail nurseries, interiorscapes, residential, public and commercial landscape areas.

DISARM G Fungicide may be applied to control foliar, soilborne, seedling, and crown diseases of production ornamentals (greenhouse, shadehouse, container grown and field grown) when applied to the surface of the growing media or when incorporated into the growing mix before planting. Make applications prior to infection as healthy roots are necessary to optimize product uptake, systemic translocation and disease protection. Surface applications of Disarm G Fungicide are not recommended where crops are watered through subsurface or flood irrigation systems.

For surface applications apply DISARM G Fungicide at 150-200 lb/Acre (3.5-4.6 lb/1,000 sq ft, 0.35 - 0.46 oz/100 sq ft). Under light to moderate disease pressure, use the lower rate (3.5 lb/1,000 sq ft). Use the higher rate (4.6 lb/1,000 sq ft) where disease pressure is heavy. Additional applications may be made at 28 day intervals as needed.

Restriction: Do not apply more than 2.2 lb active ingredient per acre per year.

Water in after application with enough water to completely wet the growing media.

DISARM G Fungicide may be incorporated into the growing mix before planting. Incorporate 18 oz Disarm G (0.0028 lb ai)

per cubic yard uniformly throughout the potting mix. If this treated potting mix is used a topdressing or soil amendment, do not use more than 1 cu ft of treated potting mix per 55.68 sq ft of soil surface.

The following diseases will be controlled by surface or potting mix applications of Disarm G Fungicide.

Foliar Diseases

Disarm G Fungicide will control certain foliar diseases of ornamentals when applications are made in advance of infection periods. The application must be made sufficiently in advance of foliar infection periods to allow the active ingredient to be translocated throughout the plants. Depending upon the plant species, the stage of growth and the environmental conditions, it may take 1-3 weeks for the fungicide to protect plants from foliar disease infection. The length of control will vary with the plant species and the pathogen.

Restriction: do not apply more than 2.2 lb of fluoxastrobin per acre per year.

Foliar Diseases (Pathogens) Controlled by Disarm G Fungicide
<p>LEAF BLIGHTS / SPOTS <i>Ascochyta</i> spp.* Alternaria Leaf Spot (<i>Alternaria</i> spp.)* Anthracnose (<i>Colletrichum</i> spp., <i>Elsinoe</i> spp.)* Cercospora Leaf Spot (<i>Cercospora</i> spp.)* Downy Mildew (<i>Peronospora</i> spp., <i>Pseudoperonospora</i> spp., <i>Plasmophora</i> spp., <i>Bremia</i> spp.)* <i>Corynespora</i> spp.* <i>Diplocarpon</i> spp.* <i>Sclerotinia</i> spp. <i>Venturia</i> spp.* Myrothecium Leaf Spot (<i>Myrothecium</i> spp.)* Septoria Leaf Spot (<i>Septoria</i> spp.)*</p>
<p>POWDERY MILDEWS <i>Erysiphe</i> spp. <i>Microsphaera azalea</i>* <i>Sphaerotheca parnosa</i>* <i>Podosphaera</i> spp.* <i>Uncinula</i> spp.*</p>
<p>RUSTS Needle Rust (<i>Melampsora</i> spp.)* <i>Phragmidium</i> spp.* <i>Puccinia</i> spp. <i>Uromyces</i> spp.*</p>
<p>FLOWER BLIGHTS Anthracnose (<i>Collectotrichum</i> spp., <i>Elsinoe</i> spp.)* Botrytis blight (<i>Botrytis</i> spp.)*</p>

Soil Borne Diseases

<p>SHOOT/STEM DISEASES (Crown Spray) Aerial/Shoot Blight (<i>Phytophthora</i> spp.)*</p>
<p>SOILBORNE DISEASES <i>Rhizoctonia solani</i> <i>Sclerotium rolfsii</i> <i>Fusarium</i> spp.*</p>

***Not for use in California**

PLANT SAFETY: DISARM G Fungicide has been shown to be safe when applied to the ornamental plants listed in the table below. However, due to the large number of genera, species and varieties of ornamental and nursery plants, it is impossible to test every variety or cultivar for tolerance to DISARM G Fungicide. Neither the manufacturer nor the seller has determined whether or not DISARM G Fungicide can be used safely on genera, species, or varieties of ornamental and nursery plants not specified on this label. The professional user should conduct small scale testing to insure plant safety prior to broad scale commercial use on plant genera and species not listed in this label.

NOTE: DO NOT USE DISARM G Fungicide ON LEATHERLEAF FERN OR OTHER FERNS GROWN UNDER NATURAL OR ARTIFICIAL SHADE.

Plants that have been shown to be tolerant to DISARM G Fungicide	
African Violet	Impatiens, Walleriana
Ageratum	Lantana
Angelonia	Lobelia
Argyranthemum	Lupine
Bacopa	Monarda
Begonia	Nemesia
Calibrachoa	Osteospermum
Chrysanthemum	Pentas
Coleus	Petunia
Dahlia	Poinsettia
Dianthus	Rose
Diascia	Scaevola
Dogwood	Snapdragon
Geranium	Torenia
Gerbera Daisy	Verbena
Hollyhock	Zinnia
Impatiens, New Guinea	

Spreader	GRANULAR SPREADER SETTINGS		
	2 lb/1,000 sq ft	3 lb/ 1,000 sq ft	4 lb/1,000 sq ft
Cyclone	3	4	4 ¼
Lely	3 II	4 II	5 II
Lesco	10	15	16
Scotts R8A/SR-1	I	K	L
Spyker	3	3 ½	4 ¼
Vicon	20	22	23

Note: Spreader settings are approximate starting points for calibrating equipment.



Arysta LifeScience North America, LLC
15401 Weston Parkway, Suite 150
Cary, NC 27513

Arysta LifeScience and the Arysta LifeScience logo registered trademarks of Arysta LifeScience, LLC

AD012213-ALLSTATES 092613