



For foliar and systemic insect control on ornamentals, and non-bearing fruit and nut trees in, nurseries and interiorscapes.

**ACTIVE INGREDIENT:**

Spirotetramat: *cis*-3-(2,5-dimethylphenyl)-8-methoxy-2-oxo-1-azaspiro[4,5]dec-3-en-4-yl-ethyl carbonate..... 22.4%

**OTHER INGREDIENTS:**..... 77.6%  
100.0%

Contains 2 pounds of spirotetramat per US gallon, or 240 grams per liter.

EPA Reg. No. 101563-130 EPA Est. No. 71106-GA-3

**STOP - Read the label before use  
KEEP OUT OF REACH OF**

**CHILDREN**

**CAUTION**

Net Contents:

250 mL (8.5 fl oz)

D00000944

84992860D 240701AV1

**envu™**

See Back Panel for First Aid Instructions and Booklet for Complete Precautionary Statements and Directions for Use.

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-424-9300  
For **PRODUCT USE** Information Call 1-800-331-2867

### FIRST AID

|   |   |
|---|---|
| <b>If on skin or clothing:</b>  | <ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>  |
| <b>If swallowed:</b>  | <ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul> |
| <b>If in eyes:</b>  | <ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first five minutes, then continue rinsing.</li></ul>   |
| <b>In case of emergency call toll free the Environmental Science U.S., LLC Emergency Response Telephone No. 1-800-424-9300. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.</b> |   |
| <b>Note To Physician:</b> No specific antidote is available. Treat patient symptomatically.   |   |

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

## **PERSONAL PROTECTIVE EQUIPMENT (PPE)**

### **Applicators and other handlers must wear:**

- Protective eyewear
- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

## **ENGINEERING CONTROLS STATEMENT**

When handlers use closed systems, or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### **User Safety Recommendations**

#### **Users should:**

- wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- remove Personal Protective Equipment immediately after handling this product.
- wash the outside of gloves before removing.
- as soon as possible, wash thoroughly and change into clean clothing.

## **ENVIRONMENTAL HAZARDS**

**For Terrestrial Use:** This pesticide is toxic to aquatic invertebrates and oysters. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. This product may contaminate water through drift of spray in wind. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

This product is potentially toxic to honey bee larvae through residues in pollen and nectar, but not to adult honey bees.

Exposure of adult bees to direct treatment or residues on blooming crops can lead to effects on honey bee larvae. See the "Directions for Use" section of this label for specific crop application instructions that minimize risk to honey bee larvae.

### **Spray Drift Reduction Management**

Do not apply when wind speed favors drift beyond the area intended for treatment. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

**Importance of Droplet Size:** An important factor influencing drift is droplet size. Select nozzles and pressure that deliver medium spray droplets as indicated in nozzle manufacturer's catalogs and in accordance with ASAE Standard S-572. Nozzles that deliver coarse spray droplets may be used to reduce spray drift provided spray volume per acre (GPA) is increased to maintain crop coverage. Low humidity and high temperature increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

**Wind Speed Restrictions:** Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed. Do not apply when winds are greater than 15 mph and avoid gusty and windless conditions. Avoiding applications when wind direction is toward an aquatic area can reduce risk exposure to sensitive aquatic areas.

**Restrictions During Temperature Inversions:** Do not make ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog. However, if fog is not present, the movement of smoke from a ground source can also identify inversions. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

**Airblast (Air Assist) Specific Recommendations:** Airblast sprayers carry droplets into the canopy via a radial or lateral directed air stream. The following drift management practices should be followed:

- Adjust deflectors and aiming devices so that spray is only directed into the canopy;
- Block off upward pointed nozzles when there is no overhanging canopy;
- Use enough air volume to penetrate the canopy and provide good coverage;
- Do not allow the spray to go beyond the edge of the cultivated area (i.e., turn off sprayer when turning at end rows);
- For applications to the outside rows, only spray inward, toward the targeted plants.

## **RUNOFF MANAGEMENT**

This product may contaminate water through runoff or drift of spray in wind. This product has a high potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

## **ENDANGERED SPECIES ADVISORY**

The use of any pesticide in a manner that may kill or otherwise harm endangered species or adversely modify their habitat is a violation of Federal law.

## **INSECT RESISTANCE STATEMENT**

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area. Kontos contains an active ingredient with a mode of action classified as a Group 23 insecticide – lipid biosynthesis inhibitor (LBI). Studies to determine cross-resistance of Group 23 insecticides with other chemical classes have demonstrated no cross-resistance. Environmental Science U.S., LLC strongly encourages that Kontos, applied alone or in tankmix combination with another Group 23 product, be applied in a block rotation or windowed approach with products from other chemical classes having a different mode of action before using additional applications of Group 23 insecticides against the same target pest. Using a block rotation or windowed approach, along with other IPM practices, is considered an effective use strategy for preventing or delaying an insect pest's ability to develop resistance to a given class of chemistry.

Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional resistance management or IPM recommendations. Also, for more information on Insect Resistance Management (IRM), visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://irac-online.org>.

## **DIRECTIONS FOR USE**

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

For foliar and systemic insect control on ornamental plants in greenhouses, nurseries, and interior plantscapes, including non-bearing fruit and nut trees (i.e., trees that will not bear fruit or nuts for one year after application).

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the REI of **24 hours** following application. **Exception:** If the product is applied by drenching the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water is:

- Long-sleeved shirt and long pants
- Protective eyewear
- Chemical-resistant gloves
- Shoes plus socks

### **APPLICATION INFORMATION**

Kontos is a Suspension Concentrate formulation. Following application to plant foliage, Kontos is fully systemic, moving through phloem and xylem to all plant tissues including new shoot, leaf and root growth. Spray adjuvants with spreading / penetrating characteristics may improve leaf uptake and systemic concentration of active ingredient. The active ingredient contained in Kontos is active primarily by ingestion. Fecundity of adult insects and mites may be reduced. Make applications as preventative treatments or to coincide with early threshold levels in developing insect and mite populations. Kontos can be applied by ground equipment (including hand-held application equipment) or through chemigation.

### **FOLIAR SPRAY APPLICATIONS**

Foliar applications must be made using properly calibrated ground sprayers (including hand-held and backpack sprayers), through properly designed, sprinkler-type, chemigation equipment (See Chemigation Application directions below).

### **Application through irrigation systems**

Kontos may be applied at rates on this label either alone or in tank mixture with other pesticides and chemicals registered for application through irrigation systems. The normal dilution ratio is 1:10 to 1:200, depending on the system. Always meter the product into the irrigation water during the first part of the irrigation cycle. The product may be mixed separately prior to injection. Agitation may be necessary if the mixture is allowed to stand more than 24 hours.

Remove scale, pesticide residue and other foreign matter from the tank and entire irrigation system.

Apply Kontos only through microirrigation (individual spaghetti tube), drip irrigation, overhead irrigation, ebb and flood, or hand-held or motorized calibrated irrigation equipment.

Do not apply this product through any other type of irrigation system. Crop injury or lack of effectiveness can result from nonuniform distribution of treated water.

If you have any questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or a person who is under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

### **SAFETY DEVICES FOR IRRIGATION SYSTEMS CONNECTED TO PUBLIC WATER SUPPLIES:**

If the source of water for your irrigation system is a public water supply, follow the instructions below.

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from

- the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
  6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
  7. Do not apply when wind speed favors drift beyond the area intended for treatment.

#### **SAFETY DEVICES FOR IRRIGATION SYSTEMS NOT CONNECTED TO A PUBLIC WATER SUPPLY:**

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where the pesticide distribution is adversely affected.
6. Systems must use a metering pump such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

**Ebb and Flood Application:** To assure accurate uptake, it is recommended that, prior to treatment, a minimum of 10 plants be brought up to a known field capacity and allowed to dry out for one or two days. Rewet these plants to determine how much water on average each plant will absorb to bring it back to field capacity. Use the volume absorbed per plant (keeping pot sizes uniform) multiplied by the number of pots being treated. Add to this volume a required minimum amount to flood the smallest treatment area. This process should minimize the return back to the storage tank. Re-use the returned volume with subsequent irrigation or nutrients on the same plants.

## **MIXING INSTRUCTIONS**

### **COMPATIBILITY / MIXING / ORDER-OF-MIXING**

Observe all cautions and limitations on labeling of all products used in mixtures.

Kontos is physically and biologically compatible with many registered pesticides and fertilizers or micronutrients. When considering mixing Kontos with other pesticides, or other additives, first contact your supplier for advice. For further information, contact your local Environmental Science U.S., LLC representative. If your supplier and Environmental Science U.S., LLC representative have no experience with the combination you are considering, conduct a test to determine physical compatibility. To determine physical compatibility, add the recommended proportions of each chemical with the same proportion of water as will be present in the chemical supply tank, into a suitable container, mix thoroughly, and allow the mixture to stand for five minutes. If the combination remains mixed, or can be readily remixed, the mixture is considered physically compatible.

The proper mixing procedure for Kontos alone or in tank mix combinations with other pesticides is:

- 1) Fill the spray tank 1/4 to 1/3 full with clean water;
- 2) While recirculating and with the agitator running, add any labeled amounts of Kontos. Allow time for thorough mixing;
- 3) Continue to fill spray tank with water until 1/2 full;
- 4) Add any other wettable powder (WP) or wettable granules (WG) products;
- 5) Allow enough time for thorough mixing of each product added to tank;
- 6) If applicable, add any remaining tank mix components: emulsifiable concentrates (EC), fertilizers, micronutrients, spray adjuvants.
- 7) Fill spray tank to desired level and maintain constant agitation to ensure uniformity of spray mixture.

## **CROP TOLERANCE**

Kontos has been evaluated for phytotoxicity on a wide range of ornamental plants. However, due to the large number of species and varieties of ornamental plants, it is impossible to test every one for tolerance to Kontos. Prior to commercial use, determine if Kontos can be used safely. In a small area, test the rates on a small number of plants for phytotoxicity prior to widespread use. Before using Kontos in tank mixture with other products and adjuvants, test the mixture on a small number of plants for phytotoxicity prior to widespread use.

If you have any questions, consult your local Environmental Science U.S., LLC representative, or call the Environmental Science U.S., LLC customer service hotline at 1-800-424-9300.

Kontos is not recommended for use on the following varieties: geraniums (*Pelargonium* spp.), orchids, hoya, *Dracaena*, *Cordyline*, *Schefflera*, neanthebella palm, and ferns.

Do not make more than one application per season to *Hydrangea*, *Impatiens* spp., crotons (*Codiaeum* spp.), *Fuschia* hybrids, *Petunia*, *Peperomia*, stock, or cyclamens (*Cyclamen* spp.).

**PRODUCED FOR**  
**Environmental Science U.S., LLC**  
**5000 CentreGreen Way, Suite 400**  
**Cary, NC 27513**

*(continued in back leaflet)*

**envu™**





GROUP 23 INSECTICIDE

 **KONTOS**<sup>®</sup>

**ACTIVE INGREDIENT:**

Spirotetramat: *cis*-3-(2,5-dimethylphenyl)-  
8-methoxy-2-oxo-1-azaspiro[4,5]dec-3-  
en-4-yl-ethyl carbonate..... 22.4%

**OTHER INGREDIENTS:** ..... 77.6%  
100.0%

**Contains 2 pounds of spirotetramat per US  
gallon, or 240 grams per liter.**

EPA Reg. No. 101563-130

EPA Est. No. 71106-GA-3

**STOP - Read the label before use  
KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See Back Panel for First Aid Instructions and  
Booklet for Complete Precautionary  
Statements and Directions for Use.

**Net Contents:**  
**250 mL (8.5 fl oz)**  
**D00000944**  
84992860D  
240701AV1

PRODUCED FOR  
Environmental Science  
U.S., LLC  
5000 CentreGreen Way,  
Suite 400  
Cary, NC 27513

PULL HERE TO OPEN 

*(continued from front leaflet)*



See Back Panel for First Aid Instructions  
and Booklet for Complete Precautionary  
Statements and Directions for Use.



(01)08101121001340

84507709D 221216AV2

### CONTAINER DRENCH APPLICATIONS - SMALL CONTAINERS

For application only to ornamental plants in greenhouses, nurseries, and interior plantscapes, including non-bearing fruit and nut trees (i.e., trees that will not bear fruit or nuts for one year after application), using soil drench, micro-irrigation, drip irrigation, overhead irrigation, ebb and flood irrigation, or hand-held motorized irrigation equipment. Do not apply more than 25 fl oz (750 ml, 0.4 lb of active ingredient) per acre of nursery per season.

| Pest  | Use Pattern  | Dosage  |  | Remarks   |
|---|--|---|--|---|
| Adelgids<br>Aphids<br>Leafhoppers<br>Mealybugs<br>Psyllids<br>Rust Mites <sup>1</sup><br>Scales (crawlers)<br>Spider Mites <sup>1</sup><br>Spittlebugs<br>Tarsonemid Mites <sup>1</sup><br>Thrips (immature) <sup>2</sup><br>Whiteflies | Containerized Plants<br><br>Herbaceous species, 1-2 plants per pot | Container Size (inches)<br><br>2<br>3<br>4<br>5<br>6<br>7<br>8<br>9<br>10<br>11<br>12 | No. pots treated with 1.7 fl oz – 3.4 fl oz (50-100 ml) of product<br><br>3000<br>2000<br>1500<br>1200<br>1000<br>850<br>750<br>650<br>600<br>550<br>500 | Exact drenching volume is dependent upon pot size, potting medium, and plant type. Apply product in sufficient volume of water to wet the potting medium, without loss of liquid from the bottom of the container.<br><br>Follow application with moderate irrigation. Irrigate carefully during the next 10 days in order to avoid loss of active ingredient from the bottom of the container.<br><br>The mode of action of Kontos is inhibition of lipid biosynthesis (IRAC Group 23 Insecticide). If retreatment is necessary, treat using a product with an alternative mode of action. |

*(continued)*

**CONTAINER DRENCH APPLICATIONS - SMALL CONTAINERS** (continued)

| Pest  | Use Pattern  | Dosage   |  | Remarks   |
|---|--|--|--|---|
| Adelgids<br>Aphids<br>Leafhoppers<br>Mealybugs<br>Psyllids<br>Rust Mites <sup>1</sup><br>Scales (crawlers)<br>Spider Mites <sup>1</sup><br>Spittlebugs<br>Tarsonemid Mites <sup>1</sup><br>Thrips (immature) <sup>2</sup><br>Whiteflies | Containerized<br>Plants<br><br>Woody<br>perennials | Container<br>Size (inches)<br><br>2<br>3<br>4<br>5<br>6<br>7<br>8<br>9<br>10<br>11<br>12 | No. pots treated<br>with<br>1.7 fl oz – 3.4 fl oz<br>(50-100 ml) of<br>product<br><br>2000<br>1350<br>1000<br>800<br>650<br>550<br>500<br>450<br>400<br>350<br>300 | Exact drenching volume is dependent upon pot size, potting medium, and plant type. Apply product in sufficient volume of water to wet the potting medium, without loss of liquid from the bottom of the container.<br><br>Follow application with moderate irrigation. Irrigate carefully during the next 10 days in order to avoid loss of active ingredient from the bottom of the container.<br><br>The mode of action of Kontos is inhibition of lipid biosynthesis (IRAC Group 23 Insecticide). If retreatment is necessary, treat using a product with an alternative mode of action. |

### CONTAINER DRENCH APPLICATIONS – LARGE CONTAINERS

For application only to ornamental plants in greenhouses, nurseries, and interior plantscapes, including non-bearing fruit and nut trees (i.e., trees that will not bear fruit or nuts for one year after application), using soil drench, micro-irrigation, drip irrigation, overhead irrigation, ebb and flood irrigation, or hand-held motorized irrigation equipment. Do not apply more than 25 fl oz (750 ml, 0.4 lb of active ingredient) per acre of nursery per season.

| Pest  | Use Pattern             | Dosage                         |  | Remarks   |
|---|-------------------------|--------------------------------|--|---|
| Adelgids<br>Aphids<br>Leafhoppers<br>Mealybugs<br>Psyllids<br>Rust Mites <sup>1</sup><br>Scales (crawlers)<br>Spider Mites <sup>1</sup><br>Spittlebugs<br>Tarsonemid Mites <sup>1</sup><br>Thrips (immature) <sup>2</sup><br>Whiteflies | Containerized<br>Plants | Container<br>Size<br>(gallons) | No. pots treated<br>with<br>1.7 fl oz – 3.4 fl oz<br>(50-100 ml) of<br>product | Exact drenching volume is dependent upon pot size, potting medium, and plant type. Apply product in sufficient volume of water to wet the potting medium, without loss of liquid from the bottom of the container.<br><br>Follow application with moderate irrigation. Irrigate carefully during the next 10 days in order to avoid loss of active ingredient from the bottom of the container.<br><br>The mode of action of Kontos is inhibition of lipid biosynthesis (IRAC Group 23 Insecticide). For resistance management purposes, if retreatment is necessary, treat using a product with an alternative mode of action. |

<sup>1</sup> Mites - If populations are heavy at the time of application, control may not be achieved rapidly enough to prevent economic damage to the plant. Make applications preventatively, or when populations are first detected, use the higher dosage (3.4 fl oz of product). Kontos will not control heavy populations of existing mites. If a second miticide application is necessary to achieve control, use a product with an alternative mode of action.

<sup>2</sup> Thrips - Full control of thrips will be observed only on foliage; thrips in buds will be suppressed; thrips controlled will only be immature thrips. If populations are heavy at the time of application, control may not be achieved rapidly enough to prevent economic damage to the plant. Make applications preventatively, or when populations are first detected on foliage. For thrips control on woody perennials use the higher dosage (3.4 fl oz product).

## FOLIAR APPLICATIONS FOR ORNAMENTALS GROWN IN GREENHOUSES

For foliar insect control in greenhouses.

### USE RESTRICTIONS

Do not apply more than 25 fl oz (750 ml, 0.4 lb of active ingredient) per acre of nursery per season.

| CROP   | PEST   | DOSAGE  | REMARKS   |
|--|--|---|---|
| Flowers<br>Foliage Plants<br>Groundcovers<br>Ornamentals in flats and plugtrays  | Adelgids<br>Aphids<br>Leafhoppers<br>Mealybugs <sup>1</sup><br>Psyllids<br>Scales (crawlers)<br>Spider Mites <sup>2</sup><br>Spittlebugs<br>Tarsonemid Mites <sup>2</sup><br>Thrips <sup>3</sup><br>Whiteflies | 1.7 fl oz – 3.4 fl oz<br>(50-100 ml) of<br>product /<br>100 gallons of<br>water<br>or<br>0.051 fl oz (1.5 ml)<br>– 0.1 fl oz (3.0 ml)<br>of product / 3-gal<br>of water | <p><b>Foliar applications:</b> Start treatments prior to establishment of high pest population and reapply on an as-needed basis.</p> <p>Apply when pests first appear or when damage is first noticed. Spray thoroughly.</p> <p>The addition of a spreader sticker may improve efficacy.</p>   |
| Shrubs<br>Evergreens<br>Trees - including<br>Non-bearing Fruit and<br>Nut Trees<br>(Non-bearing fruit and<br>nut trees are those trees<br>that will not bear fruit or<br>nuts for one year after<br>application) |  |   | <p>Make ground applications to foliage in a minimum of 10 gallons of spray solution per acre.</p> <p>Make airblast application in a minimum of 50 gallons of spray solution per acre.</p> <p>If concentrate or mist type spray equipment is used, use an equivalent amount of product on the area sprayed as would be used in a dilute application.</p> <p>Reapply at 14-28 days interval, if needed.</p> |

## FOLIAR APPLICATIONS FOR ORNAMENTALS GROWN OUTSIDE GREENHOUSES

For foliar insect control around greenhouses, field grown nurseries and container stocks, outdoor ornamentals, and ornamentals grown in flats, benches or beds.

### USE RESTRICTIONS

Do not apply more than 25 fl oz (750 ml, 0.4 lb of active ingredient) per acre of nursery per season.

Do not apply more than 5.1 fl oz of product (153 ml, 0.081 lb of active ingredient) per acre per application to outdoor ornamentals.

Do not apply until after petal fall to stone fruit, pome fruit, and tree nuts.

| CROP  | PEST   | DOSAGE   | REMARKS  |
|---|--|--|--|
| Flowers<br>Foliage Plants<br>Groundcovers<br>Ornamentals in flats<br>and plugtrays  | Adelgids<br>Aphids<br>Leafhoppers<br>Mealybugs <sup>1</sup><br>Psyllids<br>Scales (crawlers)<br>Spider Mites <sup>2</sup><br>Spittlebugs<br>Tarsonemid | 1.7 fl oz – 3.4 fl<br>oz (50-100 ml)<br>of product / 100<br>gallons of water<br>or<br>0.051 fl oz (1.5 ml)<br>– 0.1 fl oz (3.0 ml)<br>of product / 3-gal<br>of water | <b>Foliar applications:</b> Start treatments prior to establishment of high pest population and reapply on an as-needed basis. Apply when pests first appear or when damage is first noticed. Spray thoroughly. The addition of a spreader sticker may improve efficacy. Make ground applications to foliage in a minimum of 10 gallons of spray solution per acre. Make airblast application in a minimum of 50 gallons of spray solution per acre. If concentrate or mist type spray equipment is used, use an equivalent amount of product on the area sprayed as would be used in a dilute application. Reapply at 14-28 days interval, if needed. |
| Shrubs<br>Evergreens<br>Trees - including Non-bearing<br>Fruit and Nut Trees<br>(Non-bearing fruit and nut<br>trees are those trees<br>that will not bear fruit or<br>nuts for one year after<br>application) | Mites <sup>2</sup><br>Thrips <sup>3</sup><br>Whiteflies  |  |  |

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Kontos is packaged in poly-ethylene containers. Do not allow product or containers to freeze. Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking, invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away. You may contact the Environmental Science U.S., LLC Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Environmental Science U.S., LLC Emergency Response Telephone No. (800) 424-9300.

**Pesticide Disposal:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Handling:** Non-refillable container. Do not reuse or refill this container.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**Container Handling:**

**Rigid Non-refillable Containers that are too large to shake (i.e., with capacities greater than 5 gallons or 50 lbs)**

Non-refillable container. Do not reuse or refill this container. Refer to Bottom Discharge IBC or Top Discharge IBC, Drums, Kegs information as follows.

**Bottom Discharge IBC (e.g. - Schuetz Caged IBC or Snyder Square Stackable)**

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at

### **STORAGE AND DISPOSAL** *(continued)*

least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

#### **Top Discharge IBC, Drums, Kegs (e.g. – Snyder 120 Next Gen, Bonar B120, Drums, Kegs)**

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

## **IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of liability before using this product.

If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, because of manner of use and other factors beyond Environmental Science U.S., LLC control it is impossible for Environmental Science U.S., LLC to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ENVIRONMENTAL SCIENCE U.S., LLC MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Environmental Science U.S., LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ENVIRONMENTAL SCIENCE U.S., LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES, RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT ENVIRONMENTAL SCIENCE U.S., LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.

Envu and the Envu logo are trademarks and Kontos® is a registered trademark owned by Environmental Science U.S., LLC or one of its affiliates.

**PRODUCED FOR**  
**Environmental Science U.S., LLC**  
**5000 CentreGreen Way, Suite 400**  
**Cary, NC 27513**

The logo for Environmental Science U.S., LLC, featuring the word "envu" in a lowercase, bold, sans-serif font. The letters are black and have a slight shadow effect, giving them a three-dimensional appearance. The logo is positioned in the bottom right corner of the page.







## FIRST AID

### **If on skin or clothing:**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

### **If swallowed:**

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

### **If in eyes:**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first five minutes, then continue rinsing.

**In case of emergency call toll free the Environmental Science U.S., LLC Emergency Response Telephone No. 1-800-424-9300. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.**

**Note To Physician:** No specific antidote is available. Treat patient symptomatically.

PULL HERE TO OPEN ▲

84507709D 240701AV1