CENTRAL Garden & Pet

Farnan, the Farnam design, IverCare,
Guer-Grip and red to yellow color
gradation are trademarks of
Farnam Companies, Inc.
Central Garden & Pet and the Central Garden
& Pet Gesign are trademarks of Central
Salden & Pet Salden Salden

Product Code: 100504274
11-2330 • 99.0000263.01 • rev 12/11

© 2002, 2005, 2006, 2010, 2011 Farnam Companies, Inc.

## www.ivercare.com

Distributed by:
P.O. Box 34820
Phoenix, AZ 85067-4820 U.S.A.

Environmental Safety: Ivermectin and excreted ivermectin residues may adversely affect aquatic organisms. Do not contaminate ground or surface water. Dispose of the syringe in an approved landfill or by incineration.

 $\label{precentions: werCare* (ivermectin) Paste 7.87\% has been formulated specifically for use in horses only. This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.$ 

befrain from smoking and eating when handling, Wash hands after use, Avoid contact with eyes. The Material Safety Data Sheet (MSDS) contains more indromation, or safety information. To report adverse reactions in users, to obtain more information, or to obtain a MSDS, contact Farnam Companies, inc. at 800-234-2269

## WARNING: Do not use in horses intended for human consumption. Not for use in humans. Keep this and all drugs out of reach of children.

NAMEL SAFETY: IverCare® (ivermectin) Paste 1.87% may be used in horses of all agoes, including mares at any stage of pregnancy. Stallions may be treated without agoes, including mares at any stage of pregnancy. Stallions

PARASITE CONTROL PROGRAM: All horses should be included in a regular parasite control program with particular attention being paid to mares, foals and particular attention being paid to maree, foals and treatment repeated as appropriate. Consult your veterinarian for a control program to meet your specific needs. Veermectin) Paste 1.87% effectively controls of meet your specific needs. Veermectin) Paste 1.87% effectively controls characterioral paste 1.87% effectively controls gastrointestinal nematodes and bots of horses. Regular treatment will reduce the characterior pasterior and pasterior stream of the pasterior pasterior pasterior particular pasterior pasterior pasterior particular particular pasterior particular pasterior pasterio

פרכחותם מונכו מספונות.

DOSAGE AND ADMINISTRATION: This syringe contains sufficient paste to trest one 1500 lb horse at the recommended dose rate of 91 mag ivermectin per lb (200 mos 1500 lb horse at the recommended dose rate of 91 mog ivermectin per lb (200 mog/kg) body weight. Each weight, (1) While holding plunger, turn the knurled ring paste to treat 250 lb body weight. (1) While holding plunger, turn the the barrel is at on the plunger 1/4 turn to the left and slide it so the side nearest the barrel is at the prescribed weight marking. (2) Lock the ring in place by masking a 1/4 turn to the prescribed weight marking. (3) Lock the ring in place by Maske sure that the horse's mouth contains no feed. (4) Remove the cover from the fip of the syringe. (5) Insert the syning into the horse's mouth at the space between the leeth. (6) Depress the plunger as far as it will go, depositing paste on the barrel the tongue. (7) Immediately raise the horse's head for a few paste on the barrel to the control of the tongue. (7) Immediately raise the horse's head for a few

**досекся** ab

cutaneous third-stage larvae; Dermatitis caused by neck threadworm microfilariae, Strongyloides westert, **Summer Sores** caused by Habronema and Draschia spp. tourth-stage larvae) – Dictyocaulus arntieldi; Intestinal Threadworms (adults) – Gasterophilus spp. including G. intestinalis and G. nasalis, Lungworms (adults and Morms (adults) – Habronema muscae; Bots (oral and gastric stages) ednorum; Hairworms (adults) - Inchostrongylus axer; Large-mouth Stomach Oxyunis equi; Ascarids (adults and third- and tourth-stage larvae) - Parascaris Strongyles – Fourth-stage larvae; Pinworms (adults and fourth-stage larvae) -C. goldi, C. longibursatus and C. minutus, and Petrovinema poculatum, Small Cylicodontophorus spp.; Cylicostephanus spp. including C. calicatus, spp. including C. insigne, C. leptostomum, C. nassatus and C. brevicapsulatus, Cyathostomum spp. including C. catinatum and C. pateratum; Cylicocyclus compounds) – Coronocyclus spp. including C. coronatus, C. labiatus and C. labratus; Small Strongyles (adults, including those resistant to some benzimidazole class including I brevicauda and I serratus, and Craterostomum acuticaudatum; vessels), S. edentatus (also tissue stages), S. equinus, Triodontophorus spp. Large Strongyles (adults) - Strongylus vulgaris (also early forms in blood

treatment and control of the following parasites in horses.

INDICATIONS: Consult your vetennarian for assistance in the diagnosis, treatment, and control of parastitism, VerCare® (ivermectin) Paste 1.87% provides effective

Non-slip thumb "saddle" for smoother action

INFORMATION FOR HORSE OWNERS: Swelling and itching reactions after treatment with Ivermechin Paste have occurred in horses carrying heavy infections of neck threadworm microfilariae dying in large numbers. Symptomatic treatment may be advisable. Consult your veterinarian should any such reactions occur. Healing of summer sores involving extensive tissue changes may require other appropriate threspy in conjunction with treatment with Ivermectin Paste. Beinfection, and threspy in conjunction with treatment with Ivermectin Paste. Beinfection, and measures for its prevention, should also be considered. Consult your veterinarian if

the condition does not improve. Store at 20°C to 25°C (68°F to 77°F), excursions



permitted between 15°C and 30°C (between 59°F and 86°F).



With Sure-Grip

Syringe

NDC 17135-757-21



VERCARE

WESTER 1.87%

Extended Sure-Grip™ for better control

Anthelmintic and Boticide Removes worms and bots with a single dose.

Contents will treat up to 1500 lb body weight

PROUDLY MADE IN THE USA

Net Wt. 0.26 oz (7.30 g) ANADA 200-390, Approved by FDA



IVERCARE® (ivermectin) Paste 1.87%