

ACTIVE INGREDIENT: Deltamethrin: .. OTHER INGREDIENTS:96.0%

TOTAL:100.0%

DO NOT ALLOW CHILDREN TO PLAY WITH COLLAR

CAUTION See Label Booklet on back of Tin for Additional Precautionary Statements, Directions for Use and First Aid

NET WEIGHT: 2.31 oz (Two 1.16 oz Collars)

This product contains the toxic inert ingredient Bis(2-ethylhexyl) adipate (DEHA) at 18.0% Two collars provide up to a total of 12-month flea and tick protection One collar provides 6-month flea and tick protection For use on Dogs 12 weeks of age and older

READ ENTIRE LABEL BEFORE EACH USE USE ONLY ON DOGS PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Do not open protective pouch inside of the tin until ready to use. Do not let children play with this collar. Do not use on puppies under 12 weeks. Do not use on cats. Consult a veterinarian before using this product on debilitated, aged, pregnant, medicated, or nursing dogs. Sensitivities may occur after using ANY pesticide products for dogs. If signs of sensitivity occur, remove collar and bathe your dog with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. This collar is intended for use only as an insecticide and is not to be taken internally by humans or animals.

Manufactured for:



EPA Reg. No. 91300-4-53883 EPA Est. No. 091645-FRA-001



194167





FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical treatment information, you may contact SafetyCall at 1-844-685-9173, 24 hours a day, 7 days a week.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not open until ready to use. Dust may form on collar during storage. Do not get dust in mouth or eyes. Only one collar is to be worn by your dog at any one time.

Remove the collar from its packaging, separate the Peel-Away Comfort Strip and place in the trash. Place collar around dog's neck and buckle in place with the main clasp. Feed excess collar through the remaining two clasps for extra hold and a clean fit. Adjust the collar for a snug fit to create contact between the collar and the dog's hair/coat for better product release and diffusion. A properly fitted collar will allow two fingers to be placed between the collar and dog's neck. The elasticity of the collar is designed to stretch in the event that the dog becomes entangled. Leave an excess of 1-2 notches (l/4 to % inch) past the last clasp. Adjust periodically for growing dogs. Cut off and dispose of excess length. Wash hands after placing collar on dog. Replace collar every 6 six months.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Pesticide Storage: Store this product in a cool, dry place away from children and pets. Pesticide Disposal and Container Handling: Do not reuse container or used collar. Dispose of in trash.

FLEE Flea & Tick Collar. This collar kills fleas and ticks for up to 6 months. Collar is a double layer sustained-release system containing deltamethrin combining the insecticidal actions of vapors and powders working in three ways to provide head to tail protection. Thus, the collar has the effect of simultaneous usage of a spray and powder in the control of fleas and ticks, including ticks that may carry Rocky Mountain spotted fever. Use FLEE Flea & Tick Collar year-round for most effective control. Replace every 6 months.