

PEEL BACK

FORCE OF NATURE™ INSECTICIDE

may be used indoors or outdoors as a contact spray for flying and crawling pests commonly found in and around the home. This product contains Pyrethrum, derived from a Chrysanthemum flower, to give fast and effective kill on contact. Pyrethrum will also "flush" hiding insects, which then allows you to spray them for immediate contact kill.



Department of Agriculture
STATE OF HAWAII

PRECAUTIONARY STATEMENTS

LICENSED

PERIOD 2015-2017 Lic. No.

8158.3

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use this product in or on electrical equipment because of shock hazard.

EASY TO USE • NO MIXING OR MEASURING • FOR INDOOR OR OUTDOOR USE

BUSH DOCTOR®

FORCE OF NATURE™



**ACTIVE
INGREDIENTS:**
Pyrethrins..... 0.02%
*Piperonyl butoxide..... 0.20%
OTHER INGREDIENTS..... 99.78%
TOTAL 100.00%
* (butylcarbitol) (6-propylpiperonyl)
ether and related
compounds

**KILLS ON CONTACT
INSECTICIDE**

KEEP OUT OF REACH OF CHILDREN

CAUTION

GOT FUNGUS GNATS?

GET FORCE OF NATURE™

EPA Registration No. 1021-1801-71996

EPA Establishment No. 71996-CA-001

32 FLUID OUNCES (946 mL) NET WT 2 LB (0.907 kg)

PEEL BACK

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Peel back side panels for complete Directions for Use.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in a cool, dry area. Always store pesticides in the original container. Store in an upright position only. Store with spray nozzle securely closed. Do not store with spray nozzle below level of product in container. Store away from food and pet food.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

— FIRST AID —

IF SWALLOWED:

- Immediately call a Poison Control Center or doctor.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Have person sip a glass of water if able to swallow.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.

DISTRIBUTED BY

United Compost & Organics
P.O. Box 787, Arcata, CA 95518
— HUMBOLDT COUNTY —

Call 1-800-4 FOXFARM



FNINSECT-010T_01

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

—SHAKE WELL BEFORE USING—

Do not allow adults, children, or pets to enter the treated area until sprays have dried. Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift. Do not water the treated area to the point of run-off. Do not wet articles to the point of runoff or drip. Do not use treated article until spray has dried. Remove pets, birds, and cover fish aquariums before spraying. Do not make applications during rain. Do not wet plants to the point of runoff or drip. Remove or cover exposed food and drinking water before application. Remove or cover dishes, utensils, food processing equipment, and food preparation surfaces, or wash them before use.

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted use: application to vegetation.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

Bush Doctor® Force of Nature™ Insecticide may be used indoors or outdoors as a contact spray for flying and crawling pests commonly found in and around the home. This product contains Pyrethrum, derived from a Chrysanthemum flower, to give fast and effective kill on contact. Pyrethrum will also “flush” hiding insects, which then allows you to spray them for immediate contact kill. Adjust spray nozzle to give a coarse, wet spray.

FOR INDOOR HOUSEPLANTS

Phytotoxicity Note: Plant safety is an important consideration when using insecticides. However, it is not possible to evaluate the potential adverse effects of this product on numerous plant varieties that may react differently to insecticides in different growth stages or under varying environmental conditions. Before making widespread applications of this product treat a limited number of plants and observe for adverse effects over a one to 10 day period.

HOW TO APPLY:

Wearing household latex or rubber gloves, use this ready-to-use bottle to apply product. Water plant thoroughly. Turn spray nozzle to the on position, and hold spray nozzle about 12” from the plants. Spray lightly, covering insects on upper and lower leaf surfaces until surfaces are slightly wet. Avoid over-saturation. Turn off any circulating fans to avoid drift and contact with eyes and skin. Allow liquid to dry in treated areas. Wash hands thoroughly after handling and before eating or smoking. Do not allow spray to drift to non-target plants or water supplies. Avoid contamination of food and feedstuffs.

WHEN TO APPLY:

Use at first signs of insect infestation, preferably before damage and feeding occurs. Do not wait until plants are heavily infested.

WHERE TO USE:

Use on common houseplants.

Use Restrictions

- Do not water the treated area to the point of run-off.
- Do not reapply within 3 days except under extreme pest pressure.
- In case of extreme pest pressure, do not reapply within 24 hours.
- Do not apply more than 1 time per day.
- This product is not for use on plants being grown for sale or other commercial use.

PRODUCT FACTS:

Kills: Common insects found on House Plants, including: Aphids, Fungus Gnats, Mealybugs, Mites, Scale, Spider Mites, Springtails, Thrips (exposed), Whiteflies and Other Listed Insects.

Where to Use: Common Houseplants

Amount to Use: Spray lightly covering upper and lower leaf surfaces.

When to Use: At the first signs of insect infestation. Can be used up to the day of harvest. Kills eggs during dormant stage.

INSECTS:

On Houseplants: Ants, Aphids, Army worms, Asparagus Beetles, Blister Beetles, Cabbage Loopers, Cabbage worms, Caterpillars, Colorado Potato Beetles, Corn Earworms, Cucumber Beetles, Cutworms, Diamondback Moth Larvae, Earwigs, Fall Cankerworms, Fireworms, Flea Beetles, Gnats (including Fungus Gnats), Harlequin bugs, Inchworms, Japanese Beetles, Lace Bugs, Leafhoppers, Leafrollers, Leafminers, Leaf-tiers, Mealybugs, Mexican Bean Beetles, Mites, Rose Chafer, Scale, Spiders, Spider Mites, Spring Cankerworms, Springtails, Stink Bugs, Thrips (Exposed), and Whiteflies.

— OUTDOORS —

Vegetables (including in-home greenhouses): Arugula, Beets, Broccoli, Brussels Sprouts, Cabbage, Carrots, Chinese Cabbage (Bok Choy), Chinese Mustard Cabbage (Gai Choy), Cabbage (Napa), Cauliflower, Cucumber, Collards, Cilantro, Celery, Eggplant, Edible Gourds, Endive, Japanese Radish (Daikon), Lettuce (Head and Leafy), Okra, Melons (including hybrids-Cantaloupe, Casaba, Crenshaw, Honeydew, Muskmelon), Parsley, Parsnips, Potato, Peppers (including Bell, Chili, Cooking, Pimentos, Sweet), Pumpkins, Rhubarb, Rutabaga, Radish, Swiss Chard, Squash (Winter and Summer), Spinach, Tomatillo, Tomatoes, Watermelon, and Zucchini.

Small Fruits and Berries: Blackberries, Blueberries, Currants, Dewberries, Elderberries, Gooseberries, Grapes, Huckleberries, Olallieberries, Raspberries (Black and Red), Strawberries, and Youngberries.

Fruits: Apples, Apricots, Crabapples, Cherries (Sweet and Sour), Citrus Hybrids, Grapefruit, Loquat, Lemons, Limes, Nectarines, Oranges (Sweet and Sour), Oriental Pears, Pears, Pummelos, Peaches, Plums, Prunes, and Quince.

For Use on All Ornamental Plants (such as Roses, Hyacinth, Gladiolus, Freesia). All Annuals (such as Cosmos, Celosia, Ageratum, Impatiens). All Perennials (such as Gazania, Cineraria, Chrysanthemum, Zinnia). All Shrubs (such as Mountain Laurel, Juniper, Privet, Mock Orange). **All Shade Trees** (such as Oak, London Plane, Maple, Sycamore, Mulberry), and All Vines (such as Trumpet Creeper, Creeping Lantana, Wisteria, Honeysuckle).

For the Control of: Ants, Aphids, Armyworms, Asparagus Beetles, Blister Beetles, Brown Marmorated Stink Bugs (BMSB), Cabbage Loopers, 12-Spotted Cucumber Beetles, Colorado Potato Beetles, Corn Earworms, Crickets, Cross-striped Cabbageworms, Diamondback Moth Larvae, Fireworms, Flea Beetles, Fruit Worms, Fruit Tree Leafrollers, Grape Leafhoppers, Gypsy Moths (larvae and adults), Harlequin Bugs, Tomato Hornworms, Imported Cabbageworms, Leafhoppers, Leafrollers, Leaf-tiers, Mexican Bean Beetles, Potato Leafhoppers, Psyllids, Stink Bugs, Thrips, Vinegar Flies, and Webworms:

Water plant thoroughly. Turn spray nozzle to the on position and hold spray nozzle about 12” from the plants. Spray lightly, covering insects on upper and lower leaf surfaces until surfaces are slightly wet. Do not over-saturate. Do not water the treated area to the point of run-off. Spray when insects or their damage first appears. Do not wait until plants are heavily infested. Spray may be applied up to and including day of harvest.