

ferti•lome® Weed-Out Plus Lawn Fertilizer 10923

ferti•lome®



Weed-Out Plus Lawn Fertilizer 25-0-4

- Slow release formulation
- Contains Trimec® herbicide

Controls Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge, and many other broadleaf weeds which are listed on this label in St. Augustinegrass, Centipedegrass, Bahiagrass, Bermudagrass, Zoysiagrass, Bluegrass, Perennial Ryegrass, Fescue spp. and Bentgrass lawns

ACTIVE INGREDIENTS

Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid	0.29%
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid.....	0.17%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	0.07%

OTHER INGREDIENTS..... 99.47%

TOTAL 100.00%

THIS PRODUCT CONTAINS:

4.80 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton or 0.24%.
 2.80 lb 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.14%.
 1.20 lb 3,6-dichloro-o-anisic acid equivalent per ton or 0.06%.


Isomer Specific by AOAC Methods.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION

KEEP OUT OF REACH OF CHILDREN
CAUTION

See back panel for complete Precautionary Statements, including First Aid, and Directions for Use, including Storage and Disposal.

NET WEIGHT 40 LBS. (18.16 KG)



ferti•lome® Weed-Out Plus Lawn Fertilizer

ferti•lome® Weed-Out Plus Lawn Fertilizer

Covers 10,000 sq. ft.
at the 4.0 lb. application rate

ferti•lome® Weed-Out
Plus Lawn Fertilizer



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wear long-sleeved shirt, long pants, shoes plus socks and chemical-resistant gloves (made of any waterproof material). User should wash hands thoroughly with soap and water before eating, drinking, chewing gum, using tobacco, or using the toilet.

FIRST AID

If In Eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.
If On Skin Or Clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a Poison Control Center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none"> Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment. You may also call toll free 1-877-800-5556 for emergency medical treatment information.

In the event of a medical or chemical emergency contact  **North America 1-800- 255-3924**

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

USE RESTRICTIONS:

- Do not exceed specified dosages for any area. Do not apply to newly seeded lawns. During application, close spreader openings when turning.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

USE PRECAUTIONS:

- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.

INSTRUCTIONS

ferti•lome® Weed-Out Plus Lawn Fertilizer controls a wide range of lawn weeds, including Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and other listed weeds commonly found in home lawns. In addition, a 25-0-4 fertilizer provides a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds or (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in 30 days. Do not use on Dichondra. Do not overapply. Under conditions of adequate moisture, high temperatures and high humidity, over-application may cause minor transitory discoloration on St. Augustinegrass. Do not use this product on "Floritam" St. Augustinegrass.

- Mow lawn to normal height 1 to 2 days before application.
- Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
- Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

HOW MUCH TO APPLY

For Kentucky Bluegrass, Fescues, Perennial Ryegrass, Bermudagrass, Zoysiagrass, St. Augustinegrass, Bahiagrass and Centipedegrass: Apply 3.2 to 4.0 pounds of product per 1,000 square feet of turfgrass. A second or follow-up application may be required for dense infestations of weeds or for adverse seasonal conditions. The second application of 3.2 to 4.0 pounds of product per 1,000 square feet should be used 30 days after the first application.

The maximum broadcast application rate is 4.0 pounds of product per 1,000 sq. ft. per application (0.24 lb 2,4-D ae, 0.42 lb MCPP-p ae and 0.10 lb dicamba ae per acre per application). The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 8.0 pounds of product per 1,000 sq. ft. (0.48 lb 2,4-D ae, 0.84 lb MCPP-p ae and 0.20 lb dicamba ae per acre).

A uniform distribution of product on the treated area is essential. Choose a calm day to ensure uniform coverage. Do not exceed the application rate for any area.

SPREADER GUIDE

These settings can be used only as a guide, they are not absolute. The speed at which the applicator walks, spreader condition and the texture of the turf will alter the rate of application. If product remains in spreader after application to recommended square foot coverage, spread remainder over same area going at right angles.

CALIBRATION INSTRUCTIONS:

- For 3.2 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 310 square feet and adjust to apply at the rate of 1 pound of product per 310 square feet.
- For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

SPREADER	SPREADER SETTINGS	
	Lbs. of product per 1,000 sq. ft.	
	3.2 lbs.	4 lbs.
Scotts®/Republic Accugreen (Drop)	3 ½	4 ¼
Scotts®/Republic Speedy Green (Broadcast)	3	3 ¾
ferti•lome® /Earthway Ev-N-Spred (Broadcast)	13	14

10923

BROADLEAF WEEDS

Annual Yellow Sweetclover	Clover, Strawberry	Mustard	Spiny Sowthistle
Aster	Clover, Sweet	Narrowleaf Plantain	Spotted Catsear
Austrian Fieldcress	Clover, White	Narrowleaf Vetch	Spurred Spurge
Bedstraw	Cockle	Nettle	Spurweed
Beggarticks	Cocklebur	Orange Hawkweed	Stinging Nettle
Betony, Florida	Common Mullein	Oriental Cocklebur	Strawberry, India Mock
Bindweed, Field	Creeping Jenny	Oxalis	Tall Nettle
Bird Vetch	Cudweed	Parsley-Piert	Tall Vervain
Bitter Wintercress	Curly Dock	Parsnip	Tansy Ragwort
Bittercress, Hairy	Daisy, English	Pearlwort	Tansy Mustard
Bitterweed	Daisy Fleabane	Pennycress	Tanweed
Black-Eyed Susan	Daisy, Oxeye	Pepperweed	Thistle
Black Medic	Dandelion	Pigweed	Trailing Crownvetch
Black Mustard	Dichondra	Pineywoods	Tumble Pigweed
Blackseed Plantain	Dogfennel	Bedstraw	Tumble Pigweed
Blessed Thistle	Dollarweed	Plains Coreopsis	Velvetleaf
Bloodflower	False Dandelion	Plantain	Venice Mallow
Milkweed	Falseflax	Poison Ivy	Virginia Buttonweed
Blue Lettuce	False Sunflower	Poison Oak	Virginia Creeper
Blue Vervain	Fiddleneck	Pokeweed	Virginia Pepperweed
Bracted Plantain	Florida Pusley	Poorjoe	Wavyleaf Bullthistle
Bristy Oxtongue	Frenchweed	Prairie Sunflower	Western Clematis
Broadleaf Dock	Galinsoga	Prickly Lettuce	Western Salsify
Broadleaf Plantain	Goldenrod	Prickly Sida	White Mustard
Broomweed	Ground Ivy	Prostrate Knotweed	Wild Aster
Buckhorn	Gumweed	Prostrate Pigweed	Wild Buckwheat
Buckhorn Plantain	Hairy Fleabane	Prostrate Spurge	Wild Carrot
Bulbous Buttercup	Hawkweed	Prostrate Vervain	Wild Four-O'clock
Bull Thistle	Healall	Puncturevine	Wild Garlic
Bullnettle	Heartleaf Drymary	Purslane, Common	Wild Geranium
Burclover	Heath Aster	Ragweed	Wild Lettuce
Burdock	Hedge Bindweed	Red Sorrel	Wild Marigold
Burning Nettle	Hedge Mustard	Redroot Pigweed	Wild Mustard
Burweed	Henbit	Redstem Filaree	Wild Onion
Buttercup	Hoary Cross	Rough Cinquefoil	Wild Parsnip
Buttonweed	Hoary Plantain	Rough Fleabane	Wild Radish
Canada Thistle	Hoary Vervain	Russian Pigweed	Wild Rape
Carolina Geranium	Horsenettle	Russian Thistle	Wild Strawberry
Catchweed	Jimsonweed	Scarlet Pimpernel	Wild Sweet Potato
Bedstraw	Knawel	Scotch Thistle	Wild Vetch
Catnip	Knotweed	Sheep Sorrel	Woodsorrel
Catsear	Kochia	Shepherdspurse	Woolly Croton
Chickweed, Common	Lambsquarters	Slender Plantain	Woolly Morningglory
Chickweed, Mouseear	Lespedeza	Smallflower Galinsoga	Woolly Plantain
Chicory	Mallow	Soga	Wormseed
Cinquefoil	Matchweed	Smooth Dock	Yarrow
Clover, Crimson	Mexicanweed	Smooth Pigweed	Yellow Rocket
Clover, Hop	Milk Vetch	Sorrel	Yellowflower
Clover, Red	Morningglory	Sowthistle	Pepperweed
	Mouseear	Spanishneedles	
	Hawkweed	Speedwell	
	Mugwort	Spiny Amaranth	
	Musk Thistle		

25-0-4

GUARANTEED ANALYSIS

Total Nitrogen (N)	25%
2% Ammoniacal Nitrogen	
23% Urea Nitrogen*	
Soluble Potash (K ₂ O).....	4%
Boron (B)	0.02%
Copper (Cu)	0.05%
0.05% Water Soluble Copper (Cu)	
Iron (Fe)	0.10%
0.10% Water Soluble Iron (Fe)	
Manganese (Mn).....	0.05%
0.05% Water Soluble Manganese (Mn)	
Zinc (Zn)	0.05%
0.05% Water Soluble Zinc (Zn)	
Derived from: Ammoniacal Sulfate, Urea, Polymer Coated Sulfur Coated Urea, Muriate of Potash, Sodium Borate, Copper Sulfate, Ferrous Sulfate, Manganese Sulfate, Zinc Sulfate. *7.5% Slowly Available Nitrogen derived from Polymer Coated Sulfur Coated Urea.	
	F370

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused pesticide in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

10923-1-0321-REG2

Manufactured by:



Voluntary Purchasing Groups, Inc.

230 FM 87
Bonham, Texas 75418
EPA Reg. No. 2217-603-7401
EPA Est. No. 7401-TX-01
Product Questions? 855-270-4776

Information regarding the contents and levels of metals in this product is available on the internet at:
<http://www.aapfco.org/metals.html>

