

Contains permethrin and diflubenzuron, the active ingredients used in Clean-Up™ II Pour-On Insecticide with IGR.



## INSECTICIDE

# GARDATIS™ POUR-ON

## CONTROL BITING AND SUCKING LICE IN CATTLE WITH ONE APPLICATION

Gardatis™ Pour-On is a ready-to-use formula that protects cattle and horses against biting and sucking lice, flies, and ticks. The pour-on contains permethrin, diflubenzuron, and petroleum distillate, a combination proved to kill adult lice and lice eggs before they hatch, providing you with effective control with just one application. Gardatis Pour-On is approved for use on lactating and non-lactating dairy cattle, beef cattle, calves, horses, and foals. An economical product, a 0.5-gallon container can treat up to 63 animals. Reduce time spent on pest infestations with the dependable power of Gardatis Pour-On.

### KEY BENEFITS

- Protects against biting and sucking lice, flies, and ticks
- Easy-to-use formulation
- A single application kills lice at multiple stages of their life cycle
- Zero-day pre-slaughter interval and milk discard time
- Prevents emergence of horn fly larvae in the manure of treated cattle

### KEY USES

- Beef Cattle
- Calves
- Dairy Cattle
- Foals
- Horses



### PRODUCT NOTES

**EPA REGISTRATION NUMBER**  
91234-329

**ACTIVE INGREDIENT**  
Permethrin 5.0%  
Diflubenzuron 3.0%

**FORMULATION**  
Suspension Concentrate

**IRAC NUMBER**  
15  
3A

**SIGNAL WORD**  
Caution

**PACKAGE SIZE**  
6 x 0.5 gal  
2 x 2.5 gal

**RESTRICTED USE**  
No

Refer to label for specific use restrictions.



Relevant. Simple. Reliable.

### APPLICATION INFORMATION

TARGET ANIMAL	INSTRUCTIONS
<b>Beef cattle</b> <b>Calves</b> <b>Dairy cattle; lactating and non-lactating</b> <b>Mature Horses</b>	<b>Pour-On:</b> Apply 3 mL per 100 lbs body weight of the animal up to maximum of 30 mL for any one animal. Pour along back line and down their face. Product may be either administered directly from graduated tip-and-pour container, graduated applicator gun, or graduated ladle/dipper. <b>Do not</b> use this method of application on foals.
<b>Foals</b> <b>Horses</b>	<b>Wipe-On:</b> Apply 3 mL per 100 lbs body weight of the animal up to a maximum of 30 mL for any one animals. Dampen an applicator mitt, cloth, or toweling with Gardatis Pour-On. Rub over their hair with special attention to the legs, shoulders, neck, and facials areas where pests appear or tend to congregate. Apply directly onto attached ticks.  <b>Ready-To-Use Spray:</b> Use Gardatis Pour-On with IGR undiluted in a mist sprayer, applying 3 mL per 100 lbs body weight of the animal. Directly spray to the legs, shoulders, neck, and facial areas where pests appear or tend to congregate. Apply directly onto attached ticks.
<b>Instructions for all methods of application:</b> Keep away from eyes and mucous membranes of animal being treated. Repeat treatment as needed, but <b>do not</b> apply more than once every two weeks.	

Gardatis Pour-On is not effective in controlling cattle grubs. Therefore, the Gardatis Pour-On can be used on cattle at any time of the year without fear of host-parasite reactions commonly associated with grub treatment products.

A 0.5-gallon (65 fl oz) container will treat up to 63 animals.  
 A 2.5-gallon container will treat up to 315 animals.

### USE RESTRICTIONS

- **Do not** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Not for use in outdoor residential misting systems.
- Not for use in metered release systems.
- Not for use in commercial animal premise automatic misting systems.
- Not for broadcast application
- **Do not** make outdoor applications during rain.
- **Do not** apply when food or feed is present.
- Remove or cover food processing equipment and food preparation surfaces or wash them before use.
- **Do not** apply by high-pressure hand wand, handheld foggers, or fog/mist generators.
- Application is prohibited directly into sewers or drains, or to any gutter-like area where drainage to sewers, storm drains, water bodies, or aquatic habitats can occur. **Do not** allow the product to enter any drain during or after application.
- When use in dairy barns or facilities: Close milk bulk tank lids to prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.

*Refer to product label for complete application and mixing instructions.*

©2023 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state. Gardatis™ Pour-On is not manufactured, or distributed by Bayer, seller of Clean-Up™ II. Clean-Up™ is a registered trademark of Bayer. Gardatis™ Pour-On is a trademark of Atticus, LLC. F1-02162023

### KEY INSECTS

Biting lice  
 Black flies  
 Cattle ticks  
 Chewing lice  
 Deer flies  
 Horn flies  
 Horse flies  
 House flies  
 Stable flies  
 Sucking lice

*(Refer to product label for complete list)*