

| VEGETABLES Cont. | | |
|------------------|--|--|
| Crop | Diseases | Insects |
| Cucumber, Squash | Alternaria Blight, Angular Leaf Spot, Cercospora Leaf Spot, Downy Mildew, Powdery Mildew, Septoria Leaf Spot | Fireworms, Flea Beetles, Fruit Flies, Harlequin Bugs, Imported Cabbageworm, Japanese Beetles, Leafhoppers, Leafrollers, Leafhoppers, Mexican Bean Beetle, Mites, Oblique-banded Leaf Rollers, Potato Leafhoppers, Psyllids, Stink Bugs, Twelvespotted Cucumber Beetle, Webworms, Whiteflies. |
| Tomato, Pepper | Alternaria Blight, Bacterial Spot, Cercospora Leaf Spot, Downy Mildew, Early Blights, Septoria Leaf Spot | |
| Onion | Angular Leaf Spot, Cercospora Leaf Spot, Downy Mildew, Septoria Leaf Spot | |

¹For European Corn Borers, make 3-4 applications or more as eggs start to hatch and repeat at 5-7 day intervals.

STORAGE & DISPOSAL

Pesticide Storage: Store and transport in an upright position. Store in a cool, secure, dry area inaccessible to children or pets.

Pesticide Disposal: If partly filled - Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Container Handling: If empty - Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

FIRST AID

| | |
|------------------------|---|
| If on skin or clothing | Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. |
| If in eyes | Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. |
| If inhaled | Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. |
| If swallowed | Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. |

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Environmental Hazards

This pesticide is toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Note: Seller warrants that this product complies with the specifications expressed in this label. To the extent consistent with applicable law, Seller makes no other warranties, and disclaims all other warranties, express or implied, including but not limited to warranties of merchantability and fitness for the intended purpose. To the extent consistent with applicable law, Seller's liability or default, breach or failure under this label shall be limited to the amount of the purchase price. To the extent consistent with applicable law, Seller shall have no liability for consequential damages.

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

TRUSTED SINCE 1926

BONIDE

Garden Dust

Ready to Use

One-Step control of insects & disease

Use on listed garden vegetables, fruits and ornamentals

ACTIVE INGREDIENTS:

| | |
|--|----------------|
| Pyrethrins | 0.03% |
| Sulfur (4-5 microns) | 25.00% |
| Copper (basic copper sulfate)* ... (CAS No. 1344-73-6) | 5.00% |
| OTHER INGREDIENTS: | 69.97% |
| TOTAL: | 100.00% |

*Metallic copper equivalent 2.8%

EPA Est. No. 4-NY-1

EPA Reg. No. 4-473

Keep Out Of Reach Of Children

CAUTION See side panels for additional Precautionary Statements and First Aid)

Packed by weight, not volume



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use Restrictions

Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift. Do not allow adults, children, or pets to enter the treated area until dusts have settled. Do not water the treated area to the point of runoff. Do not make applications during rain.

Not for use on crops grown for resale.

This product is a specially formulated insecticide and fungicide designed for convenient use as a dust.

DIRECTIONS FOR HOME GARDEN USE

Shake container to fluff powder. Cover both tops and undersides of leaves with thin uniform film of dust. Depending on foliage density, 10 oz. will cover approximately 625 sq. ft.

Start applying when plants leaf out in the Spring before insects and diseases appear, as it is easier to prevent than cure certain diseases. Try to hit as many insects as possible. Best results are obtained in early morning or evening when wind is calm. Avoid use during high temperatures on plants susceptible to sulfur injury, such as melons, squash, cucumbers and pumpkins. Repeat treatment every 7 to 10 days, as needed. Do not apply to food crops within one day of harvest.

| ORNAMENTALS | | |
|---|----------------------------------|--|
| Crop | Diseases | Insects |
| Pine (Austrian, Ponderosa) | Pine Needle Blight | Ants (excluding fire, harvester, carpenter and Pharaoh ants), Aphids, Aster Beetles, Exposed Thrips, Flea Beetles, Gypsy Moth, Japanese Beetles, Leafhoppers, Leafhoppers, Mites, Oblique-banded Leaf Rollers, Psyllids, Stink Bugs, Webworms, Whiteflies. |
| Roses and other ornamentals such as, Azaleas, Begonias, Rhododendrons, Chrysanthemums, Marigolds and Gardenias. | Black Spot, Powdery Mildew, Rust | |

For disease control, apply to point of runoff when new growth is emerging.

| FRUIT | | |
|----------------------------------|---|---|
| Crop | Diseases | Insects |
| Apple & Pear | Anthracoze, Apple Cedar Rust, Black Rot, Fire Blight, Frog Eye, Leaf Spot, Powdery Mildew, Scab, Sooty Blotch | Aphids, Caterpillars, Cherry Fruit Flies, Exposed Thrips, Fireworms, Fruit Flies, Fruit Tree Leafroller, Grape Leafhopper, Green Peach Aphid, Gypsy Moth, Japanese Beetles, Leafhoppers, Leafrollers, Leafhoppers, Mites, Psyllids, Red-necked Borers, Webworms |
| ¹ Grape | Anthracoze, Black Rot, Powdery Mildew | |
| ² Peaches & Nectarine | Brown Rot Blossom Blight, Peach Canker, Peach Leaf Curl, Shothole, Stem Blight | |
| ³ Sour Cherry | Brown Rot Blossom Blight, Leaf Spot, Shothole, Stem Blight | |

¹Certain Vinifera and French Hybro varieties of grapes may be slightly sensitive to copper sprays resulting in marginal leaf burn; tests are recommended.

²Apply to peaches only at bud swell and pink bud stages.

³Apply to cherries only at popcorn and petal fall stages.

| VEGETABLES | | |
|----------------------|--|---|
| Crop | Diseases | Insects |
| Celery | Angular Leaf Spot, Bacterial Blights, Cercospora Leaf Spot, Early Blights, Septoria Leaf Spot | Aphids, Armyworms, Asparagus Beetles, Bean Beetles, Bean Leaf Beetle, Beet Webworms, Blister Beetles, Cabbage Looper, Cabbage Worms, Caterpillars, Celery Leafhoppers, Colorado Potato Beetle, Corn Earworm, Cucumber Beetles, Diamond Back Moth, European Corn Borers ¹ , Exposed Thrips, |
| Beans, Sweet Peas | Angular Leaf Spot, Bacterial Blights, Cercospora Leaf Spot, Downy Mildew, Early Blights, Powdery Mildew, Septoria Leaf Spot | |
| Beet, Carrot, Potato | Alternaria Blight, Angular Leaf Spot, Bacterial Blights, Cercospora Leaf Spot, Downy Mildew, Early Blights, Septoria Leaf Spot | |

Bonide Products, Inc.
6301 Sulfitt Road
Oriskany, NY 13424
©Bonide Products, Inc.
All Rights Reserved