



# Corral<sup>®</sup> 2.68G

Preemergent control of goosegrass, crabgrass, *Poa annua* and other annual grassy weeds in turfgrass and landscape ornamentals. Preemergent control of broadleaf weeds.

**This specimen label is for information only.**  
Everris makes no representations as to the accuracy of this label. It is the responsibility of the user to read and follow the label attached to the pesticide product container.

**Net Weight: 50 lbs (22.68 kg)**  
**Stock # E95750**

**ACTIVE INGREDIENT:**

Pendimethalin† . . . . . 2.68%

**OTHER INGREDIENTS:** . . . . . 97.32%

Total 100.00%

† CAS No. 40487-42-1

**EPA Reg. No. 58185-179**

**EPA Est. 82757-OH-1<sup>MF</sup>, EPA Est. 8378-IN-1<sup>K</sup>**

Superscript used is the first letter of run code on end of container.

## KEEP OUT OF REACH OF CHILDREN CAUTION

### FIRST AID

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling poison control center or doctor, or going for treatment. You may also call 1-888-241-6409 for emergency medical assistance.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION.** Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wear long-sleeved shirt and long pants, socks and shoes, and chemical resistant gloves (made of any waterproof material).

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical resistant category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Applicators must wear chemical resistant gloves such as butyl rubber > 14 mils and natural rubber > 14 mils.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**If you have questions regarding the use of this product, call TOLL FREE 1-800-492-8255**

Distributed by Everris NA Inc., PO Box 3310 • Dublin, OH 43016

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. In case of spills, either collect for use or dispose of properly.

### RECOMMENDED FOR USE BY PROFESSIONAL TURFGRASS OR NURSERY MANAGERS

#### Controls these grassy weeds as they sprout:

Barnyardgrass	Itchweed	Sandbur, Field
Crabgrass	Johnsongrass	Signalgrass
Crowfootgrass	(from seed)	Sprangletop
Fall Panicum	Junglerice	Witchgrass
Foxtail	Lovegrass	Woolly Cupgrass
Goosegrass	<i>Poa annua</i>	

#### Controls these broadleaf (dicot) weeds as they sprout:

Carpetweed	Knotweed	Purslane
Chickweed	Kochia	Pusley, Florida
Cudweed	Lambsquarters	Rocket, London
Evening Primrose	Oxalis	Shepherdspurse
Fiddleneck	Pearlwort	Smartweed
Filaree	Pigweed	Velvetleaf
Henbit	Prostrate Spurge	
Hop Clover	Puncturevine	

## DIRECTIONS FOR USE

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH ITS LABELING.**

### ORNAMENTALS

For use on container and field grown ornamentals.

#### Application Restrictions

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### FOR USE AROUND AND OVER THE TOP OF ALL ESTABLISHED LANDSCAPE, FIELD GROWN AND CONTAINER ORNAMENTALS LISTED BELOW.

Abelia	Betula	Cortaderia	Ilex	Myrtus	Platanus	Syringa
Abies	Butia	Cotoneaster	Juniperus	Nandina	Potentilla	Taxus
Acer	Buxus	Cryptomeria	Kalmia	Nerium	Prunus	Thuja
Achillea	Callistemon	Euonymus	Lagerstroemia	Ophiopogon	Pseudosuga	Tilia
Acorus	Carissa	Forsythia	Ligustrum	Oxydendron	Pyracantha	Trachycarpus
Agapanthus	Ceanothus	Franklinia	Liriope	Pachysandra	Pyrus	Tsuga
Ajuga	Chamaecyparis	Fraxinus	Lobelia	Penstemon	Quercus	Washingtonia
Aloysia	Chamaerops	Gardenia	Magnolia	Photinia	Raphiolepis	Wisteria
Arbutus	Convolvulus	Gleditsia	Malus	Picea	Rhododendron	Yucca
Arundo	Cordyline	Hedera	Miscanthus	Pieris	Solanum	
Asparagus Fern	Cornus	Hemerocallis	Moraea	Pinus	Spirea	
Berberis	Correa	Hosta	Myrica	Pittosporum	Stokesia	

#### GROUND COVERS

Aptenia	Baccharis	Fragaria	Hedera helix	Myoporum	Pelargonium	Verbena
Arctotheca	Carpobrotus	Gazania	Hypericum	Osteospermum	Rosemary	Vinca

#### PERENNIALS

Aster	Asclepias	Lythrum
-------	-----------	---------

#### ORNAMENTAL GRASSES

Ammophila	Deschampsia	Festuca	Pennisetum	Phalaris
-----------	-------------	---------	------------	----------

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

Personal protective equipment required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical resistant gloves made of any waterproof material, shoes plus socks, and protective eyewear.

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.



## DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

### TURF

#### Application Restrictions

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply more than 75 lb of product/acre (2.0 lb ai/acre) to residential lawns, recreational areas (turf) and sod farms.

For use in residential lawns, recreational areas, sod farms, commercial areas and golf courses.

For use on established bermudagrass, St. Augustinegrass, centipede-grass, zoysiagrass, bahiagrass, and tall fescue.

For goosegrass, crabgrass, foxtail, barnyardgrass, fall panicum, *Poa annua*, cudweed, hop clover, oxalis, evening primrose, chickweed, purslane, knotweed, henbit and prostrate spurge, apply in the spring or fall before they sprout.

For annual grass and dicot weeds, apply before they sprout or after established weeds are removed.

For best results, water after application ( $\frac{1}{4}$  to  $\frac{1}{2}$  inch) to remove granules from the foliage and to move the product to the soil surface. Do not destroy the chemical barrier after application by raking or aerifying.

Do not contaminate feed or foodstuffs. Keep children and pets off treated areas until material is washed into soil and foliage has dried.

To prevent *Poa annua*, henbit, and fall germinating chickweed, apply product at the suggested rate in late summer-early fall before germination. Repeat treatment yearly for continued control but not sooner than the following spring.

On newly seeded or sodded turfgrass area, wait until after the fourth mowing before application. For established turfgrass areas where reseeding is necessary, delay the product application until after seeding and after the new grass seedlings have been mowed 4 times. This may be up to 3 months from the date of seeding.

On newly sprigged areas, wait 5 months before application.

### BERMUDAGRASS PRECAUTIONS

When applied to dormant bermudagrass, greenup may be delayed 2 to 3 weeks in the spring. The delay in greenup may be more extensive on low-cut bermudagrass and on sandy soils.

Application of this product to bermudagrass overseeded with cool season turfgrass species may injure and thin the overseeding.

Application of this product to bermudagrass greens that are overseeded with cool season turfgrass species may inhibit the transition of bermudagrass, especially when the green is verticuted, aerified, etc. These practices in conjunction with an application of this product may remove the overseeding faster than the bermudagrass can fill in, resulting in thin greens.

## OTHER PRECAUTIONS – TURF

Where tall fescue is grown as a mixture with cool season grasses, this product should be used only if tall fescue is the desired turf species.

Where turf thinning or damage has occurred due to winter injury, wait until greenup, fill-in, and good rooting before making an application.

This product should not be used on winter overseedings unless enhanced reduction of the cool season grass species is desired.

Not for use on dichondra.

This product cannot be used on newly seeded or sprigged turfgrass areas or where turfgrass has been severely thinned due to winter damage, insects, disease, scalping, etc. Wait until fill-in and rooting of stolons/rhizomes is complete and desired turf density is achieved.

For established turfgrass areas where reseeding is necessary, delay product application until after new grass seedlings have been mowed 4 times.

Residual effectiveness requires a delay of four growing months after treatment before reseeding or winter overseeding and five growing months before sprigging.

Product does not control germinated or established weeds.

Allow at least a four month span between any two applications except when a repeat treatment is necessary after 6-8 weeks. Do not apply more than a cumulative total of 113 lbs. of product/acre (3.0 lbs. active ingredient/acre) per year on bentgrass and *Poa annua*; 226 lbs. of product/acre (6.0 lbs. active ingredient/acre) per year on cool season turfgrasses; and 339 lbs. of product/acre (9.0 lbs. active ingredient/acre) per year on warm season turfgrasses.

This product may cause discoloration and thinning where overapplications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature – below 40° F, excessive soil moisture, scalping, etc.).

If metallic surfaces (golf course equipment, automobiles, etc.) are present at the application site, do not use this product under windy conditions (5-10 MPH) due to its staining potential.

## APPLICATION RATES

TURFGRASS SPECIES	WEEDS	RATES OF CORRAL® 2.68G	COMMENTS
Warm Season Grasses Bahia grass Bermudagrass Centipedegrass St. Augustine grass Tall Fescue Zoysiagrass	barnyardgrass chickweed crabgrass fall panicum foxtail knotweed <i>Poa annua</i> prostrate spurge purslane	75 lbs./acre prior to weed germination	Apply a repeat application of 75 lbs./acre after 6-8 weeks if necessary
	goosegrass	75 lbs./acre prior to weed germination	Apply a repeat application of 75 lbs./acre to extend control of goosegrass
	cudweed evening primrose henbit hop clover	75 lbs./acre prior to weed germination	
Bermudagrass Greens	barnyardgrass chickweed crabgrass fall panicum foxtail knotweed <i>Poa annua</i> prostrate spurge purslane	75 lbs./acre prior to weed germination	Apply a repeat application of 75 lbs./acre after 6-8 weeks if necessary
	cudweed evening primrose goosegrass henbit hop clover	75 lbs./acre prior to weed germination	Apply a repeat application of 57 lbs./acre after 6-8 weeks
Cool Season Grasses Fine Fescue Kentucky Bluegrass Perennial Ryegrass Tall Fescue	barnyardgrass chickweed crabgrass fall panicum foxtail knotweed oxalis <i>Poa annua</i> prostrate spurge purslane	57-75 lbs./acre as the initial application prior to weed germination	Apply a repeat application of 57 lbs./acre after 6-8 weeks for acceptable control at the lower rate
	goosegrass	57-75 lbs./acre in the spring prior to weed germination.	Apply a repeat application after 6-8 weeks at 57 lbs./acre
Bentgrass and Established <i>Poa annua</i> (Fairway cut. Not for use on greens or tees)	barnyardgrass chickweed crabgrass fall panicum foxtail knotweed oxalis <i>Poa annua</i> prostrate spurge purslane	57 lbs./acre in the spring prior to weed germination.	Apply a repeat application of 57 lbs./acre after 6-8 weeks where heavy weed infestations are expected
	goosegrass	57 lbs./acre in the spring prior to weed germination	Apply a repeat application of 57 lbs./acre after 6-8 weeks

## SUGGESTED SPREADER SETTINGS FOR TURF

To provide proper distribution calibrate spreader before application.

Spreader	Ground or PTO Speed	Width of Coverage	RECOMMENDED RATE RANGE (lbs. of product/acre)	
			57 lbs. product/A	75 lbs. product/A
Scotts Drop®	3 mph	overlap wheels	2½	2½
Gandy (30")	3 mph	overlap wheels	17	19
Scotts Rotary R-7, R-7X	3 mph	8 ft.	B	B½
Scotts Rotary R-8A, R-8 (cone 9)	3 mph	9 ft.	H	H½
Lely Models (see below)	4.5 mph	18 ft.	2½ III	3 III
Vicon (Model 02) (Model 03)	4.5 mph 4.5 mph	18 ft. 10 ft.	10 7	12 9

Lely models WTR, WFR, HR, and 1250. PTO at 450 rpm.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a clean, dry place. Reseal opened bag by folding top down and securing.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

## DISCLAIMER AND LIMITATION OF LIABILITY

### IMPORTANT NOTICE FROM EVERRIS NA INC. ("Everris").

#### PLEASE READ BEFORE USE.

To the extent consistent with applicable law, user or buyer accepts the conditions, disclaimer of warranties and limitations of liability. Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once for full refund.

**CONDITIONS:** The directions for use of this product are believed to be adequate and the user or buyer must always follow the label directions carefully and exercise judgment and caution when using this product under their growing conditions. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness, unsatisfactory or substandard results or other unintended consequences may result because of such factors as weather conditions, presence or absence of other materials, or the manner of use or application, all of which are beyond the control of Everris. All such risks shall be assumed by the user or buyer.

**WARRANTY:** This product corresponds to all claims and descriptions set forth on the label and is reasonably fit for the purposes set forth in the directions for use on the label when used in accordance with those directions. This warranty is subject to the conditions and limitations stated herein. Everris recognizes that the rights and remedies of the user or buyer are subject to the provisions of the applicable law, but makes no other warranties or representations, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Everris is authorized to make any warranties beyond those contained herein or to modify the warranties contained therein. Subject to the user's or buyer's rights and remedies under the applicable law, Everris disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** Subject to the user's or buyer's rights and remedies under the applicable law, the exclusive remedy of the user or buyer and the liability of Everris or its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid by the user or buyer for the quantity of this product involved or, at Everris' election, the replacement of the product. To the extent consistent with applicable law, Everris must have prompt notice of any claim so that a timely investigation of buyer's or user's claim can be made. Buyer and all users shall promptly notify Everris of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

Distributed by:  
Everris NA Inc.  
PO Box 3310  
Dublin, OH 43016  
1-800-492-8255

Product of USA