

LIQUI-VICT 2X CANINE ANTHELMINTIC- pyrantel pamoate suspension
Happy Jack Inc

Happy Jack®
LIQUI-VICT 2X™

INDICATIONS FOR USE

To prevent reinfestation of (Toxocara canis) in puppies and adult dogs and in lactating bitches after whelping.

For the removal of large roundworms (Toxocara canis and Toxascaris leonina) and hookworms (Ancylostoma caninum and Uncinaria stenocephala) in dogs and puppies.

Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

WARNING: Keep out of Reach of Children

Read Accompanying Professional Information

LOT
EXP

RECOMMENDED STORAGE:
STORE BELOW 86°F (30°C)

Manufactured for:
HAPPY JACK, INC.
Snow Hill, NC 28580
www.happyjackinc.com

Take Time
(eye symbol)
Observe Label
Directions

IMPORTANT: SHAKE WELL BEFORE USING

DIRECTIONS FOR USE

For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent T.canis reinfestation. Administer one full teaspoon (5 mL) for each 10 pounds of body weight.

For the removal of large roundworms (ascarids) and hookworms. Administer one full teaspoon (5 mL) for each 10 pounds of body weight. It is not necessary to withhold food prior to treatment. If medication is to be dispensed, client can be advised that dogs usually find this wormer very palatable and will lick the dose from the bowl willingly. If there is a reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. The presence of large roundworms (ascarids) and hookworms should be confirmed by lab fecal examination.

It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2 to 4 weeks after first treatment.

PRINCIPAL DISPLAY PANEL - 60 mL Bottle Label

Happy Jack

LIQUI-VICT 2X

(pyrantel pamoate)

Canine Anthelmintic Suspension

Active Ingredient:

4.54 mg of pyrantel base

as pyrantel pamoate per ml.

FOR ANIMAL USE ONLY

ANADA 200-007 Approved by FDA

NET CONTENTS – 2 FL. OZ. (60ML)

INDICATIONS FOR USE
To prevent reinfection of (*Toxocara canis*) in puppies and adult dogs and in lactating bitches after whelping.

For the removal of large roundworms. (*Toxocara canis* and *Toxascaris leonina*) and hookworms. (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies.

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

WARNING: Keep Out Of Reach Of Children
Read Accompanying Professional Information

LOT EXP [REDACTED]

RECOMMENDED STORAGE:
STORE BELOW 30°C (86°F)

Manufactured for:
HAPPY JACK, INC.
Snow Hill, NC 28580
www.happyjackinc.com

Take Time
Observe Label
Directions



LIQUI-VICT 2X™
(pyrantel pamoate)
Canine Anthelmintic
Suspension

Active Ingredient:
4.54 mg of pyrantel base
as pyrantel pamoate per ml.

FOR ANIMAL USE ONLY
ANADA 200-007 Approved by FDA
NET CONTENTS - 2 Fl. Oz. (60ML)

IMPORTANT: SHAKE WELL BEFORE USING

DIRECTIONS FOR USE: For maximum control and prevention of reinfection, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent T.canis reinfection. Administer one full teaspoonful (5 mL) for each 10 lb of body weight.

For the removal of large roundworms (ascarids) and hookworms. Administer one full teaspoonful (5 mL) for each 10 lb of body weight. It is not necessary to withhold food prior to treatment. If medication is to be dispensed, client can be advised that dogs usually find this wormer very palatable and will lick the dose from the bowl willingly. If there is a reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. The presence of large roundworms (ascarids) and hookworms should be confirmed by lab fecal examination.

It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2 to 4 weeks after first treatment.

LIQUI-VICT 2X CANINE ANTHELMINTIC

pyrantel pamoate suspension

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:23851-312
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
pyrantel pamoate (UNII: 81BK194Z5M) (pyrantel - UNII:4QIH0N49E7)	pyrantel	4.54 mg in 1 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:23851-312-02	60 mL in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200007	11/01/1996	

Labeler - Happy Jack Inc (003189354)

Establishment

Name	Address	ID/FEI	Business Operations
ABS Corporation		048022727	MANUFACTURE

Establishment

Name	Address	ID/FEI	Business Operations
COSMA Spa		428655732	API MANUFACTURE

Revised: 7/2013

Happy Jack Inc