



HEXCEL™

EW



Contains Hexythiazox, the active ingredient used in Hexygon® IQ.

EMULSION FOR COMMERCIAL USE ONLY

ACTIVE INGREDIENT: Hexythiazox	(% by weight)
<i>trans</i> -5-(4-Chlorophenyl)-N-cyclohexyl-4-methyl-2-oxothiazolidine-3-carboxamide.....	11.93%
OTHER INGREDIENTS:	88.07%
TOTAL	100.00%

Contains 1.0 lb. of Hexythiazox per gallon.

Contains Petroleum Distillate

EPA Reg. No. 91234-325

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See below for additional Precautionary Statements.

FIRST AID	
If swallowed:	<ul style="list-style-type: none"> ▪ Call a poison control center or doctor immediately for treatment advice. ▪ Do not induce vomiting unless told to do so by the poison control center or doctor. ▪ Do not give any liquid to the person. ▪ Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> ▪ Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. ▪ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ▪ Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15 - 20 minutes. ▪ Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> ▪ Move person to fresh air. ▪ If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. ▪ Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
Contains petroleum distillate. Vomiting may cause aspiration pneumonia.	

**For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident,
Call CHEMTREC Day or Night Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)**

Hexcel™ EW is not manufactured, or distributed by Gowan Company, seller of Hexygon® IQ Miticide.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate or viton \geq 14 mils
- Shoes plus socks

Remove and wash contaminated clothing before reuse. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. **Do not** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **Do not** contaminate water when cleaning equipment or when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame. **Do not** mix or allow coming in contact with oxidizing agent. Hazardous chemical reaction may occur.

PRODUCT INFORMATION

Hexcel EW is for use as a foliar application in labeled crops. **Hexcel EW** provides effective control of Tetranychid spider mite species. It is an emulsion to be mixed with water and applied as a spray.

Hexcel EW controls mites through activity on eggs and immature stages. Control is achieved from either direct contact with the spray or from contact with treated plant surfaces. Through its ovicidal activity, **Hexcel EW** controls newly deposited mite eggs and eggs which are laid after application. **Hexcel EW** is also highly effective in controlling immature motile stages of target mite species that are sprayed or move onto treated surfaces. Adult mites are not directly affected. However, eggs produced by females in contact with treated surfaces will be rendered nonviable. **Hexcel EW** provides residual control of pest mite species. The degree and duration of control is dependent on the rate used, growth stage of the mite, species of mite, and climatic conditions under which the material is applied.

Apply **Hexcel EW** prior to adult mite build up. **Hexcel EW** will not control adult spider mites. Use higher rates on moderate-to-high mite infestations or for larger plants with a dense canopy. If adult mites are present in medium-to-high populations, better results may be obtained by using **Hexcel EW** in combination with a registered contact adulticide. The use of less than label rates is not recommended with **Hexcel EW**.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS

- Apply this product only as specified on the EPA approved label and any supplemental labeling.
- **Do not** graze or feed livestock on cover crops growing in treated areas.
- **Do not** make more than one application of **Hexcel EW** or any other Hexythiazox product per year.
- **Do not** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- **Do not** plant rotational crops other than those on this label within 120 days of this application.
- For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate or viton \geq 14 mils
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants in farms, forests, nurseries, or greenhouses.

REENTRY: Keep children, pets and unprotected persons out of treated areas until sprays have dried.

MANAGING MITE RESISTANCE

For resistance-management, **Hexcel EW** contains a Group 10A acaricide. Any mite population may contain individuals naturally resistant to **Hexcel EW** and other Group 10A acaricides. The resistant individuals may dominate the mite population if this group of acaricides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed. To reduce the potential for developing mite resistance, rotate to an acaricide with a different mode of action. Monitor treated pest populations for resistance development. Read product label before applying any acaricide and follow label directions.

To delay mite resistance, take the following steps:

- Rotate the use of **Hexcel EW** or other Group 10A acaricides within a growing season, or among growing seasons, with different groups that control the same pests. Avoid application of more than the maximum seasonal use rate or the total number of consecutive sprays of **Hexcel EW** per season.
- Use tank mixtures with acaricides from a different group that are equally effective on the target pest when such use is permitted. **Do not** rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - o Individual acaricides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - o Mixtures with components having the same IRAC mode of action classification are not recommended for mite resistance management.
 - o When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - o Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
 - o The mite resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual acaricidal activity. Mixtures of acaricides with unequal periods of residual mite activity may offer a mite resistance management benefit only for the period where both acaricides are active.
- Adopt an integrated pest management program for acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Atticus, LLC at 984-465-4800.

MIXING DIRECTIONS

- Turn on spray tank agitation. Pour the specified amount of this material on the surface of water in a nearly filled spray tank.
- Add balance of water to fill the tank.
- Keep agitator running during filling and spraying operations. **Do not** allow mixture to stand.

Tank Mix Compatibility: **Hexcel EW** is compatible with most crop protection chemicals. However, all possible combinations have not been evaluated. Before full-scale mixing of **Hexcel EW** with other products, test mix small proportionate quantities of each to ensure compatibility. **Do not** mix or allow coming in contact with oxidizing agents. Hazardous Chemical reaction may occur.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Phytotoxicity: As is common with most emulsion formulations, adverse effects such as spotting or discoloration of the treated surfaces can occur. Some conditions known to contribute to phytotoxicity include, but are not limited to, high temperatures, poor spray drying conditions, excessive spray deposit or run-off, certain spray mixtures, stage of crop development or tank mixes with other pesticides.

APPLICATION INFORMATION

Use sufficient spray volume to obtain thorough, uniform coverage of all plant surfaces. Apply using ground equipment, chemigation or air unless otherwise specified under the crop comments or on supplemental labeling supplied by Atticus, LLC. This product is for use in commercial plantings. It is also intended for use in home plantings and ornamental landscaping by licensed or certified pesticide applicators.

Important: **Hexcel EW** is a 1 pound active ingredient per gallon emulsion to be diluted with water for application in commercial plantings, home plantings and ornamental landscaping.

For best results, apply **Hexcel EW** at first sign of mites before adult mite build up. **Hexcel EW** will not control adult spider mites. Use higher rates on moderate to high mite infestations, large trees with dense foliage, or for extended residual control. The lowest rate of **Hexcel EW** may be used in conjunction with other IPM strategies, or for shorter residual control later in the season. The use of less than label rates is not recommended since it will result in poor performance and contribute to resistance development. If adult mites are present in medium to high populations, better results may be obtained using **Hexcel EW** in combination with a registered contact adulticide. **Hexcel EW** may be tank mixed with horticultural grade oil. Refer to all precautions regarding phytotoxicity on the oil label.

Do not make more than one application of hexythiazox per year.

Chemigation Systems: **Hexcel EW** may be applied through irrigation systems (chemigation) to any crop on the **Hexcel EW** label unless otherwise specified under the crop comments or on supplemental labeling supplied by Atticus, LLC. **Do not** allow chemigation to run off field.

Types of Irrigation Systems: Apply **Hexcel EW** only through sprinkler, including center pivot, lateral move, Low Energy Precision Applications (LEPA), end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. **Do not** apply **Hexcel EW** through any other type of irrigation system.

USE DIRECTIONS FOR ALL RECOMMENDED TYPES OF IRRIGATION SYSTEMS

Uniform Water Distribution and System Calibration: The irrigation system must provide uniform distribution of treated water. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. For best results apply at 100% input/travel speed, for center pivots or 0.10 inch (2,716 gallons) up to 0.15 inch (4,073 gallons) of water/A, for other systems. Higher labeled rates of **Hexcel EW** may be necessary for chemigation applications.

Chemigation Monitoring: A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Drift: **Do not** apply when wind speed favors drift beyond the area intended for treatment.

Required System Safety Devices: The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must

also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Using Water from Public Water Systems: Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regular serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually

shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Cleaning the Chemical Injection System: In order to accurately apply pesticides, the chemical injection system must be kept clean; free of chemical or fertilizer residues and sediments. Refer to your owner's manual or ask your equipment supplier for the cleaning procedure for your injection system.

Flushing the Irrigation System: At the end of the application period, allow time for all lines to flush the pesticide through all nozzles or emitters before turning off irrigation water. To ensure the lines are flushed and free of pesticides, a dye indicator may be injected into the lines to mark the end of the application period.

Equipment Area Contamination Prevention

It is recommended that nozzles in the immediate area of control panels, chemical supply tanks, pumps and system safety devices be plugged to prevent chemical contamination of these areas.

Center-Pivot and Automatic-Move Linear Systems: Inject the specified dosage per acre continuously for one complete revolution or move of the system. **DO NOT USE END GUNS.** The system should be run at maximum speed.

Solid Set and Manually Controlled Linear Systems: Injection should be during the last 30 to 60 minutes of regular irrigation period or as a separate 30 to 60 minute application not associated with a regular irrigation. **DO NOT USE END GUNS.**

APPLICATION DIRECTIONS

CROP	PEST	RATE	COMMENTS
CHRISTMAS TREE, CHRISTMAS TREE PLANTATIONS	Pacific Spider Mite, Spruce Spider Mite, Two-spotted Spider Mite	12 - 24 fl. oz./acre OR 4 - 8 fl. oz./100 gal (0.094 - 0.19 lb. a.i./A)	Ground Application: Apply at specified rates. Air Application: Apply in 2 - 10 gallons finished spray per acre. For best coverage, apply in two opposing passes of 5 gallons each. Chemigation: See details under Chemigation Systems.
<ul style="list-style-type: none"> Do not make more than one application per calendar year. 			
GREENHOUSE TOMATOES (Including tomatoes grown for transplant)	Tetranychid Mites (such as European Red Mite and Two-spotted Spider Mite)	12 - 24 fl. oz./acre OR 4 - 8 fl. oz./100 gal OR 0.28 - 0.55 oz./1000 sq. ft. (0.094 - 0.19 lb. a.i./A)	Use sufficient volume of water to obtain complete coverage but not less than 20 gallons per acre.
<ul style="list-style-type: none"> Do not make more than one application per calendar year. Do not apply more than 24 ounces (0.19 lb. a.i.) of formulated product per acre per calendar year. Pre-Harvest Interval (PHI): 1 day. 			

(continued)

APPLICATION DIRECTIONS (continued)

CROP	PEST	RATE	COMMENTS
<p>ORNAMENTAL PLANTS AND NONBEARING FRUIT TREES AND VINES Growing in Nurseries, Greenhouses, Shadehouses and Christmas Tree Plantations</p> <p>ESTABLISHED ORNAMENTAL LANDSCAPE PLANTINGS Including those in: Interiorscapes, Residences, Public Areas, Commercial Areas, Institutional Areas, Industrial Areas, Rights of way and other easements, and Recreational sites such as campgrounds, golf courses, parks and athletic fields</p> <p>*ORNAMENTAL LAWNS AND TURF (Including residential) *Not Registered for Use in California</p> <p>*NON-RESIDENTIAL TURFGRASS (Including turf grown for sod, turf grown to produce seed, or turf for research purposes) *Not Registered for Use in California</p>	<p>Arborvitae Spider Mite, European Red Mite, Honeylocust Spider Mite, Pacific Spider Mite, Southern Red Mite, Spruce Spider Mite, Strawberry Spider Mite, Two-spotted Spider Mite, Willamette Mite, Banks Grass Mite, Clover Mite, Brown Wheat Mite</p>	<p>12 - 24 fl. oz./acre OR 4 - 8 fl. oz./100 gal OR 0.28 - 0.55 oz./1000 sq. ft. (0.094 - 0.19 lb. a.i./A)</p>	<p>When using high-volume dilute sprays, do not exceed the maximum per acre rate or apply less than the minimum labeled rate. For concentrate equipment, use the specified rate per acre.</p> <p>Apply to lawns and turf at the first signs of egg deposition.</p>
<p>RESIDENTIAL POME FRUIT GROUP 11-10 (such as apple, pear)</p> <p>RESIDENTIAL STONE FRUIT GROUP (such as peach, plum, nectarine)</p> <p>RESIDENTIAL NUT TREES GROUP AND PISTACHIOS (such as almond, pistachio)</p>	<p>European Red Mite, Pacific Spider Mite, Strawberry Spider Mite, Two-spotted Spider Mite, Willamette Mite, Pecan Leaf Scorch Mite, Citrus Red Mite</p>	<p>12 - 24 fl. oz./acre OR 4 - 8 fl. oz./100 gal (0.094 - 0.19 lb. a.i./A)</p>	<p>When using high-volume dilute sprays, do not exceed the maximum per acre rate or apply less than the minimum labeled rate. For concentrate equipment, use the specified rate per acre.</p>
<ul style="list-style-type: none"> ▪ Pre-Harvest Interval (PHI): <ul style="list-style-type: none"> o Residential Pome Fruit: 28 days. o Residential Stone Fruit: 7 days. o Residential Nut Trees and Pistachio: 7 days. 			
<p>ORCHIDS</p>	<p>Arborvitae Spider Mite, European Red Mite, Honeylocust Spider Mite, Pacific Spider Mite, Southern Red Mite, Spruce Spider Mite, Strawberry Spider Mite, Two-spotted Spider Mite, Willamette Mite</p>	<p>12 - 24 fl. oz./acre OR 4 - 8 fl. oz./100 gal OR 0.28 - 0.55 fl. oz./1000 sq. ft. (0.094 - 0.19 lb. a.i./A)</p>	<p>When using high-volume dilute sprays, do not exceed the maximum per acre rate. For concentrate equipment, use the specified rate per acre.</p> <p>Note: If used on orchids, apply prior to bloom formation.</p>
<p>PEPPER/EGGPLANT SUBGROUP 8-10B African eggplant; bell pepper; eggplant; martynia; nonbell pepper; okra; pea eggplant; pepino; roselle; scarlet eggplant; cultivars, varieties, and/or hybrids of these</p>	<p>Tetranychid Mites (such as Two-spotted Spider Mite), Pacific Spider Mite, Strawberry Spider Mite</p>	<p>12 - 24 fl. oz./acre OR 4 - 8 fl. oz./100 gal OR 0.28 - 0.55 fl. oz./1000 sq. ft. (0.094 - 0.19 lb. a.i./A)</p>	<p>Ground Application: Apply a minimum of 20 gallons finished spray per acre.</p> <p>Air Application: Apply a minimum of 20 gallons finished spray per acre.</p> <p>Chemigation: See details under Chemigation Systems.</p>
<ul style="list-style-type: none"> ▪ Do not make more than one application per calendar year. ▪ Do not apply more than 24 ounces of formulated product per acre per calendar year. ▪ Pre-Harvest Interval (PHI): 1 day. 			

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Nonrefillable container. Do not reuse or refill this container. If empty: Offer for recycling if available or discard in a sanitary landfill. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

For plastic containers ≤ 5 gallons: Nonrefillable Container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

For plastic containers > 5 gallons: Nonrefillable container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Recap and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED, OR DRINKING WATER!

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

Hexcel is a trademark of Atticus, LLC.

Hexygon® is a registered trademark of Gowan Company.

20230322a