

Hi-Yield® Garden & Farm Insect Control

CONTROLS:

Ants
Aphids
Beetles
Borers
Crickets
Fire Ants
Flies
Grasshoppers
Mosquitoes
Ticks
Plus over
150 more
insects
listed
on label.



CONTROLS INSECTS ON:

Dogs
Flowers
Fruit Trees
Indoor Plants
Lawn Areas
Livestock
Nut Trees
Outdoor Pests
Poultry
Roses
Trees and
Shrubs
Vegetables



Do Not Use On Cats

- Controls Over 150 Insects, Outdoor Use - Residential Homeowner
- FOR TERMITES: OUTSIDE, POST-CONSTRUCTION USE ONLY • Plus Borer And Carpenter Ant Control

ACTIVE INGREDIENT:

* Permethrin..... 10.0%

OTHER INGREDIENTS**..... 90.0%

TOTAL..... 100.0%

*(3-phenoxyphenyl)methyl
(+)-3-(2,2-dichloroethenyl)-2,2-dimethylcyclo-
propanecarboxylate,

** Contains petroleum distillates.

KEEP OUT OF REACH OF CHILDREN

WARNING

See Back Label Panel and Label Booklet
for Additional Precautionary
Statements and Directions For Use

NET CONTENTS ONE PINT (16 FL. OZS. / 473 ML)

Hi-Yield® Garden & Farm Insect Control

PEEL FROM CORNER OF BOOKLET

FIRST AID

If On Skin Or Clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If In Eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
If Inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
If Swallowed	<ul style="list-style-type: none">• Immediately call a poison control center or doctor.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give any liquid to the person.• Do not give anything by mouth to an unconscious person.

Note to Physician and Veterinarian: This product contains petroleum distillates. Vomiting may cause aspiration pneumonia.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

In the event of a medical or chemical emergency, contact ChemTel 
North America 1-800-255-3924 or Worldwide Intl. + 01- 813-248-0585

Manufactured By:



230 FM 87

BONHAM, TEXAS 75418

EPA Reg. No. 7401-456

EPA Est. No. 7401-TX-01

Product Questions?

855-270-4776



Manufactured By:



230 FM 87 • BONHAM, TX 75418

EPA Reg. No. 7401-456

EPA Est. No. 7401-TX-01

Visit Us At: www.fertilome.com

Product Questions? 855-270-4776

33005-1019-CL



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes skin irritation. Harmful if inhaled. Causes moderate eye irritation. Do not get on skin or on clothing. Avoid breathing spray mist. Avoid contact with eyes or clothing. Wear long sleeved shirt and long pants, socks, chemical resistant footwear and gloves. Wash hand thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before re-use.

DO NOT USE ON CATS: May be toxic or potentially fatal if applied to or ingested by cats. Accidental application to cats and/or grooming a recently treated dog may result in tremors and/or uncoordinated muscle movements. If this occurs, immediate veterinary care should be provided.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively foraging the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

Do Not use or store near heat or open flame. Do Not use this product in or around electrical equipment due to the possibility of shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

1. Treatment to soil or vegetation around structures;
2. Applications to lawns, turf, and other vegetation;
3. Applications to building foundations, up to a maximum height of 3 feet.

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

Do not water the treated area to the point of run-off.

Do not make applications during rain.

Not for use in outdoor residential misting systems.

USE ONLY ON DOGS OR LIVESTOCK. DO NOT APPLY TO CATS.

Except when applying to dogs or livestock, do not allow adults, children or pets to enter treated areas until sprays have dried.

Remove pets, birds and cover fish aquaria before spraying.

Except when applying to dogs or livestock, do not apply this product that will contact adults, children or pets, either directly or through drift.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

READ ENTIRE LABEL BEFORE EACH USE

HOW TO MIX AND APPLY

Mix the labeled amount of Hi-Yield® Garden & Farm Insect Control as directed and apply in accordance with specific directions for use. Always calculate the area to be treated by measuring the length and the width then multiply together to determine the square footage of the area to be treated, where appropriate. Apply Hi-Yield® Garden & Farm Insect Control as a coarse spray with a pump-up or other type of hand-held sprayer. When mixing, first fill the sprayer with half of the required amount of water, then measure and add the appropriate amount of Hi-Yield® Garden & Farm Insect Control. Complete filling the sprayer with the remaining amount of required water. Shake the sprayer to mix the product and water. Thorough spray coverage is necessary, but avoid spray runoff or dripping.

For large area coverage such as on lawns, this product may also be used through a hose-end type sprayer such as the ferti•lome® Air-O-Matic Hose-End Sprayer. Always follow the directions provided with the hose-end sprayer type for proper dilution and application of Hi-Yield® Garden & Farm Insect Control.

HOW TO MIX AND APPLY (Continued)

Do not use food utensils, such as measuring spoons or cups, for food purposes after use in measuring pesticides.

MEASUREMENT CONVERSION CHART

1 teaspoon = 1/6 fl.oz.; 1 ounce = 6 teaspoons;
3 teaspoons = 1 tablespoon;
2 tablespoons = 1 ounce
Tbs = Tablespoon; tsp = Teaspoon

ORNAMENTAL HOUSE PLANTS

Contact and residual spray. Mix 1 1/2 teaspoons in 1 gallon of water. Use sufficient spray to obtain full and uniform coverage. Avoid spraying to the point of runoff. Adjust spray volume used to plant size.

ORNAMENTAL HOUSE PLANTS

Plants	Insects	Instructions & Restrictions
<p>INDOORS: To protect plants in greenhouses and lathhouses, and indoor landscaping. Ageratum, Ardisia, Aster, Azalea, Baby's Breath, Begonia, Birdsnest Fern, Bleeding Heart, Carnation, Chrysanthemum, Coleus, Crown of Thorns, Cockscomb, Cyclamen, Dracaena, Dumbcane, English Ivy, Fuchsia, Gladiolus, Grape Ivy, Hypoestes Ivy, Juniper, Lily, Marigold, Orchid, Palm, Pansy, Peperomia, Petunia, Philodendron,</p>	<p>Aphids, Armyworms, Corn Earworms, Exposed Thrips, Cankerworms (Fall and Spring), Inchworms, Japanese Beetles, Leafminers, Leafrollers, Mealybugs, Scales, Spider Mites, Thrips (exposed), Whiteflies</p>	<p>Spray directly on insects. Remove infested plants from living areas or areas likely to be occupied by children, pets (particularly fish and cats) prior to applications. When plants are dry, they can be returned to their original location. Do not use more product than is recommended as this may cause plant injury. Use with care on ferns, palms and fuchsia. As with all plants listed above, especially carnations, chrysanthemums and roses, there exists a wide variety of species and cultivars, and, therefore, a wide variety of susceptibility to injury. (Con't on next page.)</p>

ORNAMENTAL HOUSE PLANTS (Continued)

Plants	Insects	Instructions & Restrictions
(Continued) Piggyback Plant, Poinsettia, Portulaca, Pothos, Prayer Plant, Purple Passion, Rabbits Foot Fern, Rose, Schefflera, Snapdragon, Snake Plant, Statice, Velvet Plant, Verbena, Zinnia	(Continued)	(Continued) To test for possible phytotoxicity, spray a few plants as described above, wait 24 hours, and check for possible injury. Repeat applications at 4 to 14 days intervals maybe made only as necessary for insect control. Repeat applications may be necessary to control adult Japanese beetles on roses.
Chrysanthemums	Armyworms, Cabbageworms, Cutworms, Leafrollers	
Velvet Plants	Fungus Gnats	

ROSES AND FLOWERS

Mix 1 1/2 teaspoons in 1 gallon of water. Apply as a uniform spray, thoroughly wetting all plant parts. Do not spray to the point of runoff. Try to cover underside of leaves and penetrate dense foliage. Adjust spray volume used to plant size.

Plants	Insects	Instructions & Restrictions
Ageratum, Aster, Azalea, Begonia, Coleus, Common Ninebark and Snowberry, Exacum, Gladiolus, Gold Bells, Hypoestes, Ivy, Lilac, Marigold, Mock-orange, Nannyberry, Orchid, Pansy, Pea Shrub, Petunia, Poinsettia, Rhododendron, Rose, Snapdragon, Zinnia	Aphids, Bagworms, Cicadas, Cankerworms (fall), Inchworms, Leafminers, Leafrollers, Mealybugs, Spider Mites, Thrips (exposed), Webworms (fall), Whiteflies	Spray in the late afternoon or evening when the temperature ranges from 50° to 75°F and when there is little or no wind. Spray at the first sign of insects. Repeat at 4 to 8 day intervals only as necessary for insect control. Application can be made up to the day of collection.

ORNAMENTAL TREES AND SHRUBS

Mix 1 1/2 teaspoons in 1 gallon of water. Apply as a uniform spray, thoroughly wetting leaves and branches. Do not spray to the point of runoff. Try to cover underside of leaves and penetrate dense foliage. Adjust spray volume used to plant size.

Plants	Insects	Instructions & Restrictions
Arizona Cypress, Azalea, Birch, Non-bearing Cherry, Non-bearing Citrus, Conifers, Elm, English Ivy, Euonymus, Weeping Fig, Fir, Honeysuckle, Lilac, Mock-orange, Oak, Palm, Philodendron, Pine, Poinsettia, Tulip, Rhododendron, Taxus	Aphids, Bagworms, Cicadas, Cankerworms (fall), Inchworms, Leafminers, Leafrollers, Mealybugs, Spider Mites, Thrips (exposed), Webworms (fall), Whiteflies	Spray in late afternoon or evening, when the temperature ranges from 50° to 75°F and when there is little or no wind. Spray at the first sign of insects. Repeat applications at 4 to 8 day intervals only as needed to control infestations. Applications can be made up to the day of collection.

ORNAMENTAL TREES AND SHRUBS *(Continued)*

Plants	Insects	Instructions & Restrictions
Conifer	Budworms	Spray in late afternoon or evening, when the temperature ranges from 50° to 75°F and when there is little or no wind. Spray at the first sign of insects. Repeat applications at 4 to 8 day intervals only as needed to control infestations. Applications can be made up to the day of collection.
Elm	Elm Leaf Beetles, Elm Spanworms, Fall Webworms	
Douglas and Grand Firs	Tussock Moths	
Non-bearing Citrus	Citrus Black Flies	
Oak	Oakworms	
Pine	Pine Beetles, Pine Moths, Pine Needle Miners, Needle Scales	

To Control and Prevent Bark Beetles & Boring Insects (including, but not limited to: Ash Borer, Bronze Birch Borer, Rhododendron Borer, Elm Bark Beetles and Turpentine Beetles): Mix 1 ½ to 3 fl. ozs. of this product per 1 gallon water. Treat lower branches and trunk areas until thoroughly wet but avoid applying to the point of runoff. Apply prior to adult emergence. Use the higher rate for heavy pest infestation. Use the lower rate for preventative control. Apply in early May and again in early June.

NUISANCE PESTS IN OUTDOOR AREAS

Mix at the rate of 6 $\frac{2}{3}$ fl. oz. per 1 gallon water to treat 1,600 linear feet (or 1 $\frac{2}{3}$ fl. oz. per quart water to treat 400 linear feet) with a 6-inch spray band.

Where	Insects	Instructions & Restrictions
Outside Surfaces of Buildings	Ants, Clover Mites, Crickets, Earwigs, Millipedes, Sowbugs (Pillbugs)	To help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of buildings, including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen. Repeat treatment every 14 days only as needed to maintain insect control.
	Ants, Crickets, Earwigs, Fleas, Lone Star Ticks	For localized infestations of these insects in areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For Ants, thoroughly wet hills and runways. Repeat applications only when reinfestation occurs.

NUISANCE PESTS IN OUTDOOR AREAS (Continued)

Where	Insects	Instructions & Restrictions
Other Outdoor Areas	Flies (Cluster and House), Flying Moths, Gnats, Mosquitoes	Use as an aid in reducing the annoyance from these insects, spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages and other places where these insects alight or congregate.
Other Outdoor Areas	Tent Caterpillars	Apply when Caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when Caterpillars have returned to their tents. Wet tents with spray on all sides. A few Caterpillars may be seen escaping from the tent; however, they should die within several hours. Treat new tents that may appear.

FOR LAWN AREAS (Bentgrass, Bermudagrass, Bluegrass, Dichondra, Fescue, Irish Moss, Merlon & St. Augustine Grass): For control of *Ants, Armyworms, Brown Dog Ticks, Chinch Bugs, Crickets, Cutworms, Fire Ants, Fleas, Grasshoppers, Japanese Beetle Grubs, Mole Crickets, Sod Webworms*, consult the following chart:

Pests	Instructions & Restrictions
Armyworms, Cutworms, Chinch Bugs, Fleas, Japanese Beetle Grubs, Mole Crickets, Sod Webworms, Brown Dog Ticks, Crickets, Grasshoppers, Ants	For best results, thoroughly wet down grass at least 24 hours before applying. Lawn soil should be moist prior to application. Home lawns should be no taller than 3 inches at time of application. Slightly more water may be used as long as the amount of product per area is used as follows: For heavy infestations, repeat application after 2 weeks. For Ants, Crickets, Grasshoppers mix 1/2 fl. oz. (1 Tbs.) in 3 gallons of water and apply to cover 1,000 sq. ft. For all other insects, mix 1 1/2 fl. oz. (3 Tbs.) in 10 gals. of water and apply to cover 1,000 sq. ft.

Pests	Instructions & Restrictions
Fire Ants	<p>Mix 1 1/2 fl. oz. (3 Tbs.) in 1 gallon of water to treat one mound. Apply the solution as a gentle rain to each Fire Ant mound using a sprinkler can. Thoroughly wet the mound and the surrounding area to a 4-foot diameter. For best results, apply in cool weather, 65° - 85° F, or in early morning or late evening hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment. Treat all mounds in the vicinity, and treat all colonies which have not yet constructed mounds. Do not use equipment that produces a pressurized spray since this will disturb Fire Ants and cause migration, reducing product effectiveness. Use this same method for Fire Ants that exist in containerized plants.</p>

VEGETABLES (For Home Vegetable Gardens Only): Mix at the rate in the table below in 4 gallons of water to cover 4,000 square feet. Spray at first sign of insects. Wet the plants for thorough and uniform coverage but avoid runoff. Try to cover the underside of leaves and penetrate dense foliage. Treat in late afternoon or evening, when the temperature is cooler and when there is little or no wind.

Vegetable	Target Pest(s)	Instructions & Restrictions	Rate/4 Gal. Water
Asparagus (pre-harvest)	Cutworm, Asparagus Beetle	Do not apply more than 4 applications per season. Can be applied up to 3 days prior to harvest.	1 ½ fl. oz. (3 Tbs)
Asparagus (post-harvest)	Larval stages of Asparagus Beetle, Tarnished Plant Bug, Lygus Bugs and Adult Japanese Beetle	Treat ferns or brush growth after spear harvest when insects, are present.	1 ½ fl. oz. (3 Tbs)
Celery - See Leafy Vegetables			

Vegetable	Target Pest(s)	Instructions & Restrictions	Rate/4 Gal. Water
Cole Crops: Broccoli, Brussels Sprouts, Cabbage, Cauliflower	Beet Armyworm, Diamond Back Moth, Imported Cabbageworm, Cabbage-aphid, Cabbage Looper	Apply every 5 days only as needed to maintain insect control. Can be applied up to 1 day prior to harvest. Do not make more than 8 applications on Cauliflower, Brussels Sprouts and Broccoli and 10 applications on Cabbage per season.	1 fl. oz. (2 Tbs)

Vegetable	Target Pest(s)	Instructions & Restrictions	Rate/4 Gal. Water
Cucurbits: Balsam Pear (Bitter Melon), Chinese Wax Gourd (Chinese Preserving Melon), Citron Melon, Cucumber, Gherkin, Edible Gourds; Melons including hybrids such as Cantaloupe, Casaba, Crenshaw, Honeydew, Honey Balls, Mango Melon, Muskmelon, Persian Melon, Pumpkin, Summer & Winter Squash, Watermelon	Aphids, Cabbage Looper, Cucumber Beetle (Adults), Cutworms, Leafhoppers, Leafminers, Melonworms, Pickleworm, Plant Bugs, Squash Bugs, Flea Beetle	Do not make more than 8 applications per season. Ensure uniform coverage. Can be applied up to day of harvest	1 fl. oz. (2 Tbs.)

Vegetable	Target Pest(s)	Instructions & Restrictions	Rate/4 Gal. Water
Eggplant	Colorado Potato Beetle, Vegetable Leaf Miner	Do not exceed 16 applications per season. Can be applied up to 3 days prior to harvest.	3 fl. oz. (6 Tbs.)
Horseradish (foliar application)	Imported Crucifer Weevil	Do not exceed 3 foliar applications per season. Can be applied up to 22 days prior to harvest.	3 fl. oz. (6 Tbs)

Vegetable	Target Pest(s)	Instructions & Restrictions	Rate/4 Gal. Water
Leafy Vegetables: Amaranth, Arrugula, Celery, Celtuce, Chervil, Corn Salad, Chrysanthemum (edible-leaved & garland), Cress (garden & upland), Dandelion, Dock, Endive, Fennel, Lettuce (head & leaf) Orach, Parsley, Purslane (garden & winter), Rhubarb, Spinach, Swiss Chard	Alfalfa Looper, Aphids, Beet Armyworm, Cabbage Looper, Corn Earworm, Cutworms, European Corn Borer, Fall Armyworm, Green Cloverworms, Southern Armyworm, Tobacco Budworm, Vegetable Leafminer	Apply every 5 to 10 days only as needed for insect control. Do not make more than 7 applications per crop. Can be applied up to 1 day of harvest.	2 fl. oz. (4 Tbs)
Melons-See Cucurbits			

Vegetable	Target Pest(s)	Instructions & Restrictions	Rate/4 Gal. Water
Potatoes	Beet Armyworm, Colorado Potato Beetle, Potato Leafhopper, Potato Flea Beetle, Aster Leafhopper, Potato Aphid, Potato Tuberworm, Cabbage Looper, Cutworms, European Corn Borer, Potato Psyllid, Tarnished Plant Bug	Do not apply more than 12 applications. Can be applied up to 7 days prior to harvest.	1 1/2 fl. oz. (3 Tbs)
Peppers	Vegetable Leafminer, Cabbage Looper, Flea Beetle, Corn Earworm, Pepper Weevil, European Corn Borer	Do not make more than 8 applications per season. Can be applied up to 3 days prior to harvest.	2 fl. oz. (4 Tbs.)

Vegetable	Target Pest(s)	Instructions & Restrictions	Rate/4 Gal. Water
Strawberries, Raspberries, Blueberries	Strawberry Weevil, Aphids, Spider Mite, Lygus Bug, Flea Beetle, Omnivorous Leafhopper, Spittlebug, Japanese Beetle, Field Cricket, Red Spider, Leafroller, Weevils	Do not make more than 8 applications per season. Do not apply within 14 days of harvest.	1 1/5 fl. oz. (7 tsp.)
Sweet Corn	Corn Earworm, European Corn Borer, Fall Armyworm, Southern Armyworm	Apply every 5 days only as needed to maintain pest control. Do not make more than 6 applications per season. Can be applied up to 1 day prior to harvest.	1 1/2 fl. oz. (3 Tbs.)

Vegetable	Target Pest(s)	Instructions & Restrictions	Rate/4 Gal. Water
Tomatoes (Do not apply on Cherry Tomatoes and varieties yielding Tomatoes less than one inch in diameter.)	Beet and Southern Armyworm, Tomato Pinworm, Tomato Fruitworm, Horn Worms, Granulate Cutworms, Colorado Potato Beetle, Cabbage Looper	Do not make more than 6 applications per season. Can be applied up to date of harvest.	1 1/2 fl. oz. (3 Tbs.)

TREE FRUIT AND NUT CROPS GROWN AROUND THE HOME:

Mix 2 fl. oz. (4 Tbs.) in 4 gallons of water. Spray at the first sign of insects. Wet trees for thorough and uniform coverage but avoid applying to the point of runoff. Try to cover underside of leaves and penetrate dense foliage.

Trees (Fruit Or Nut)	Target Pest(s)	Instructions & Restrictions
Almonds, Pecans	Navel Orange Worm, Peach Twig Borer	Apply when insects appear. For Almonds, do not make more than 2 applications during the period when the outer hull of the almond has started to open. Do not make more than 5 applications per season. Can be applied up to 7 days prior to harvest.
Apples	Plum Curculio, Redbanded Leafroller, Rosy Apple Aphid, Spotted Tentiform Leafminer, Tarnished Plant Bug, White Apple Leafhopper, Obliquebanded Leafroller, Green Fruitworm, Japanese Beetles, Codling Moth	Apply when insects appear. Do not make more than 3 applications per season. Do not apply after petal fall.

Trees (Fruit Or Nut)	Target Pest(s)	Instructions & Restrictions
Peaches	Lesser Peachtree Borer, Peach Twig Borer, Plum Curculio, Tarnished Plant Bug, Green Fruit Worm, Oriental Fruit Moth	Apply when insects appear. Do not apply within 7 days of harvest. Do not make more than 8 applications per season.
Pears	Aphids, Pear Psylla, Green Fruitworm, and Codling Moth	Apply when insects appear. Do not apply more than 3 applications per season. Can be applied up to 14 days prior to harvest.

ANIMAL AND ANIMAL QUARTERS: For use on dogs and residential (personal) domestic animal use.

RESTRICTIONS

- **DO NOT APPLY TO CATS.**
- Do not apply this product in barns or stables where animals intended for slaughter or human consumption will be maintained.
- Do not apply when food, feed or water is present.

FOR USE ONLY ON DOGS AND LISTED

"DOMESTIC" LIVESTOCK

Treatment Of Animals And Animal Quarters: To aid in control of Horn Flies, Face Flies, Houseflies, Mosquitoes, Gnats, Stable Flies, Horse Flies, Deer Flies, Blood Sucking Lice, Poultry Lice, Bedbugs, Mites, Fleas and Ticks: Mix this product at a rate of 1 part concentrate in 19 parts water (6.7 fluid ounces product per 1 gallon water) or as directed below. When used in dairy barns or facilities, close milk bulk tank lids to prevent contamination.

For Livestock: To protect cattle (beef or dairy), goats, sheep, hogs and horses from Horn Flies, Face Flies, Houseflies, Mosquitoes, and Gnats: Mix this product at a rate of 1 part concentrate in 19 parts water (6.7 fluid ounces product per 1 gallon water). Apply a light mist sufficient to wet the surface of the hair. For Face Flies, spray face and head, but do not spray into the eyes.

To Control Stable Flies, Horse And Deer Flies: Apply at a rate of 2 fluid ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or only at intervals as necessary to provide continued protection.

To Control Blood-Sucking Lice: Apply to the infested areas of the animal using a stiff brush to move the spray to the base of the hair. Mix this product at a rate of 1 part concentrate in 19 parts water (6.7 fluid ounces product per 1 gallon water). Repeat treatment every 3 weeks only as necessary for Lice protection.

To Control Poultry Lice: Mix this product at a rate of 1 part concentrate in 19 parts water (6.7 fluid ounces product per 1 gallon water). Spray roosts, walls and nests or cages to thoroughly wet but not to the point of runoff. This should be followed by spraying over the birds with a fine mist.

To Control Bedbugs And Mites In Poultry Houses: Mix this product at a rate of 1 part concentrate in 19 parts water (6.7 fluid ounces product per 1 gallon water). Thoroughly spray crevices of roost poles, cracks in walls, and cracks in nests where the Bedbugs and Mites hide. Do not apply to the point of runoff.

On Dogs: To kill Fleas, Ticks, Lice - Mix this product at a rate of 1 part concentrate in 19 parts water (6.7 fluid ounces product per 1 gallon water). Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly. Reapply every 35 days only as needed to control pests. Do not spray into eyes or face. Avoid contact with genitalia. **Do not treat puppies less than twelve weeks old.** Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing dogs, or dogs on medication. **DO NOT use this product on cats or kittens.** Sensitivities may occur after using any pesticide product for dogs. If signs of sensitivity occur, bathe your dogs with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

WOOD-INFESTING INSECTS INCLUDING TERMITES*:
For Beetles (Furniture, Deathwatch, False Powderpost, Longhorned, Powderpost), Carpenter Ants and other wood-infesting Ants, Carpenter Bees and Termites* including Dry-wood Termites*: Treat outside surfaces of buildings, porches, wooden decks and patios, wooden fences, window frames, doorways, foundations, eaves, patios, garages, garbage areas and other areas where you may find these pests. Mix 2 1/2 fl. ozs. with 1 gallon of water to treat 100 sq.ft. Spray infested surfaces until thoroughly wet, avoiding dripping and runoff. Apply evenly on wood surfaces. To treat small areas, apply by brushing or spraying the diluted spray. For large areas, use a coarse, low pressure spray. Repeat treatment only as needed to maintain effectiveness. Do not spray into electrical outlets or utilities.

Spray infested areas and areas likely to be infested, such as wood surfaces, voids and channels in damaged wood, spaces between wooden parts of a structure, and between wood and foundations where infestation is likely to occur, also around fenceposts, wood piles, (do not use on firewood) and tree stumps.

Suitable protective clothing should be worn in confined or overhead areas to prevent exposure to eyes and skin. The use of eye protection, protective head covering, chemical resistant gloves and footwear, a long-sleeved shirt and long pants or coveralls are recommended. Do not permit contact with treated surfaces until spray has dried. Treated confined areas such as detached garages, under porches and decks, should be ventilated during and immediately following application before the areas are re-occupied.

FOR CARPENTER ANTS: Spray around doors and windows and other places where Ants may enter or crawl and hide. Also treat cracks and crevices or openings into wall voids or other areas where these Ants may be present.

FOR CARPENTER BEES: Spray or inject into holes and tunnels already made by the Bees with a tank sprayer, and spray areas where they are beginning to build tunnels into the wood.

FOR TERMITES* INCLUDING DRY-WOOD TERMITES*: Treat as for other wood-infesting insects. This product will kill workers and winged reproductive forms of Termites in the treated areas. Re-treat only as needed to maintain effectiveness.

* For localized control only. Such applications for Termites are not a substitute for mechanical alternation, soil treatment or foundation treatment, but are merely a supplement. For active Termite infestations, get a professional inspection.

SOIL TERMITE TREATMENT DIRECTIONS

SUBTERRANEAN TERMITES: DO NOT USE INSIDE DWELLINGS. For use around existing structures, not for pre-construction use. For post-construction treatment around residential buildings.

Precaution: When treating adjacent to an existing structure, the applicator must check the area to be treated, and immediately adjacent areas of the structure, for visible and accessible cracks and holes to prevent any leaks or significant exposures to persons occupying the structure. People present or residing in the structure during application must be advised to remove their pets and themselves from the structure if they see any signs of leakage. After application, the applicator is required to check for leaks. All leaks resulting in the deposition of termiticide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site. Treated confined areas such as detached garages, under porches and decks should be ventilated during and immediately following application before the areas are re-occupied.

Wear suitable protective clothing (goggles, protective head covering, chemical resistant gloves, footwear, a long-sleeved shirt and long pants or overalls) in confined or overhead areas to prevent exposure to eyes and skin.

Do not allow people or pets to contact contaminated areas or to reoccupy contaminated areas of the structure until the clean up is completed.

Mix 6.7 fl. oz. of this product per 1 gallon of water (makes 0.5% water emulsion). Apply as specified below. One treatment should last up to 5 years if treated soil is left undisturbed. Kills and controls subterranean termites.

Chemicals for soil treatment are used to establish a barrier against termites. The chemical emulsion must be adequately dispersed in the soil to provide a barrier between the wood in the structure and the termite colonies in the soil. Barriers may be established in the soil on the outside perimeter of the house by trenching using a rate of 1 gallon of emulsion per 2 ½ linear feet per foot of depth [or, 4 gallons of emulsion per 10 linear feet per foot of depth]. A complete treatment may require treatment with another EPA registered product to the inside perimeter of the foundation and to other crucial areas. For these situations, contact a licensed pest control operator for treatment. Do not use under slabs, in crawl spaces, or indoors. If termites are found within the house or structure, contact a licensed pest control operator for treatment.

Remove all wood debris and wood forms. Around piers, pipes, chimney bases and along foundations of poured concrete, dig a trench 6 inches wide and 4 inches deep. For brick and block foundations, dig the trench about 12 inches deep. If the footing is more than 12 inches deep, make holes with a crowbar, pipe or rod about 1 foot apart that extend from the trench bottom to the top of the footing. Make holes closer in hard-packed clay soils. Never dig below the top of the footing. Apply the prepared emulsion (or solution) in the trench and to the backfill at the rate of 1 gallon per 2 ½ linear feet per foot of depth.

For raised porches, terraces and entrance slabs outdoors, drill holes at one foot intervals, 6 inches from the foundation, and pour one-half gallon of emulsion (or solution) into each hole. Refill holes.

Be sure to treat the interiors of accessible hollow block foundations at the same rate as above. Do not use in structural voids insulated with rigid foam. Be careful to have adequate ventilation when treating areas under structure. For severe infestations or treatment under slabs with heat ducts or radiating heat pipes, in basements and buildings with sufficient crawl space, etc., contact a licensed pest control operator or termite control service.

NOTE: Avoid contamination of public and private water supplies by following these precautions. Do not allow a hose or any type of faucet extension to reach into the termite chemical solution while filling or mixing this material with water. Use anti-backflow or anti-siphonage equipment on all filling equipment. These devices can be purchased at hardware or plumbing supply stores. Use of anti-backflow equipment will help ensure that the user of this chemical will not contaminate domestic water supplies.

Refer to FHA (Federal Housing Administration), state and local specifications for safe distances of treatment areas from wells. Do not treat structures that have cisterns, wells, French drains, or sump systems. Soil in the vicinity of wells, cisterns or ponds should not be treated. For those situations, contact the local Cooperative Extension Service or consult a commercial pest control service. Do not treat soil that is water saturated or frozen or in any conditions where runoff or movement from the treatment area (site) is likely to occur.

Retreatment for subterranean termites can only be performed if there is clear evidence of reinfestation or disruption of the barrier due to construction, excavation, or landscaping and/or evidence of the breakdown of the termiticide barrier in the soil. These vulnerable or reinfested areas maybe retreated in accordance with application techniques described in this product's labeling. The timing and type of these retreatments will vary, depending on factors such as termite pressure, soil types, soil conditions or other factors which may reduce the effectiveness of the barrier.

Annual retreatment of the structure is prohibited unless there is clear evidence that reinfestation or barrier disruption has occurred.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store product in its original container in a cool, dry, locked place out of reach of children and domestic animals. Always store container in an upright position.

PESTICIDE DISPOSAL: If empty: Non-refillable container. Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT. If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

This page left blank intentionally.

This page left blank intentionally.