

# ImiGold™ 0.5 G

Turf & Ornamental Insecticide

**For systemic insect control in turfgrass and landscape ornamentals**

**ACTIVE INGREDIENT**

Imidacloprid – 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine ..... 0.5%

**OTHER INGREDIENTS** ..... 99.5%

**TOTAL** ..... 100.0%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**FIRST AID**

<b>If Swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on Skin or Clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If in Eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Contact the Rocky Mountain Poison Center at 1-866-673-6671 for emergency medical treatment information.

**NOTE TO PHYSICIAN:** No specific antidote is available. Treat symptomatically.

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC at 1-800-424-9300**

EPA Reg. No. 70506-142  
EPA Est. No. 61842-CA-001

Batch/Lot # \_\_\_\_\_

**Net Contents: 30 lbs.**

 **UPI United Phosphorus, Inc.**  
630 Freedom Business Center, Suite 402  
King of Prussia, PA 19406  
1-800-438-6071 • www.upi-usa.com

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Caution. Harmful if swallowed or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

**ENVIRONMENTAL HAZARDS**

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not formulate this product into other end-use products.

**TURF & ORNAMENTAL APPLICATIONS INFORMATION**

**Use Precautions**

- Do not apply more than 80 lbs. ImiGold 0.5 G Insecticide (0.4 lb. a.i.) per acre per year.
- Apply so as to avoid runoff or puddling of irrigation water.
- Do not apply to areas which are waterlogged or saturated, which will not allow penetration into the root zone.
- Before application, ensure that application equipment is properly calibrated.
- Do not use this product on commercial sod farms.
- Do not use this product in commercial greenhouses, nurseries, on sod farms, or on grass grown for seed.
- Treated areas may be replanted with any crop specified on an Imidacloprid label, or with any crop for which a tolerance exists for the active ingredient. For crops not listed on an Imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12-month plant-back interval should be observed.

**Turfgrass Applications**

Apply ImiGold 0.5 G to turf and ornamentals in sites such as:

- |                               |                                    |
|-------------------------------|------------------------------------|
| Airports                      | Home lawns                         |
| Athletic fields               | Multi-family residential complexes |
| Business and office complexes | Parks                              |
| Cemeteries                    | Playgrounds                        |
| Golf courses                  | Shopping complexes                 |

Due to the residual activity of the active ingredient, applications of ImiGold 0.5 G Insecticide may be made before target pests lay their eggs. For best control, apply before the eggs hatch, basing the need for application on historical monitoring, current season trapping, or other methods. All applications should be followed by rainfall or irrigation to move the active ingredient through the thatch.

When infested turfgrass areas are waterlogged, or soil beneath turf is saturated with water, avoid making applications because it is difficult to obtain thorough and consistent distribution of the product. For best results, ensure that rainfall or irrigation after application will vertically penetrate the soil, allowing the active ingredient to be carried into the zone where the pests are found.

PESTS	RATE IMIGOLD 0.5 G	INSTRUCTIONS
Larvae of: Annual bluegrass weevil Asiatic garden beetle Billbugs Black turfgrass ataeenius Cutworms (suppression) European chafer European Crane Fly Green June beetle Japanese beetle Northern masked chafer Oriental beetle <i>Phyllophaga</i> spp. Southern masked chafer	50 to 80 lbs. per acre or 1.2 to 1.8 lbs. per 1000 sq. ft.	For best control of grubs, billbugs, European crane fly and annual bluegrass weevil, apply before egg hatch. <u>Chinchbugs (suppression)</u> : apply before first instar nymphs hatch. <u>Mole crickets</u> : apply before or during the peak egg hatch period. When adults or large nymphs are present and actively tunneling, ImiGold 0.5 G Insecticide application should be accompanied by a curative insecticide. Consult your local State Agricultural Experiment Station or State Extension Turf Specialists for assistance regarding timing of application. NOTE: For best control, irrigation or rainfall should occur within 24 hours after application to move the active ingredient through the thatch. On golf courses, irrigate treated areas following application. Avoid mowing turf or lawn area until sufficient irrigation or rainfall has occurred. Do not apply more than 80 lbs. ImiGold 0.5 G Insecticide (0.4 lb. of active ingredient) per acre per year.
Chinch bug (suppression) Mole crickets	80 lbs. per acre or 1.8 lbs. per 1000 sq. ft.	

### Ornamental Applications

Apply ImiGold 0.5 G Insecticide to plants used for aesthetic purposes or climatic modification that are being grown in interior landscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

ImiGold 0.5 G Insecticide can be used as a broadcast application on ornamentals in commercial and residential landscapes. Because ImiGold 0.5 G Insecticide is a systemic product that is translocated upward within the plant system, best control will be achieved when applications are followed by sufficient mechanical incorporation, irrigation or rainfall to move the active ingredient into the soil. Apply before anticipated pest infestation for best control.

#### Application

Apply ImiGold 0.5 G Insecticide uniformly over the area being treated with drop-type or rotary-type spreaders. Avoid the use of spreaders which apply material in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions, and check frequently to be sure equipment is working properly and distributing granules uniformly and accurately.

ORNAMENTAL APPLICATIONS		
Landscape plantings including trees, shrubs, evergreens, flowers, foliage plants, ground covers		
PESTS	RATE	INSTRUCTIONS
White grub larvae (such as Japanese beetle larvae, Chafers, <i>Phyllophaga</i> spp., Asiatic garden beetle, Oriental beetle)	1.4 to 1.8 lbs. per 1000 sq. ft.	Apply as a broadcast treatment. Before applying, remove black plastic or other weed barrier material that will not permit water to reach soil surface. NOTE: For best control, mechanical incorporation, irrigation or rainfall should occur within 24 hours after application to move the active ingredient into the soil. Refer to label sections below for application equipment and spreader settings.

### Spreader Settings for ImiGold 0.5 G Insecticide

The following chart can be used as a guide in establishing an accurate rate of ImiGold 0.5 G Insecticide. Application rates will vary depending upon speed, equipment wear and condition of spreader. Carefully calibrate application equipment prior to commercial use of this product.

SPREADER	SPEED (mph)	SETTINGS	
		60 lbs./A	80 lbs./A
Cyclone Model CB4000	3	3.25	3.50
Earthway Drop	3	9	10
Earthway 10' Rotary	3	11	12
Gandy Turf Tender 24"	3	16	19
Gandy Turf Tender 42"	3	21	23
Lely	4.5	3.50	-
Lesco Model 6000	3	B ½	C
PSB Prizelawn CBR-11	3	3.25	3 1/2
Scott's Proturf R-8A	3	H	I
Scott's Proturf SR-1	3	H	I
Scott's PF-1, PF-2, PF-3, PF-4 Drop	3	4	4.50
Scott's SS-1 Drop	3	4	4.50
Spyker Model 34	3	3.1	3.2
Vicon Model PS-203	4.5	13	16

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**Pesticide Storage:** Store in a cool, dry area and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material is spilled for any reason or cause, carefully sweep material into a pile and dispose of as directed for pesticides below. Refer to Precautionary Statements on label for hazards associated with the handling of this material. In spill or leak incidents, keep unauthorized people away.

**Pesticide Disposal:** Pesticides wastes are toxic. Improper disposal of excess pesticide, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional Office for guidance.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## IMPORTANT INFORMATION READ BEFORE USING PRODUCT

### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of United Phosphorus, Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of United Phosphorus, Inc. and Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold United Phosphorus, Inc. and Seller harmless for any claims relating to such factors.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UNITED PHOSPHORUS, INC. AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.**

To the extent consistent with applicable law, United Phosphorus, Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UNITED PHOSPHORUS, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UNITED PHOSPHORUS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

United Phosphorus, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of United Phosphorus, Inc.

ImiGold is a trademark of United Phosphorus, Inc.

© 2009 United Phosphorus, Inc. All rights reserved.

Rev. 7/31/09