



Imida^{Pro}2SC

Insecticide

NET CONTENTS: 1 Gallon

Insecticide for foliar and systemic insect control in turf grass (including sod farms), landscape ornamentals, fruit and nut trees and interior plantscapes.

ACTIVE INGREDIENT:

Imidacloprid, 1 -[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidimine21.4%

OTHER INGREDIENTS:.....78.6%

TOTAL:100.0%

Contains 2 pounds of imidacloprid per gallon.

EPA Reg. No. 83529-2-72159

EPA Est. No. 37429-GA-01

DISTRIBUTED BY:

AGRISEL
AGRI-CHEMICALS
Quality & Better Value

P.O. Box 3528 • Suwanee, GA 30024

SHAKE WELL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE INSIDE BOOKLET.

Net Contents 1 Quart (32 oz)

FIRST AID

IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.

For 24-Hour medical emergency assistance (human or animal) call in case of emergency call CHEMTREC toll free at 1-800-222-1222. For chemical emergency assistance (spill, leak, fire, or accident) call CHEMTREC at 1-800-424-9300.

Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Keep children or pets off treated area until spray is dry.

Personal Protective Equipment (PPE):

Applicators and Other Handlers Must Wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves of any weatherproof material such as barrier

laminate, butyl rubber, nitrile rubber, neoprene rubber, polyethylene, polyvinyl chloride (PVC) or viton. If you want more options, follow the instructions for category A on an EPA chemical-resistant category chart.

- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

User Should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product.

Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly into water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift into blooming crops or weeds if bees are visiting the treatment area.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not formulate this product into other end-use products.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent): Then offer to recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticides as directed below. In spill or leak accidents, keep unauthorized people away. You may contact CHEMTREC at 1-800-424-9300 for decontamination procedures or any other assistance that may be necessary.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the areas during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours..

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene runner, natural rubber, polyethelene, polyvinyl chloride (PVC) or viton.
- Shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and pets off treated area until dry.

APPLICATION TO TURFGRASS

ImidaPro 2SC will control the following soil inhabiting pests found in turfgrass.

PEST	SCIENTIFIC NAME
Northern & Southern masked chafers	<i>Cyclocephala borealis</i> , <i>C. immaculata</i> , and/or <i>C. lurida</i>
Asiatic garden beetle	<i>Maladera castanea</i>
European chafer	<i>Rhizotrogus majalis</i>
Green June beetle	<i>Cotinis nitida</i>
May or June beetle	<i>Phyllophaga spp.</i>
Japanese beetle	<i>Popillia japonica</i>
Oriental beetle	<i>Anomala orientalis</i>
Billbugs	<i>Spherophorus spp.</i>
Annual bluegrass weevil	<i>Hyperodes spp.</i>
Black turfgrass ataenius	<i>Ataeneus spretulus</i> and <i>Aphodius spp.</i>
European Crane Fly	<i>Tipula paludosa</i>
Mole crickets	<i>Scapteriscus spp.</i>

ImidaPro 2SC will suppress cutworms and chinch bugs.

ImidaPro 2SC can be applied on turfgrass in the following sites:

Home lawns	Golf courses	Playgrounds
Business and office complexes	Airports	Athletic fields
Shopping complexes	Cemeteries	Sod farms
Multi-family residential complexes	Parks	

ImidaPro 2SC has adequate residual activity that applications can be made preceding the egg laying activity of the target pests. High levels of control can be achieved when applications are made prior to or during egg hatch of the target pests. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Optimum control will be achieved when applications are made prior to egg hatch of target pests, followed by sufficient irrigation or rainfall is required to move the active ingredient through the hatch.

Avoid application when infested areas are waterlogged or soil beneath turf is saturated with water. These conditions prevent thorough and consistent distribution. Best results are achieved when rainfall or irrigation after application will penetrate vertically in the soil column carrying the active ingredient into the zone where insects are normally located.

Do not exceed a total of 1.6 pints application (0.4 lb active ingredient) per acre per year.

APPLICATION EQUIPMENT FOR USE IN TURFGRASS

ImidaPro 2SC should be diluted with enough water to provide adequate volume to promote thorough distribution into the pest zone. Use only accurately calibrated equipment for application to turfgrass. Apply a uniform, coarse droplet spray, using a low pressure setting to eliminate off target drift. Calibration should be performed on a regular basis to ensure that equipment is distributing product properly.

APPLICATION TO ORNAMENTALS

ImidaPro 2SC can be applied to ornamental plants in commercial and residential landscapes and interior plantscapes. ImidaPro 2SC is systemic insecticide that is transported within the plant system from the roots to upper foliage. ImidaPro 2SC must be applied into a growing area of the plant that allows absorption of the active ingredient. Adding soluble nitrogen type fertilizers to the spray solution when appropriate can promote the uptake of the active ingredient. Application can be made by foliar application or soil applications; including soil injection, drenches, and broadcast sprays. Foliar applications offer locally systemic activity against insect pests. The systemic translocation of the active ingredient will be slower when applied to woody plants with soil applications. This delay can take 60 days or longer depending on species and size of plant. To offset this, make applications before anticipated pest infestation.

For outdoor ornamentals, broadcast applications cannot exceed a total of 1.6 pints (0.4 lb of active ingredient) per acre per year.

ANT MANAGEMENT PROGRAMS

Using ImidaPro 2SC to control aphids, scale insects, mealybugs and other sucking pests on ornamentals with ant populations because it removes honeydew as a food source. To enhance control of ants, supplement with residual sprays, bait placements or other ant control tactics.

NOTE: Not for use in commercial greenhouses, nurseries, or on grasses grown for seed, or on commercial fruit and nut trees.

APPLICATION EQUIPMENT FOR FOLIAR APPLICATIONS

ImidaPro 2SC mixes readily with water and may be used in many types of application equipment. Mix ImidaPro 2SC with the required amount of water and apply as desired dependent upon the selected use pattern. Add a commercial spreader/sticker promote coverage on hard to wet foliage such as holly, pine, or ivy. If concentrate or mist type spray equipment is used, an equivalent amount of product should be used on the area sprayed, as would be used in a dilute application.

ImidaPro 2SC is compatible with many commonly used fungicides, miticides, liquid fertilizers, and other insecticides.

If applicator has prior experience with a particular tank mix, physical compatibility should be checked by making a small clear jar test using correct proportions of products to be tank mixed.

Do not apply through any irrigation system.

RESTRICTIONS

- Do not graze treated areas or use clippings from treated areas for feed or forage.
- Avoid runoff or puddling of irrigation water following application.
- Keep children and pets off treated areas until dry.
- Avoid application of ImidaPro 2SC to areas which are water logged or saturated, which will not allow penetration into the root zone of the plant.
- Do not apply more than 1.6 pt (0.4 lb. of active ingredient) per acre per year.
- Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop which a tolerance exists for the active ingredient.
- For crops not listed on an imidacloprid label, or for crops for which no tolerance for the active ingredient have been established, 12-month plant-back interval should be observed.

Use Site: TURF GRASSES

To control larvae of:

European Crane fly
Annual bluegrass weevil
Green June beetle
Asiatic garden beetle
Japanese beetle

Billbug
Northern masked chafer
Black turfgrass ataenius
Oriental beetle

Cutworms (suppression)
Phyllophaga spp.
European chafer

Apply 1.25 to 1.6 pints per acre (equivalent to 0.46 to 0.6 fl. oz. (14 to 17 mL) per 1,000 sq. ft.). Make application prior to egg hatch of grubs, billbugs, annual bluegrass weevil, and European Crane fly to maximize control. Be sure to read "Application Equipment" Section of this label. For chinch bugs (suppression) and mole crickets apply 1.6 pints per acre (equivalent to 0.6 fl. oz. (17 mL) per 1,000 sq. ft.). For suppression of chinch bugs, make application before the hatching of the first instar nymphs. For control of mole crickets make application before or during

the peak egg hatch period. If adults or large nymphs are actively tunneling, ImidaPro 2SC application should be combined with a curative insecticide. Follow label instructions for other insecticides when tank-mixing. Consult your local turf, state Agricultural Experiment Station, or State Extension Service Specialist for more specific information regarding timing of application.

Notes and Restrictions for Turf Grass:

Irrigation or rainfall should occur within 24 hours after application to move the active ingredients through the hatch.

Do not apply more than 1.6 pt. (0.4 lb. of active ingredient) per acre per year.

Avoid mowing turf or lawn area, until after adequate irrigation or rainfall has occurred so that uniformity of application will not be affected.

Use Site: TREES, SHRUBS, EVERGREENS, FLOWERS, FOLIAGE PLANTS, GROUNDCOVERS AND INTERIOR PLANTSCAPES

(Only in and around Industrial and Commercial Buildings and Residential Areas)

To control:

Adelgids	Leafhoppers
Aphids	(including glassy-winged sharpshooter)
Japanese beetles	Mealybugs
Lace bugs	Psyllids
Leaf beetles	Sawfly larvae
(including elm and viburnum leaf beetles)	Thrips (suppression)
	Whiteflies

Apply 1.5 fluid ounces (45 mL) per 100 gal. of water. Make foliar applications before high pest populations become established. Reapply on an as needed basis. To control White grub larvae, (such as Japanese beetle larvae, Chafers, *Phyllophaga* spp., Asiatic garden beetle, Oriental beetle) apply 0.46 to 0.6 fluid ounces (14 to 17 mL) per 1,000 sq. ft. Make broadcast applications by mixing specified rate in enough water to uniformly cover the treated area. Do not use less than 2 gallons of water per 1,000 sq. ft. If necessary, irrigate thoroughly to incorporate ImidaPro 2SC into the upper soil profile. Refer to use directions specific for FLOWERS and GORUNDCOVERS concerning

additional use directions.

Use Site: TREES SHRUBS, FLOWERS AND GROWDCOVERS

(Only in and around Industrial and Commercial buildings and residential areas, and state, national, and private wooded and forested areas)

To control:

Adelgids

Japanese beetles

Leafminers

Sawfly larvae

Aphids

Lace bugs

Mealybugs

Soft scales

Armored scales (suppression)

Leaf beetles

(including elm and viburnum leaf beetles)

Pine tip moth larvae

Eucalyptus longhorned borer

Flatheaded borers (including bronze birch and alder borer)

Thrips (suppression)

Black vine weevil larvae

Psyllids

White grub larvae

Asian longhorned beetle

Leafhoppers

(including glassywinged sharpshooter)

Royal Palm Bugs

Whiteflies

Emerald ash borer

For TREES, apply 0.1 to 0.2 fl. oz. (3 to 6 mL) per inch of trunk diameter (D.B.H.).

Application Technique:

Soil Injection

GRID SYSTEM: Make applications in a grid pattern on 2.5 foot centers within the drip line of a tree.

CIRCLE SYSTEM: Make applications in holes evenly spaced approximately 2-3 feet apart in a circle within the drip line of the tree. Larger trees may require additional application circles.

BASAL SYSTEM: Make applications into holes evenly spaced around the base of the tree trunk no more than 6 to

12 inches from the base.

Soil Drench

Apply no less than 10 gallons of water per 1,000 square feet as a drench around the base of the tree, directed to the root zone. Any plastic or other barrier that may prevent drench solution from reaching the root zone must be removed.

Notes and Restrictions for Trees:

- Use sufficient water to be able to inject equal amount of solution into each hole.
- Use low pressure and sufficient solution for thorough distribution into the treatment zone.
- Maintain soil moisture for 7 to 10 days.
- Do not use less than 4 holes per tree.
- No Soil Injection Applications Allowed In Nassau or Suffolk Counties of New York.
- Application to trees already heavily infested with borers listed may not prevent the eventual loss of trees.

For SHRUBS apply 0.1 to 0.2 fl. oz. (3 to 6 mL) per foot of shrub height.

Application Techniques:

Soil Injection:

Apply to individual plants using dosage indicated.

Soil Drench:

Uniformly apply the dosage in no less than 10 gallons of water per 1,000 square feet as a drench around the base of the tree, directed to the root zone. Remove any plastic or any other barrier that will stop solution from reaching the root zone.

Notes and Restrictions for Shrubs:

- Mix required dosage in sufficient water to inject an equal amount of solution into each hole.
- Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Maintain soil moisture for 7 to 10 days.
- Do not use less than 4 holes per shrub.

• No Soil Injection Applications Allowed in Nassau or Suffolk Counties in New York.

For FLOWERS and GROUNDCOVERS apply 0.46 to 0.6 fluid ounces (14 to 17 mL) per 1,000 sq. ft.

Apply as a broadcast treatment and incorporate into the soil before planting or after plants are established. If application is made to established plants, best control is achieved by irrigating after application.

Use Site: POME FRUITS

(Only in and around Residential Areas)

Includes: Apple, Crabapple, Loquat, Mayhaw, Pear, Pear (Oriental), Quince

To control:

Aphids

(except Woolly apple aphid)

Leafhoppers

(including glassy-winged sharpshooter)

Leafminer

Mealybugs*

San Jose Scale*

- Apply 1.5 fluid ounces (45 mL) per 100 gal or 6.0 fluid ounces per acre. Application rate per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees.

Notes and Restrictions for residential Pome Fruits:

- Apply specified dosage as foliar spray as needed after petal-fall is complete.
- For control of rosy apple aphid, apply prior to leafrolling caused by the pest.
- For first generation leafminer control, make first application as soon as petal fall is complete. Greatest leafminer control will result from the earliest possible application. For second and succeeding generations of leafminer, optimal control is obtained from applications made early in the adult flight against egg and early instar larvae. A second application may be required 10 days later if severe pressure continues or if generations are overlapping. A single application may result in suppression only. ImidaPro 2SC will not control late stage larvae.
- For San Jose Scale, time applications to the crawler stage. Treat each generation.
- For late season (preharvest) control of leafhopper species, apply ImidaPro 2SC while most leafhoppers are in the nymphal stage.
- For optimal control of mealybug, insure good spray coverage of the trunk and scaffolding limbs or other resting

sites of the mealybug.

- Do not apply more than 6.0 fluid ounces per acre in a single application. Do not make more than 5 applications.
 - Allow 10 or more days between applications. Allow at least 7 days between last application and harvest.
- Not for use in California for control on pears.

PECAN*

(Only in and around residential areas)

(*Use on pecans not permitted in California unless directed by specific supplemental labeling)

To control:

Yellow pecan aphid

Black margined aphid

Pecan leaf phylloxera

Pecan spittlebug

Pecan stem phylloxera

Apply 1.5 fl. oz. (45 mL) per 100 gal or 6.0 fl. oz./A¹. Make foliar applications as pest pressure begins to increase. Make a second application 10 to 14 days after first if field scouting reveals continued pest pressure. Use of an organosilicone based spray adjuvant at recommended rate can insure thorough coverage of foliage.

Notes and Restrictions for residential Pecan Trees.

- Do not apply more than a total of 18.0 fluid ounces of ImidaPro 2SC per acre per year.
- Do not make more than 3 applications.
- ¹Application rate per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees.

GRAPES

(For use only in and around industrial and commercial buildings and residential areas)

To control Leafhoppers (including glassy-winged sharpshooter) and Mealybugs apply 1.5 fl. oz. (45 mL) per 100 gal. or 3.0 fl. oz./A (90 mL/A). Apply label rate as a foliar spray using 200 gallons of water per acre.

Notes and Restrictions for Residential Grapes:

- Do not apply more than a total of 6.0 ounces of ImidaPro 2SC per acre per year.
- Allow at least 14 days between applications.
- Applications may be applied up to and including day of harvest.

CONDITIONS OF SALE AND WARRANTY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, use or buyer accepts the following Conditions, Disclaimer or Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Agrisel USA, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, AGRISEL USA, INC. MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Agrisel USA, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. AGRISEL USA, INC. DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT AGRISEL USA, INC.'S ELECTION, THE REPLACEMENT OF PRODUCT.