



# Insecticide in Water-Soluble Packets

For foliar and systemic insect control in turfgrass (Including sod farms), landscape ornamentals, fruit and nut trees, and interior plantscapes.

### **ACTIVE INGREDIENT:**

lmidacloprid, 1-[(6-Chloro-3-pyridinyl) methyl]-N-nitro-2-imidazolidinimine	. 75.0%
OTHER INGREDIENTS:	. 25.0%
TOTAL:	100.0%

Do not remove packets from container except for immediate use.

Keep water soluble packets in this container and store in a cool, dry place but not below freezing (32° F).

# KEEP OUT OF REACH OF CHILDREN CAUTION

EPA Reg. No. 53883-513 EPA Est. No. 33967-NJ-001

Manufactured for: Control Solutions, Inc. 5903 Genoa Red Bluff Road Pasadena, Texas 77507

	FIRST AID
IF SWALLOWED	<ul> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
IF ON SKIN OR CLOTHING	<ul> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of soap and water for 15 to 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
IF IN EYES	<ul> <li>Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product contain	er or label with you when calling a poison control center or doctor or going for treatment. You may also contact SafetyCall® (8)

897-8050 for emergency medical treatment information.

NOTE TO PHYSICIAN No specific antidote is available. Treat the patient symptomatically

## PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or vapor. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Keep children and pets away from treated area until dry.

## Personal Protective Equipment (PPE)

WPS Uses: Applicators and other handlers who handle this product for any use covered by the Worker Protection Standard (40 CFR part 170) – in general, agricultural plant use e.g. use on sod farms, must wear:

Long-sleeved shirt and long pants

- Chemical resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.
- · Shoes plus socks

Follow manufacturer's instructions for cleaning, maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Non-WPS Uses: Applicators and other handlers who handle this product for any use NOT covered by the Worker Protection Standard (40 CFR part 170)

- Shirt and pants
- · Shoes plus socks

## **Engineering Controls Statements:**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## **User Safety Recommendations**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
   Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change

# **ENVIRONMENTAL HAZARDS**

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops/plants or weeds. Do not apply this product or allow it to drift to blooming crops/plants or weeds if bees are foraging. This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater ntamination.

## PROTECTION OF POLLINATORS



other insect pollinators.

APPLICATION RESTRICTIONS EXIST FOR THIS PRODUCT BECAUSE OF RISK TO BEES AND OTHER INSECT POLLINATORS. FOLLOW APPLICATION RESTRICTIONS FOUND IN THE DIRECTIONS FOR USE TO PROTECT POLLINATORS.



in the Directions for Use for each application site for specific use restrictions and instructions to protect bees and

## This product can kill bees and other insect pollinators.

Bees and other insect pollinators will forage on plants when they flower, shed pollen, or produce nectar.

Bees and other insect pollinators can be exposed to this pesticide from:

- Direct contact during foliar applications, or contact with residues on plant surfaces after foliar applications
- Ingestion of residues in nectar and pollen when the pesticide is applied as a seed treatment, soil, tree injection, as well as foliar applications.

When Using This Product Take Steps To:

 Minimize exposure of this product to bees and other insect pollinators when they are foraging on pollinator attractive plants around the application site.
 Minimize drift of this product on to beehives or to off-site pollinator attractive habitat. Drift of this product onto beehives or off-site to pollinator attractive habitat can result in bee kills.

nformation on protecting bees and other insect pollinators may be found at the Pesticide Environmental Stewardship website at: http:// pesticidestewardship.org/PollinatorProtection/Pages/default.aspx.

Pesticide incidents (for example, bee kills) should immediately be reported to the state/tribal lead agency. For contact information for your state, go to: www.aapco.org/officials.html. Pesticide incidents should also be reported to the National Pesticide Information Center at: www.npic.orst.edu or directly to EPA at: beekill@epa.gov

## Do not formulate this product into other end-use products.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

See individual sites for specific pollinator protection application restrictions. If none exist under the specific site, for foliar applications, follow these application directions for food/feed crops and commercially grown ornamentals that are attractive to pollinators, and for non-agricultural uses

FOR FOOD/FEED CROPS AND COMMERCIALLY GROWN ORNAMENTALS NOT UNDER CONTRACT FOR POLLINATION SERVICES BUT ARE ATTRACTIVE TO POLLINATORS

Do not apply this product while bees are foraging. Do not apply this product until flowering is complete and all petals have fallen unless one of the following conditions is met:

- The application is made to the target site after sunset
- The application is made to the target site when temperatures are below 55°F
- The application is made in accordance with a government-initiated public health response
  The application is made in accordance with an active state-administered apiary registry program where beekeepers are notified no less than 48-hours
- prior to the time of the planned application so that the bees can be removed, covered or otherwise protected prior to spraying
- The application is made due to an imminent threat of significant crop loss, and a documented determination consistent with an IPM plan or predetermined economic threshold is met. Every effort should be made to notify be the bees can be removed, covered or otherwise protected prior to spraying.



Non-Agricultural Uses:

Do not apply QUALI-PRO IMIDACLOPRID 75 WSB Insecticide while bees are foraging. Do not apply QUALI-PRO IMIDACLOPRID 75 WSB Insecticide to plants that are flowering. Only apply after all flower petals have fallen off.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted- entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has be

- eated, such as plants, soil, or water is:

  Coveralls

  Chemical resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep children and pets off treated area until dry.

State Specific Restrictions

. The state of Arizona has not approved this product for use on agricultural sites. Do not use this product on uses considered by the Arizona statutes to

## TURFGRASS (INCLUDING SOD FARMS, LANDSCAPE ORNAMENTALS, FRUIT & NUT TREES, AND INTERIOR PLANTSCAPES)

Mixing: Inside each foil pouch is a clear inner water soluble packet containing the QUALI-PRO IMIDACLOPRID 75 WSB. Do not allow packets to become wet prior to adding to the spray tank. Do not handle the clear inner packets with wet hands or wet gloves. Rough handling may cause breakage. Reseal outer carton to protect remaining packets.

To prepare the spray mixture, remove the outer foil pouch and drop the required number of unopened, clear water soluble packets, as determined under **Applications**, into the spray tank while filling with water to the desired level. Operate the agitator while mixing. Depending on water temperature and amount of agitation, the packets should completely dissolve within a few minutes after they are added to the water. Allow longer time for the clear packet to dissolve when using cooler water.

Do not use QUALI-PRO IMIDACLOPRID 75 WSB packets in a tank-mix with products that contain Boron or release free chlorine. The resultant reaction of PVA and boron or free chlorine is a plastic which is not soluble in water or solvents such as diesel oils, kerosene, gasoline or alcohol. Do not attempt to use the packets directly in diesel oils or summer spray type oils as in ULV or LV uses. PVA packets are water soluble not oil soluble. Use of chlorinated water is acceptable.

QUALI-PRO IMIDACLOPRID 75 WSB has been found to be compatible with commonly used fungicides, miticides, liquid fertilizers, and other commonly used insecticides. Check physical compatibility using the correct proportion of products in a small jar test if local experience is unavailable.

# **Restrictions**Do not apply through any irrigation system.

Do not graze treated areas or use clippings from treated areas for feed or forage. Avoid runoff or puddling of irrigation water following application. Keep children and pets off treated area until dry. Do not apply QUALI-PRO IMIDACLOPRID 75 WSB to areas which are water logged or saturated, or frozen, which will not allow penetration into the root zone of the plant. Do not apply more than 8.6 oz. (0.4 lb of active ingredient) per acre per year.

Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient. For crops not listed on an imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12- month plant-back interval must be observed.

Not for use in commercial greenhouses, nurseries, or on grasses grown for seed, or on commercial fruit and nut trees.

Do not apply this product, by any application method, to linden, basswood or other Tilia species.

### TURFGRASS

QUALI-PRO IMIDACLOPRID 75 WSB can be used as directed on turfgrass in sites such as home lawns, business and office complexes, shopping centers, residential complexes, golf courses, airports, cemeteries, parks, playgrounds, athletic fields and sod farms to control soil inhabiting pests as listed in the table below. as in sites such as home lawns, business and office complexes, shopping

Effective control can be achieved when applications are made before or during the egg laying period. The timing of applications can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. For best control, apply prior to egg hatch of the target pests followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

Irrigation or rainfall must occur within 24 hours after application to move the active ingredient through the thatch. Do not make applications when turfgrass areas are waterlogged or the soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The area to be treated must be able to accept vertical penetration of rainfall or irrigation water. Avoid mowing turf or lawn area until after sufficient irrigation or rainfall has occurred so that uniformity of application will not be affected.

Do not apply more 8.6 oz. (0.4 lb of active ingredient) per acre per year.

Application Instructions
Apply QUALI-PRO IMIDACLOPRID 75 WSB in sufficient water to provide adequate distribution in the treated area. The use of accurately calibrated equipment normally used for the application of turfgrass insecticides is required. Use equipment which will produce a uniform, coarse-droplet spray, using a low pressure setting to eliminate off target drift. Check calibration periodically to ensure that equipment is working properly.

APPLICATION INSTRUCTIONS ON TURFGRASS			
PEST	USE DIRECTIONS	RATE	
Larvae of: European Crane Fly, (Tipula paludosa), Annual bluegrass weevil (Hyperodes spp.), Green June beetle (Cotinis nitida), May or June beetle (Phyllophaga spp.), Asiatic garden beetle (Maladera castanea), Japanese beetle (Popillia japonica) Billbug (Spherophorus spp.), Northern masked chafer and Southern masked chafer (Cyclocephala borealis, C. immaculata, and/or C. lurida), Black turfgrass ataenius (Ataenius spretulus and Aphodius spp.) Oriental beetle (Anomala orientalis), Cutworms (suppression) (Phyllophaga spp.), European chafer (Rhizotroqus majalis)	For best results, for the control of grubs, billbugs, European crane fly and annual bluegrass weevil, make application prior to egg hatch of the target pest.  Refer to <b>Application Instructions</b> section of the label.	1.6 oz. (1 packet) per 8,250 to 11,000 sq. ft.	
Chinchbug (suppression), Mole crickets ( <i>Scapteriscus</i> spp.)	For suppression of chinchbugs, make application prior to hatching of the instar nymphs.  For control of mole crickets make application prior to or during the peak egg hatch period. When adults or large nymphs are present and actively tunneling, application should be accompanied by a curative insecticide. Follow label instructions for other insecticides when tank-mixing.	1.6 oz. (1 packet) per 8,250 sq. ft.	

# RESTRICTIONS

- Do not apply this product in a way that will contact people or pets.
  Do not allow children or pets to enter treated areas until sprays have dried.
  Do not apply to plants in bloom or allow this product to contact plants if bees are foraging.

## **ORNAMENTALS**

QUALI-PRO IMIDACLOPRID 75 WSB is for use on ornamentals in com nercial and residential landscapes and interior plantscapes

QUALI-PRO IMIDACLOPRID 75 WSB mixes with water and may be used in a variety of application equipment. Mix product with the required amount of water and apply as directed for the selected use pattern

When making foliar applications on hard-to-wet foliage such as holly, pine, or ivy, the addition of a spreader/sticker is recommended. If concentrate or mist type spray equipment is used, an equivalent amount of product should be used on the area sprayed, as would be used in a dilute application. **Application Instructions** 

## OLIALI-PRO IMIDACI OPRID 75 WSB is a systemic product and will be translocated upward into the plant system from root untake. Place the product where the growing portion of the target plant can absorb the active ingredient. The addition of a nitrogen containing fertilizer, where applicable, into the solution may enhance the uptake of the active ingredient. Application can be made by foliar application or soil applications; including soil injection, drenches, and broadcast sprays. Foliar applications offer locally systemic activity against insect pests.

When making soil applications to plants with woody stems, systemic activity will be delayed until the active ingredient is translocated throughout the plant. In some cases, this translocation delay could take 60 days or longer. For this reason, applications should be made prior to anticipated pest infestation to achieve optimum levels of control.

For outdoor ornamentals, broadcast applications cannot exceed a total of 8.6 oz (0.4 lb of active ingredient) per acre per year

## Ant Management Programs

Use QUALI-PRO IMIDACLOPRID 75 WSB to control aphids, scale insects, mealybugs and other sucking pests on ornamentals to limit the honeydew available as a food source for ant populations. QUALI-PRO IMIDACLOPRID 75 WSB applications can then be supplemented with residual sprays, bait placements or other ant control tactics to further reduce the pest population

APPLICATION INSTRUCTIONS  Trees, Shrubs, Evergreens, Flowers, Foliage Plants, Ground Covers, and Interior Plantscapes For use only in and around industrial and commercial buildings and residential areas.		
PEST	RATE	INSTRUCTIONS
Adelgids Aphids Japanese beetles Lace bugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassywinged sharp- shooter) Mealybugs Psyllids Sawfly larvae Thrips (suppression) Whiteflies	1.6 oz. (1 packet ) per 300 gals. of water	Foliar Applications: Start treatments prior to establishment of high pest populations and reapply on an as needed basis.
White grub larvae (such as Japanese beetle larvae, Chafers, <i>Phyllophaga</i> spp. Asiatic garden beetle, Oriental beetle)	1.6 oz. (1 packet) per 8,250 to 11,000 sq. ft.	Broadcast Applications: Mix required amount of product in sufficient water to uniformly and accurately cover the area being treated. Do not use less than 2 gallons of water par 1000 sq ft. Irrigate thoroughly to incorporate QUALI-PRO IMIDACLOPRID 75 WSB into the upper soil profile.  Refer to FLOWERS and GROUNDCOVERS for additional use directions.

## APPLICATION INSTRUCTIONS

## cial buildings and residential areas and state, national, and private wooded and

forested areas to control the insect pests listed below.			
Adelgids Aphids Armored scales (suppression) Black vine weevil larvae	TREES 1.6 oz. (1 packet)	Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. Do not use less than 4 holes per tree.	
Eucalyptus longhorned borer Flatheaded borers (including bronze birch borer and alder borer) Japanese beetles	per 24 to 48 inches	SOIL INJECTION': For application to trees - Grid System: Holes should be spaced on 2.5 foot centers, in a grid pattern, extending to the drip line of the tree.	
Lace bugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassywinged sharpshooter) Leafminers		Circle System: Apply in holes evenly spaced In circles, (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line.	
		Basal System: Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base.	
Mealybugs		For application to shrubs - Apply to individual plants using dosage indicated.	
Pine tip moth larvae Royal Palm Bugs Sawfly larvae Soft Scales Thrips (suppression) White grub larvae Whiteflies		SOIL DRENCH: Uniformly apply the dosage in no less than 10 gallons of water per 1000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.	
		For Control of Specified Borers: Application to trees already heavily infested may not prevent the eventual loss of the trees due to existing pest damage and tree stress.	
	FLOWERS & GROUNDCOVERS	Apply as a broadcast treatment and incorporate into the soil before planting, or apply prior to bloom or after all flower petals have fallen off for established plants. If application is made to established plants, irrigate thoroughly after application.	
	1.6 oz. (1 packet) per 8,250 to 11,000 sq. ft.		

No Soil Injection Applications Allowed in Nassau or Suffolk Counties of New York.

### RESTRICTIONS

- Do not apply this product in a way that will contact people or pets.

APPLICATION INSTRUCTIONS For use only in and around residential areas.			
CROP	PEST	RATE	INSTRUCTIONS
Apple Cychemple (avecut Mechycopy) (1 poetset)	Apply specified dosage as foliar spray as needed after petal-fall is complete.  For control of rosy apple aphid, apply prior to leafrolling caused by the pest.		
Loquat, Mayhew, Pear, Pear (Oriental), Quince	aphid) Leafhoppers (including glassywinged sharpshooter) Leafminer Mealybugs² San Jose Scale²	per 300 gals. of water (2.1 oz. per acre¹)	For first generation leafminer control, make first application as soon as petal-fall is complete Greatest control will result from the earliest possible application. For second and succeeding generations, best control is obtained from applications made early in the adult flight agains egg and early instar larvae. A second application may be required 10 days later if severe pressure continues or if generations are overlapping. A single application may result is suppression only. QUALI-PRO IMIDACLOPRID 75 WSB will not control late stage larvae.
			For San Jose Scale, time applications to the crawler stags. Treat each generation.
			For late season (preharvest) control of leafhopper species, apply QUALI-PRO IMIDACLOPRID 75 WSB while most leafhoppers are In the nymphal stage.
			For optimal control of mealybug, insure good spray coverage of the trunk and scaffolding limbs or other resting sites.
			Do not apply more than 2.1 ounces per acre in a single application. Do not make more than applications or exceed a total of 8.6 ounces (0.4 lb ai) per acre per year.
		Allow 10 or more days between applications. Allow at least 7 days between last application and harvest.	
Pecan <sup>3</sup>	Yellow pecan aphid Black margined aphid Pecan leaf phyfloxera	1.6 oz. (1 packet) per 300 gals.	Make foliar applications as pests begin to build before populations become extreme. Two applications at a 10 to 14 day interval may be required to achieve control. Scout and retreal friededc.
Pecan spittlebug Pecan stem phyfloxera (2.7)	of water (2.1 oz. per acre¹)	Thorough uniform coverage of foliage is necessary for optimal control. Addition of ar organosilicone-based spray adjuvant at a rate not to exceed the adjuvant manufacturer's recommended use rate may improve coverage.	
			Do not apply more than a total of 6.3 ounces of QUALI-PRO IMIDACLOPRID 75 WSB pe acre per year. Do not make more than 3 applications. Allow 10 or more days between applications. Allow at least 7 days between last application and harvest.

standard of 400 gallons of dilute spray solution per acre for large trees.

Not for use in California for control on pears

Use on pecans not permitted in California unless directed by state-specific 24(c) labeling

APPLICATION INSTRUCTIONS  For use only in and around industrial and commercial buildings and residential areas.			
CROP	PEST	RATE	INSTRUCTIONS
Grapes	Leafhoppers (including glassy-winged sharpshooter) Mealybugs	1.6 oz. (1 packet) per 300 gal. of water (1.0 oz. per acre)	Apply specified dosage as a foliar spray using 200 gallons of water per acre. Do not apply more than a total of 2.0 ounces of QUALI-PRO IMIDACLOPRID 75 WSB per acre per year. Allow at least 14 days between applications. Applications may be applied up to and including day of harvest.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or dispo

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Do not store below freezing (32°F). Exposure to moisture or excessive handling of water soluble packets may cause breakage. Store water soluble packets in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site (in the treatment area) or at an approved was

CONTAINER DISPOSAL: Nonrefillable Container (flexible-bag-all weights): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Then dispose of empty bag in sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable Container (rigid-fifty lbs. or less): Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Handle and open container carefully. Do not cut water soluble packets when opening. If container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to **PRECAUTIONARY STATEMENTS** on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed above. In spill or leak incidents, keep unauthorized people away

# LIMITATION OF WARRANTY AND LIABILITY

Read the entire directions for use, conditions of warranties and limitat ons of liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following CONDITIONS, DISCLAIMER OF WARRANTIES, and LIMITATIONS OF LIABILITY.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Control Solutions, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law. Control Solutions, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Control Solutions, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Control Solutions, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Control Solutions, Inc.'s election, the replacement of product.

Quali-Pro is a registered trademark of Makhteshim Agan of North America. Inc