



# ***IPRODIONE 2F SELECT™***

FUNGICIDE

## ***Professional Non-Residential Disease Control***



### **ACTIVE INGREDIENT**

Iprodione

### **% ACTIVE INGREDIENT**

23.0%

### **FORMULATION**

Flowable

### **CHEMICAL FAMILY**

Dicarboximide

### **MODE OF ACTION**

Signal Transduction

### **RESTRICTED USE**

None

### **SIGNAL WORD**

Caution

### **PACKAGING INFORMATION**

2 x 2.5 gal.

### **LABELED USES**

Golf Courses  
Non-Residential  
Ornamentals  
Non-Residential Turf  
Sod Farms

**Iprodione 2F Select** is a foliar-applied fungicide that may be deployed for the control of disease on non-residential turf including golf courses, sod farms and institutional areas. It can be affectively applied on spring, summer, fall and winter diseases. Iprodione 2F Select is compatible and may be applied with, most commonly used, registered fungicides including TM 4.5F Select to control a broad spectrum of disease and aid in resistance management.

- Up to 28 days residual
- Effective for all seasons
- Non-Residential use only
- Preventive and curative effects
- Labeled for Turf and Ornamental Applications

**IMPORTANT:  
ALWAYS READ AND FOLLOW  
LABEL INSTRUCTIONS.**



**PRIME SOURCE**

# IPRODIONE 2F SELECT™

FUNGICIDE

## **Turf Applications:**

Applied according to good turf management practices, Iprodione 2F Select is an effective tool in controlling spring, summer and fall diseases and in controlling certain winter diseases.

## **Summer Stress Complex/Summer Decline:**

Tank-mix applications of 4 to 8 fl. oz. per 1,000 sq. ft. of Chipco® Signature™ Fungicide with 3 to 4 fl. oz. per 1,000 sq. ft. Iprodione 2F Select to help manage these diseases.

## **Pythium Blight:**

Pythium blight will be controlled by a tank-mix of Iprodione 2F Select with one of the following: fosetyl-al or propamocarb hydrochloride fungicides. Follow the label directions and use precautions for all tank-mix partners and ensure that the use of that fungicide is registered for use on turf.

## **Gray Snow Mold (Typhula spp.):**

To use Iprodione 2F Select as a tank-mix in sites where continuous snow cover occurs, use 4 to 8 fl. oz. per 1,000 sq. ft. with one of the following registered products: chlorothalonil, such as Chlorothalonil 720 Select, or pentachloronitrobenzene (PCNB) fungicides. Use the rates according to the directions on the labels of the registered tank mix partner products. For optimum control, application must be made in autumn prior to snow cover. Ensure thorough coverage is attained by dilution in sufficient water (1 to 5 gallons of spray solution per 1,000 sq. ft.). If the turf has been frozen prior to application, use the higher rate of Iprodione 2F Select, and make a repeat application only if a thaw occurs and all snow cover has melted.

## **Restrictions:**

- Do not apply to turf cut higher than 1" on golf holes where water bodies are present.
- Except for use on golf courses, if applying this product adjacent to a water body such as a lake, reservoir, river, permanent stream, marsh or natural pond, estuary, or commercial fish pond, there must be at least a 25-foot vegetative buffer strip between the water body and the point of application.
- Do not mow or irrigate treated areas until the grass is completely dry, which typically takes 24 hours post-treatment.
- Do not mix with any sticker, extender, or wetting agents.
- Do not apply more than 35 fl. oz. product/1,000 sq. ft. per year.
- Do not apply more than 6 applications per year.
- Do not allow animals to graze on treated turf and do not feed clippings from treated turf to livestock or poultry.



## **LABELED TURF DISEASES INCLUDE:**

Anthracnose  
Brown Patch  
Corticium Red Thread  
Curvularia  
Dollar Spot  
Fusarium Blight  
Fusarium Patch  
Gray Snow Mold  
Large Patch  
Leaf Spot  
Necrotic Ring Spot  
Pink Snow Mold

## **LABELED ORNAMENTAL DISEASES INCLUDE:**

Aerial Web Blight  
Alternaria Leaf Blight  
Alternaria Leaf Spot  
Blossom Blight  
Botrytis Blight  
Botrytis Storage Rot  
Cylindrocladium Blight and Wilt  
Daffodil Leaf Scorch  
Fusarium Corm Rot  
Fusarium Leaf Spot  
Helminthosporium Leaf Spot  
Ink Spot  
Ray Blight  
Rhizoctonia Stem and Root Rot  
Tulip Fire



Prime Source, LLC  
10025 US 264 Alternate  
Middlesex, NC 27557  
[www.Prime-Sourcecellc.com](http://www.Prime-Sourcecellc.com)

Download a  
specimen label at  
[Prime-Sourcecellc.com](http://Prime-Sourcecellc.com)