

- Water-based fogging concentrate
- Effective and fast acting flushing action, knockdown and kill of insects

KEEP OUT OF REACH OF CHILDREN

### CAUTION

See back for first aid and precautionary statements.

#### ACTIVE INGREDIENTS:

Pyrethrins.....	1.00%
* Piperonyl butoxide, Technical.....	2.00%
** N-Octyl bicycloheptene dicarboximide .....	3.00%
OTHER INGREDIENTS .....	94.00%
	<hr/> 100.00%

\* Equivalent to 1.6% (butylcarbityl) (6-propylpiperonyl) ether and 0.4% related compounds

\*\* MGK® 264, Insecticide Synergist

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

#### READ ALL DIRECTIONS COMPLETELY BEFORE USE.

**GENERAL:** This concentrate is designed for use full strength or diluted as desired, according to the dilution table listed below. Achieve optimum performance using a mechanical aerosol generator to deliver ULV fog droplets sized less than 50 microns in diameter (with 80% or more of the droplets under 30 microns in diameter). Apply as a space, area or contact spray in residential and commercial structures and animal housing.

Do not apply directly to food. Do not allow spray to contact stored food or food contact surfaces. Do not apply as a space spray while food processing is underway. Cover or remove foods before treatment. Cover or wash all food processing surfaces and equipment with an effective cleaning compound followed by a potable water rinse. In the home cover all food processing surfaces, dishes, and utensils before treatment or thorough wash before use following treatment.

Avoid contamination of food or feedstuffs. Remove pets and birds and cover fish aquariums and turn off aquarium air pump motor before spraying.

**In health care facilities:** Remove patients from area to be treated prior to treatment. Ventilate treated area for two hours after spraying. Do not return patients to treated areas until ventilation is complete. When used as a coarse wet spray, the sprayed surfaces must be dried before patients are allowed to reoccupy treated areas.

**NOTE:** Some wood and plastic finishes can be marred or damaged by insecticide applications. To avoid damage, cover those items and do not treat directly. If in doubt, test by applying this material to an inconspicuous area or surface and evaluate before application.

**NOTE:** If there is a need to treat such areas as conduits, motor housings, junction and switch boxes or other electrical equipment, turn off electrical equipment AND shut off power to the area prior to application. After application, allow time for insecticide to dry before restoring power to the area, starting motors or placing switches in the "ON" position.

**AREAS for use in and around:**

- Animal Areas and Quarters
- Animal Shelters
- Apartments
- Arcades
- Athletic Fields
- Attics
- Auditoriums
- Bakeries
- Bars
- Basements
- Bathrooms
- Beverage Plants
- Boats
- Bottling Plants
- Brandy Storage Warehouses
- Breweries
- Brewery Warehouses
- Buildings
- Buses
- Cabinets
- Cabins
- Cafeterias
- Campers
- Canneries
- Cargo Holds
- Cattle Barns
- Churches
- Circus Tents
- City Missions
- Closets
- Club Houses
- Commercial Buildings
- Communication Centers
- Concession Stands
- Condominiums
- Correctional Facilities
- Court Houses
- Crawl Spaces
- Crematoriums
- Dairy and Livestock Barns
- Day Care Centers
- Delicatessens
- Departments Stores
- Diners
- Doghouses
- Drains and Sewers
- Dried Fruit Processing Plants
- Dried Fruit Warehouses
- Drugstores
- Dumpsters
- Dwellings
- Egg Processing Plants
- Factories
- False Ceilings
- Federally Inspected Meat and Poultry Establishments
- Flour Mills
- Food Handling Establishments
- Food Processing Plants
- Food Storage Facilities
- Freight Containers
- Fruit Packing Sheds
- Funeral Parlors
- Garages
- Garbage Bins
- Garbage Cans
- Garbage Compactors
- Garbage Trucks
- Gas Stations
- Goat Houses
- Grain Elevators
- Grain Harvesting and Handling Equipment
- Grain Mills
- Granaries
- Grocery Stores
- Grocery Warehouses
- Healthcare Facilities
- Homes
- Horse Stables and Barns
- Horse Trailers
- Hospices
- Hospitals
- Hotels
- Indoor Eating Establishments
- Industrial Installations
- Institutional Dining Areas
- Institutions
- Jails
- Kennels
- Laboratories
- Laundromats
- Lavatories
- Libraries
- Liquor Stores
- Livestock Barns
- Loafing Sheds
- Locker Rooms
- Marinas
- Mausoleums
- Meat Packing Plants
- Meat Rendering Plants
- Mess Halls
- Milk Rooms
- Mills
- Mini-storages
- Missions
- Mobile Canteens
- Morgues
- Mortuaries
- Motels
- Movie Theaters
- Museums
- Mushroom Processing Plants
- Nursing Homes
- Office Buildings
- Offices
- Other Public Buildings
- Parking Garages
- Passenger Rail Cars
- Peanut Processing Plants
- Peanut Storage Warehouses
- Pool Halls
- Poultry Houses
- Poultry Processing Plants
- Prisons
- Public Buildings
- Public Storages
- Rabbit Housing
- Rabbit Processing Plants
- Railroad Cars
- Recreational Vehicles
- Research Animal Quarters
- Restaurants
- Restrooms
- Retail Stores
- Rice Mills
- Schools
- Shade Houses
- Ship Holds
- Ships
- Smoke Shops
- Sports Facilities
- Sports Stadiums
- Stable Bins
- Stables
- Storage Bins (areas)
- Stored Food in Multi-walled Paper or Cloth Bags
- Stored Seed Warehouses
- Stores
- Supermarkets
- Swine Housing
- Taverns
- Textile Mills and Warehouses
- Theaters
- Tobacco Processing Facilities
- Tobacco Warehouses
- Tool Sheds
- Townhouses
- Trains
- Transportation Equipments
- Truck Trailers
- Trucks
- Tunnels, Utility
- Utility Rooms
- Vegetable Packing Facilities
- Vending Machines
- Veterinary Hospitals
- Wall and Equipment Voids (attics, walls, ceilings)
- Warehouses
- Washrooms
- Wholesale Stores
- Wine Cellars
- Wine Storage Warehouses
- Wineries
- Zoos

**FOR THE CONTROL OF INSECTS including:**

- Almond Moths
- American Black Flour Beetles
- Angoumois Grain Moths
- Ants
- Barn Flies
- Bean Weevils
- Bed Bugs
- Bees
- Beetles
- Black Carpet Beetles
- Black Fungus Beetles
- Blow Flies
- Book Lice
- Bottle Flies
- Boxelder Bugs
- Broad-horned Flour Beetles
- Brown Dog Ticks
- Brown Spider Beetles
- Cadelles
- Carpet Beetles
- Centipedes
- Cereal Beetles
- Cheese Mites
- Cheese Skippers
- Chocolate Moths
- Cigarette Beetles
- Clothes Moths
- Clover Mites
- Cluster Flies
- Cockroaches
- Cocoa Bean Moths
- Coffee Bean Weevils
- Confused Flour Beetles
- Corn Sap Beetles
- Crickets
- Dark Mealworms
- Darkling Beetles
- Depressed Flour Beetles
- Dermestid Beetles
- Drain Flies
- Dried Fruit Beetles
- Drugstore Beetles
- Earwigs
- Elm Leaf Beetles
- European Grain Moths
- Face Flies
- Firebrats
- Flat Grain Beetles
- Fleas
- Flies
- Flying Moths
- Foreign Grain Beetles
- Fruit Flies
- Fungus Beetles
- Fungus Gnats
- Gnats
- Grain Mites
- Granary Weevils
- Grasshoppers
- Ground Beetles
- Hairy Fungus Beetles
- Hairy Spider Beetles
- Hide Beetles
- Horn Flies
- Hornets
- House Flies
- Indian Meal Moths
- Khapra Beetles
- Large Black Flour Beetles
- Larger Grain Borers
- Lesser Grain Borers
- Lesser Houseflies
- Lesser Mealworms
- Longheaded Flour Beetles
- Maize Weevils
- Meal Worms
- Mediterranean Flour Moths
- Merchant Grain Beetles
- Mexican Grain Beetles
- Midges
- Miller Moths
- Millipedes
- Mites
- Mosquitoes
- Mushroom Flies
- Palmetto Bugs
- Pillbugs
- Pink Scavenger Caterpillars
- Red Flour Beetles
- Red Horned Grain Beetles
- Rice Flour Beetles
- Rice Moths
- Rusty Grain Beetles
- Saw-toothed Grain Beetles
- Scorpions
- Sheep Ticks
- Silverfish
- Skipper Flies
- Slender-horned Flour Beetles
- Small Flying Moths
- Small-eyed Flour Beetles
- Sowbugs
- Spider Beetles
- Spiders
- Spinous Ear Ticks
- Square-necked Grain Beetles
- Stable Flies
- Tobacco Moths
- Trogoderma Beetles
- Two Banded Beetles
- Vinegar Flies
- Warehouse Beetles
- Wasps
- Waterbugs
- Webbing Clothes Moths
- Weevils
- White Marked Spider Beetles
- Yellow Jackets
- Yellow Mealworms

**APPLICATION RATES**

Type of pest	Method of application	Recommended rate
Flying insects and related pests	As a ULV, fog or space spray	1/2 to 1 fluid ounce of concentrate per 1,000 cubic feet of space
	As a crack and crevice injection	1/2 to 1 fluid ounce of concentrate per 1,000 cubic feet of space
Crawling insects and related pests	As a ULV, fog or space spray	1 fluid ounce per 1,000 cubic feet of space
	As a contact, surface or crack and crevice spray	6 to 32 fluid ounces in one gallon of water per 1,000 square feet of surface area
	As a crack and crevice injection	1 fluid ounce per 1,000 cubic feet of space

## DILUTION TABLE

Application method	% pyrethrins in solution	Mixture	Application rate	Lbs. pyrethrins
ULV, fog or space spray	1.00%	Use full strength/undiluted	1 fluid ounce per 1,000 cubic feet of space	0.0006 lbs. per 1,000 cubic feet
	0.50%	1 part concentrate to 1 part water		0.00032 lbs. per 1,000 cubic feet
Contact, surface or crack and crevice spray	0.25%	32 fluid ounces of concentrate in one gallon of water	1 gallon of solution per 1,000 square feet of surface area	0.02 lbs. per 1,000 square feet
	0.15%	18 fluid ounces of concentrate in one gallon of water		0.012 lbs. per 1,000 square feet
	0.10%	12 fluid ounces of concentrate in one gallon of water		0.008 lbs. per 1,000 square feet
	0.05%	6 fluid ounces of concentrate in one gallon of water		0.004 lbs. per 1,000 square feet
Crack and crevice injection	1.00%	Use full strength/undiluted	1 fluid ounce per 1,000 cubic feet of space	0.0006 lbs. per 1,000 cubic feet
	0.50%	1 part concentrate to 1 part water		0.00032 lbs. per 1,000 cubic feet

## SPACE SPRAY APPLICATIONS

**General instructions for use as a space application in listed areas:** Close doors, windows and shut off all ventilating equipment. Use a mechanical aerosol generator to deliver ULV fog droplets sized less than 50 microns in diameter (with 80% or more of the droplets under 30 microns in diameter). Apply at the rate of 1/2 to 1 fluid ounce of undiluted concentrate per 1,000 cubic feet of space, or dilute with water according to the dilution table. Direct spray toward upper corners of room and ceiling. Do not remain in treated areas after application. Leave room closed for at least 1/2 hour. Ventilate thoroughly before occupants are allowed to re-enter. Repeat treatment when reinfestation occurs. Do not use equipment designed for deep void or crack and crevice injection for space spray applications in accessible areas.

Do not apply directly to food. Do not allow spray to contact stored food. Do not apply as a space spray while food processing is underway. Foods must be removed or covered during treatment. All food processing surfaces and equipment must be covered or washed with an effective cleaning compound followed by a potable water rinse. In the home all food processing surfaces, dishes and utensils must be covered during treatment or thoroughly washed before use. Food processing operations may continue when this product is applied as a wet spray as a spot, or crack and crevice treatment, with care and in accordance with the directions and precautions listed on this label.

**When applying as a space spray to kill flying insects:** Use a mechanical aerosol generator to deliver ULV fog droplets sized less than 50 microns in diameter (with 80% or more of the droplets under 30 microns in diameter). Following the general and general space spray instructions on the label, apply undiluted at a rate of 1/2 to 1 fluid ounce per 1,000 cubic feet. If diluted solution is preferred, mix 1 part concentrate to 1 part water and apply at the rate of 1 fluid ounce of diluted solution per 1,000 cubic feet of space.

**When applying as a space spray to kill crawling insects:** Use a mechanical aerosol generator to deliver ULV fog droplets sized less than 50 microns in diameter (with 80% or more of the droplets under 30 microns in diameter). Following the general use and general space spray instructions on the label, treat as a space spray. First, treat cracks and crevices in woodwork, walls, floors, behind and beneath all equipment, cupboards, obstructions or dark harborage areas, underneath sinks, behind pipes, under pallets and in all places that harbor insects. Contact as many insects as possible with the application. Then, apply undiluted concentrate at the rate of 1 fluid ounce per 1,000 cubic feet of space.

**Dried fruit products in storage or processing:** Following the general use and general space spray instructions, do not apply to fruit directly. Apply one fluid ounce of concentrate, or diluted concentrate, per 1,000 cubic feet of space. Disperse this product as a fine mist in the air above the trays and shelves. Repeat treatment when reinfestation occurs.

**For crack and crevice injection:** Inject into cracks and crevices or void spaces where insects may be harboring, living and breeding. Always follow equipment manufacturer's

instructions for proper equipment set up and operating pressures. Place injector tip into cracks, crevices, holes and other small openings. Release approximately one-second of product. For light infestations move injector tip along cracks while treating at the rate of three linear feet per second. For heavy infestations move injector tip along at one linear foot per second. For closed voids calculate the void's cubic areas and treat at the rate of five to ten seconds per three cubic feet. Several holes may be required in long-running voids.

## CONTACT OR SURFACE SPRAY APPLICATIONS

**General instructions for use as a contact or surface spray:** To kill accessible stages of listed insects, mix 6 to 32 fluid ounces of concentrate with water to make one gallon of spray solution. Using a compressed air sprayer or similar type of sprayer, apply as a coarse, wet spray. Apply 1 gallon of spray solution mixture per 1,000 square feet of surface area. To kill listed insects, spray thoroughly into all cracks, molding, crevices, walls, floors and other surfaces, storage and handling areas. Treat areas where insects hide, travel or rest, such as around drains, sinks, under pallets, wall voids, nests, ant trails, water pipes, around and behind heavy machinery and equipment. Contact as many insects as possible. Leave room ventilation or air conditioning equipment on during application. Food processing operations may continue when this mixture is applied as a spot or crack and crevice treatment, with care and in accordance with the directions and precautions listed on this label.

**For use in transit protection: Do not use in aircraft.** To kill accessible stages of listed insects, mix 6 to 32 fluid ounces of concentrate with water to make one gallon of spray solution. Using a compressed air sprayer or similar type of sprayer, apply as a coarse, wet spray. Apply 1 gallon of spray solution mixture per 1,000 square feet of surface area. Clean and spray truck beds, boxcars, ships holds and other listed transit equipment before loading. Thoroughly treat walls and floors. Spray each layer of cartons or containers lightly as they are stacked for transit.

**For use on stored grain products:** To assist in the kill of insects in stored grain, apply to the surface of binned grain at a rate of 1 gallon per 1,000 square feet. Apply at monthly intervals with the first appearance of insects. In case of severe infestations of flying moths, break up webbing before spraying and make a second application in two weeks

## ANIMAL HOUSING APPLICATIONS

**General instructions for use in and around animal housing and related facilities:** For use in and around listed livestock areas to kill accessible stages of listed insects. Do not apply spray directly to animals that will be used for milk or meat production. If facilities are to be used for milk and/or meat animals, remove animals before application. Milk and/or meat animals may re-occupy the facility once the insecticide has dried. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying.

**As a space spray in animal housing:** Following the general and general animal housing instructions, use a mechanical aerosol generator to deliver ULV fog droplets sized less than 50 microns in diameter (with 80% or more of the droplets under 30 microns in diameter). Apply at the rate of 1/2 to 1 fluid ounce of undiluted concentrate per 1,000 cubic feet of space above the animals, or dilute with water according to the dilution table. Direct application above the animals, towards upper portions of the enclosure, filling the room with mist. Repeat treatment when reinfestation occurs.

**As a contact or surface spray in animal housing:** To kill accessible stages of listed insects, mix 6 to 18 fluid ounces of concentrate with water to make one gallon of spray solution. Following the general and general animal housing instructions use a compressed air sprayer, or similar type of sprayer, to apply solution as a coarse, wetting spray. Apply approximately 1 gallon of spray solution mixture per 1,000 square feet of surface area. Spray walls, cracks and crevices, foundation, rafters, floor and any area where insects hide or rest. Animals may re-occupy the facility once the insecticide has dried.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in a cool, dry place. Keep container closed.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of container in a sanitary landfill or by other procedures approved by state and local authorities.

### CONTAINERS ONE GALLON AND UNDER:

**If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. For terrestrial uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your state water board or regional office of the EPA.

## KEEP OUT OF REACH OF CHILDREN

### CAUTION

### FIRST AID

#### IF SWALLOWED:

- Immediately call a poison control center or doctor.
- Do not induce vomiting unless told to do so by a poison control center or a doctor.
- Do not give any liquid to the person.
- Do not give anything by mouth to an unconscious person.

#### IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.

**JETSTREAM™**  
Water-Based Contact Insecticide

JetStream is a trademark of McLaughlin Gormley King Company. McLaughlin Gormley King Company, MGK and Premium Pyrethrum are registered trademarks of McLaughlin Gormley King Company. All other marks are property of their respective holders. Manufactured by McLaughlin Gormley King Company®. ©2009 McLaughlin Gormley King Company. All rights reserved.  
8810 Tenth Avenue North, Minneapolis, MN 55427  
037-1629/5M-0209

