



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

LUNA[®] EXPERIENCE

Reduced Application Rate for use in almond

EPA Reg. No. 264-1091

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF ARIZONA AND CALIFORNIA

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

ALMOND		
Disease Control	Product Rate	Use Information
Brown rot blossom blight <i>(Monilinia laxa)</i> <i>(Monilinia fructicola)</i> Shot hole <i>(Wilsonomyces carpophilus)</i> Powdery mildew <i>(Sphaerotheca pannosa)</i> <i>(Podosphaera tridactyla)</i> Jacket rot Green fruit rot <i>(Botrytis cinerea)</i> Alternaria leaf spot <i>(Alternaria alternata)</i> Anthracnose <i>(Colletotrichum acutatum)</i> Scab <i>(Cladosporium carpophilum)</i> Rust <i>(Tranzschelia discolor)</i>	6.0 to 8.0 fl oz/acre	Begin applications preventatively. Apply on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals. Brown rot Blossom blight: Applications should be made at the critical timings for disease control. Typically this begins at pink bud stage (about 5% bloom). If conditions are conducive for disease development, apply again at full bloom and at petal fall. Refer to University and/or extension guidelines for best application timings.
Disease Suppression	Product Rate	Use Information
Hull rot <i>(Rhizopus spp.)</i> <i>(Monilinia spp.)</i>	6.0 to 17.0 fl oz/acre	For best results apply early hull split and again 14-days later. Refer to University and/or extension guidelines for best application timings.
Restrictions: Do not apply more than 34 fl oz of LUNA [®] EXPERIENCE per acre per season. Do not apply LUNA [®] EXPERIENCE within 35 days of harvest. Do not cut cover crops in treated areas for feed, or allow livestock to graze treated areas. To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA [®] EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.		

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).