

Lamb Milk Replacer

Guaranteed Analysis

Crude Protein	Min	24.0%
Crude Fat	Min	30.0%
Crude Fiber	Max	0.15%
Calcium	Min	0.75%
Calcium	Max	1.25%
Phosphorus	Min	0.70%
Ash	Max	8.0%
Sodium	Max	0.60%
Sodium	Min	1.10%
Selenium	Min	0.30 ppm
Zinc	Min	60 ppm
Vitamin A	Min	27,000 IU/lb
Vitamin D3	Min	5,000 IU/lb
Vitamin E	Min	150 IU/lb

Ingredients

Whey protein concentrate, animal fat (preserved with BHA, BHT, citric acid, and ethoxyquin), dried whey product, dried whey, coconut oil, calcium propionate, citric acid (preservatives), dextrose, dried corn syrup, calcium carbonate, dried skimmed milk, L-lysine, DL-methionine, dicalcium phosphate, sodium silico aluminate, artificial flavor, vitamin E supplement, sodium selenite, maltodextrin, ferrous sulfate, lecithin, ethoxylated mono-diglycerides, propylene glycol, magnesium sulfate, choline chloride, zinc sulfate, vitamin A supplement, manganese sulfate, ascorbic acid, mineral oil, niacin supplement, biotin, vitamin D3 supplement, calcium pantothenate, calcium iodate, vitamin b12 supplement, menadione sodium bisulfite complex (source of vitamin K activity), riboflavin supplement, folic acid, thiamin mononitrate, cobalt sulfate, pyridoxine hydrochloride.

Feeding Directions

Using the enclosed scoop*, mix level measures of Lamb Milk Replacer into 1 pint of warm water. The following amounts of reconstituted Lamb Milk Replacer solution should be fed as indicated below. Amounts are based on an average lamb birth weight of 10 to 12 pounds. Adjust as needed for body weight and appetite.

Age in days	Number of feedings daily	Ounces of Lamb Milk Replacer solution per feeding	Total Ounces of Lamb Milk Replacer solution per day
Day One	4	Colostrum	Colostrum
2-10	3	6-8 ounces	18-24 ounces
11-Weaning	3	8-10 ounces	24-30 ounces

Whenever possible, provide the lamb with at least one feeding of colostrum from the ewe. If the ewe cannot provide colostrum, use [Manna Pro Colostrum Supplement](#). Provide a warm enclosed area for lambs separated from ewes. Feed milk replacer from a SUCKLE bottle using a lamb nipple. Starting the 2nd week feed lamb starter free choice. Sprinkle dry Lamb Milk Replacer over lamb starter to get lambs nibbling on starter feed. Dry feed intake is the best indicator of the appropriate time to wean. Lambs may be weaned when feed intake is greater than 0.5 pounds per day of lamb starter. At weaning, lambs will be at least 30 days old and weigh 25 pounds or more.

*The enclosed scoop holds about 1 ounce of milk replacer powder by weight.